
10 INFINITE SERIES

10.1 Sequences

Preliminary Questions

1. What is a_4 for the sequence $a_n = n^2 - n$?

SOLUTION Substituting $n = 4$ in the expression for a_n gives

$$a_4 = 4^2 - 4 = 12.$$

2. Which of the following sequences converge to zero?

(a) $\frac{n^2}{n^2 + 1}$ (b) 2^n (c) $\left(\frac{-1}{2}\right)^n$

SOLUTION

- (a) This sequence does not converge to zero:

$$\lim_{n \rightarrow \infty} \frac{n^2}{n^2 + 1} = \lim_{x \rightarrow \infty} \frac{x^2}{x^2 + 1} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{x^2}} = \frac{1}{1 + 0} = 1.$$

- (b) This sequence does not converge to zero: this is a geometric sequence with $r = 2 > 1$; hence, the sequence diverges to ∞ .

- (c) Recall that if $|a_n|$ converges to 0, then a_n must also converge to zero. Here,

$$\left| \left(\frac{-1}{2}\right)^n \right| = \left(\frac{1}{2}\right)^n,$$

which is a geometric sequence with $0 < r < 1$; hence, $\left(\frac{1}{2}\right)^n$ converges to zero. It therefore follows that $\left(\frac{-1}{2}\right)^n$ converges to zero.

3. Let a_n be the n th decimal approximation to $\sqrt{2}$. That is, $a_1 = 1$, $a_2 = 1.4$, $a_3 = 1.41$, etc. What is $\lim_{n \rightarrow \infty} a_n$?

SOLUTION $\lim_{n \rightarrow \infty} a_n = \sqrt{2}$.

4. Which of the following sequences is defined recursively?

(a) $a_n = \sqrt{4 + n}$ (b) $b_n = \sqrt{4 + b_{n-1}}$

SOLUTION

- (a) a_n can be computed directly, since it depends on n only and not on preceding terms. Therefore a_n is defined explicitly and not recursively.

- (b) b_n is computed in terms of the preceding term b_{n-1} , hence the sequence $\{b_n\}$ is defined recursively.

5. Theorem 5 says that every convergent sequence is bounded. Determine if the following statements are true or false and if false, give a counterexample.

- (a) If $\{a_n\}$ is bounded, then it converges.
(b) If $\{a_n\}$ is not bounded, then it diverges.
(c) If $\{a_n\}$ diverges, then it is not bounded.

SOLUTION

- (a) This statement is false. The sequence $a_n = \cos \pi n$ is bounded since $-1 \leq \cos \pi n \leq 1$ for all n , but it does not converge: since $a_n = \cos \pi n = (-1)^n$, the terms assume the two values 1 and -1 alternately, hence they do not approach one value.

- (b) By Theorem 5, a converging sequence must be bounded. Therefore, if a sequence is not bounded, it certainly does not converge.

- (c) The statement is false. The sequence $a_n = (-1)^n$ is bounded, but it does not approach one limit.

Exercises

1. Match each sequence with its general term:

$a_1, a_2, a_3, a_4, \dots$	General term
(a) $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \dots$	(i) $\cos \pi n$
(b) $-1, 1, -1, 1, \dots$	(ii) $\frac{n!}{2^n}$
(c) $1, -1, 1, -1, \dots$	(iii) $(-1)^{n+1}$
(d) $\frac{1}{2}, \frac{2}{4}, \frac{6}{8}, \frac{24}{16}, \dots$	(iv) $\frac{n}{n+1}$

SOLUTION

(a) The numerator of each term is the same as the index of the term, and the denominator is one more than the numerator; hence $a_n = \frac{n}{n+1}, n = 1, 2, 3, \dots$

(b) The terms of this sequence are alternating between -1 and 1 so that the positive terms are in the even places. Since $\cos \pi n = 1$ for even n and $\cos \pi n = -1$ for odd n , we have $a_n = \cos \pi n, n = 1, 2, \dots$

(c) The terms a_n are 1 for odd n and -1 for even n . Hence, $a_n = (-1)^{n+1}, n = 1, 2, \dots$

(d) The numerator of each term is $n!$, and the denominator is 2^n ; hence, $a_n = \frac{n!}{2^n}, n = 1, 2, 3, \dots$

2. Let $a_n = \frac{1}{2n-1}$ for $n = 1, 2, 3, \dots$. Write out the first three terms of the following sequences.

(a) $b_n = a_{n+1}$ (b) $c_n = a_{n+3}$ (c) $d_n = a_n^2$ (d) $e_n = 2a_n - a_{n+1}$ **SOLUTION**(a) The first three terms of $\{b_n\}$ are:

$$b_1 = a_2 = \frac{1}{2 \cdot 2 - 1} = \frac{1}{3}, \quad b_2 = a_3 = \frac{1}{2 \cdot 3 - 1} = \frac{1}{5}, \quad b_3 = a_4 = \frac{1}{2 \cdot 4 - 1} = \frac{1}{7}.$$

(b) The first three terms of $\{c_n\}$ are:

$$c_1 = a_4 = \frac{1}{2 \cdot 4 - 1} = \frac{1}{7}, \quad c_2 = a_5 = \frac{1}{2 \cdot 5 - 1} = \frac{1}{9}, \quad c_3 = a_6 = \frac{1}{2 \cdot 6 - 1} = \frac{1}{11}.$$

(c) Note

$$a_1 = \frac{1}{2 \cdot 1 - 1} = 1, \quad a_2 = \frac{1}{2 \cdot 2 - 1} = \frac{1}{3}, \quad a_3 = \frac{1}{2 \cdot 3 - 1} = \frac{1}{5}.$$

Thus,

$$d_1 = a_1^2 = 1^2 = 1, \quad d_2 = a_2^2 = \left(\frac{1}{3}\right)^2 = \frac{1}{9}, \quad d_3 = a_3^2 = \left(\frac{1}{5}\right)^2 = \frac{1}{25}.$$

(d) The first three terms of $\{e_n\}$ are:

$$e_1 = 2a_1 - a_2, \quad e_2 = 2a_2 - a_3, \quad e_3 = 2a_3 - a_4.$$

Thus, we must compute a_1, a_2, a_3 and a_4 . We set $n = 1, 2, 3$ and 4 in the formula for a_n to obtain:

$$a_1 = \frac{1}{2 \cdot 1 - 1} = 1, \quad a_2 = \frac{1}{2 \cdot 2 - 1} = \frac{1}{3}, \quad a_3 = \frac{1}{2 \cdot 3 - 1} = \frac{1}{5}, \quad a_4 = \frac{1}{2 \cdot 4 - 1} = \frac{1}{7}.$$

Therefore,

$$e_1 = 2 \cdot 1 - \frac{1}{3} = \frac{5}{3}, \quad e_2 = 2 \cdot \frac{1}{3} - \frac{1}{5} = \frac{7}{15}, \quad e_3 = 2 \cdot \frac{1}{5} - \frac{1}{7} = \frac{9}{35}.$$

In Exercises 3–12, calculate the first four terms of the sequence, starting with $n = 1$.

3. $c_n = \frac{3^n}{n!}$

SOLUTION Setting $n = 1, 2, 3, 4$ in the formula for c_n gives

$$c_1 = \frac{3^1}{1!} = \frac{3}{1} = 3, \quad c_2 = \frac{3^2}{2!} = \frac{9}{2},$$

$$c_3 = \frac{3^3}{3!} = \frac{27}{6} = \frac{9}{2}, \quad c_4 = \frac{3^4}{4!} = \frac{81}{24} = \frac{27}{8}.$$

$$4. b_n = \frac{(2n-1)!}{n!}$$

SOLUTION Setting $n = 1, 2, 3, 4$ in the formula for b_n gives

$$b_1 = \frac{(2 \cdot 1 - 1)!}{1!} = \frac{1}{1} = 1, \quad b_2 = \frac{(2 \cdot 2 - 1)!}{2!} = \frac{6}{2} = 3,$$

$$b_3 = \frac{(2 \cdot 3 - 1)!}{3!} = \frac{120}{6} = 20, \quad b_4 = \frac{(2 \cdot 4 - 1)!}{4!} = \frac{5040}{24} = 210.$$

$$5. a_1 = 2, \quad a_{n+1} = 2a_n^2 - 3$$

SOLUTION For $n = 1, 2, 3$ we have:

$$a_2 = a_{1+1} = 2a_1^2 - 3 = 2 \cdot 4 - 3 = 5;$$

$$a_3 = a_{2+1} = 2a_2^2 - 3 = 2 \cdot 25 - 3 = 47;$$

$$a_4 = a_{3+1} = 2a_3^2 - 3 = 2 \cdot 2209 - 3 = 4415.$$

The first four terms of $\{a_n\}$ are 2, 5, 47, 4415.

$$6. b_1 = 1, \quad b_n = b_{n-1} + \frac{1}{b_{n-1}}$$

SOLUTION For $n = 2, 3, 4$ we have

$$b_2 = b_1 + \frac{1}{b_1} = 1 + \frac{1}{1} = 2;$$

$$b_3 = b_2 + \frac{1}{b_2} = 2 + \frac{1}{2} = \frac{5}{2};$$

$$b_4 = b_3 + \frac{1}{b_3} = \frac{5}{2} + \frac{2}{5} = \frac{29}{10}.$$

The first four terms of $\{b_n\}$ are 1, 2, $\frac{5}{2}$, $\frac{29}{10}$.

$$7. b_n = 5 + \cos \pi n$$

SOLUTION For $n = 1, 2, 3, 4$ we have

$$b_1 = 5 + \cos \pi = 4;$$

$$b_2 = 5 + \cos 2\pi = 6;$$

$$b_3 = 5 + \cos 3\pi = 4;$$

$$b_4 = 5 + \cos 4\pi = 6.$$

The first four terms of $\{b_n\}$ are 4, 6, 4, 6.

$$8. c_n = (-1)^{2n+1}$$

SOLUTION for $n = 1, 2, 3, 4$ we have

$$c_1 = (-1)^{2 \cdot 1 + 1} = (-1)^3 = -1;$$

$$c_2 = (-1)^{2 \cdot 2 + 1} = (-1)^5 = -1;$$

$$c_3 = (-1)^{2 \cdot 3 + 1} = (-1)^7 = -1;$$

$$c_4 = (-1)^{2 \cdot 4 + 1} = (-1)^9 = -1.$$

The first four terms of $\{c_n\}$ are -1, -1, -1, -1.

$$9. c_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$$

SOLUTION

$$c_1 = 1;$$

$$c_2 = 1 + \frac{1}{2} = \frac{3}{2};$$

$$c_3 = 1 + \frac{1}{2} + \frac{1}{3} = \frac{3}{2} + \frac{1}{3} = \frac{11}{6};$$

$$c_4 = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} = \frac{11}{6} + \frac{1}{4} = \frac{25}{12}.$$

10. $a_n = n + (n + 1) + (n + 2) + \cdots + (2n)$

SOLUTION The general term a_n is the sum of $n + 1$ successive numbers, where the first one is n and the last one is $2n$. Thus,

$$a_1 = 1 + 2 = 3;$$

$$a_2 = 2 + 3 + 4 = 9;$$

$$a_3 = 3 + 4 + 5 + 6 = 18;$$

$$a_4 = 4 + 5 + 6 + 7 + 8 = 30.$$

11. $b_1 = 2$, $b_2 = 3$, $b_n = 2b_{n-1} + b_{n-2}$

SOLUTION We need to find b_3 and b_4 . Setting $n = 3$ and $n = 4$ and using the given values for b_1 and b_2 we obtain:

$$b_3 = 2b_{3-1} + b_{3-2} = 2b_2 + b_1 = 2 \cdot 3 + 2 = 8;$$

$$b_4 = 2b_{4-1} + b_{4-2} = 2b_3 + b_2 = 2 \cdot 8 + 3 = 19.$$

The first four terms of the sequence $\{b_n\}$ are 2, 3, 8, 19.

12. $c_n = n$ -place decimal approximation to e

SOLUTION Using a calculator we find that $e = 2.718281828 \dots$. Thus, the four first terms of $\{c_n\}$ are

$$c_1 = 2.7; \quad c_2 = 2.72; \quad c_3 = 2.718; \quad c_4 = 2.7183.$$

13. Find a formula for the n th term of each sequence.

(a) $\frac{1}{1}, \frac{-1}{8}, \frac{1}{27}, \dots$

(b) $\frac{2}{6}, \frac{3}{7}, \frac{4}{8}, \dots$

SOLUTION

(a) The denominators are the third powers of the positive integers starting with $n = 1$. Also, the sign of the terms is alternating with the sign of the first term being positive. Thus,

$$a_1 = \frac{1}{1^3} = \frac{(-1)^{1+1}}{1^3}; \quad a_2 = -\frac{1}{2^3} = \frac{(-1)^{2+1}}{2^3}; \quad a_3 = \frac{1}{3^3} = \frac{(-1)^{3+1}}{3^3}.$$

This rule leads to the following formula for the n th term:

$$a_n = \frac{(-1)^{n+1}}{n^3}.$$

(b) Assuming a starting index of $n = 1$, we see that each numerator is one more than the index and the denominator is four more than the numerator. Thus, the general term a_n is

$$a_n = \frac{n + 1}{n + 5}.$$

14. Suppose that $\lim_{n \rightarrow \infty} a_n = 4$ and $\lim_{n \rightarrow \infty} b_n = 7$. Determine:

(a) $\lim_{n \rightarrow \infty} (a_n + b_n)$

(b) $\lim_{n \rightarrow \infty} a_n^3$

(c) $\lim_{n \rightarrow \infty} \cos(\pi b_n)$

(d) $\lim_{n \rightarrow \infty} (a_n^2 - 2a_n b_n)$

SOLUTION

(a) By the Limit Laws for Sequences, we find

$$\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n = 4 + 7 = 11.$$

(b) By the Limit Laws for Sequences, we find

$$\lim_{n \rightarrow \infty} a_n^3 = \lim_{n \rightarrow \infty} (a_n \cdot a_n \cdot a_n) = \left(\lim_{n \rightarrow \infty} a_n \right) \cdot \left(\lim_{n \rightarrow \infty} a_n \right) \cdot \left(\lim_{n \rightarrow \infty} a_n \right) = \left(\lim_{n \rightarrow \infty} a_n \right)^3 = 4^3 = 64.$$

(c) By Theorem 4, we can “bring the limit inside the function”:

$$\lim_{n \rightarrow \infty} \cos(\pi b_n) = \cos \left(\lim_{n \rightarrow \infty} \pi b_n \right) = \cos \left(\pi \lim_{n \rightarrow \infty} b_n \right) = \cos(7\pi) = -1.$$

(d) By the Limit Laws of Sequences, we find

$$\lim_{n \rightarrow \infty} (a_n^2 - 2a_n b_n) = \lim_{n \rightarrow \infty} a_n^2 - \lim_{n \rightarrow \infty} 2a_n b_n = \left(\lim_{n \rightarrow \infty} a_n \right)^2 - 2 \left(\lim_{n \rightarrow \infty} a_n \right) \left(\lim_{n \rightarrow \infty} b_n \right) = 4^2 - 2 \cdot 4 \cdot 7 = -40.$$

In Exercises 15–26, use Theorem 1 to determine the limit of the sequence or state that the sequence diverges.

15. $a_n = 12$

SOLUTION We have $a_n = f(n)$ where $f(x) = 12$; thus,

$$\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} 12 = 12.$$

16. $a_n = 20 - \frac{4}{n^2}$

SOLUTION We have $a_n = f(n)$ where $f(x) = 20 - \frac{4}{x^2}$; thus,

$$\lim_{n \rightarrow \infty} \left(20 - \frac{4}{n^2} \right) = \lim_{x \rightarrow \infty} \left(20 - \frac{4}{x^2} \right) = 20 - 0 = 20.$$

17. $b_n = \frac{5n-1}{12n+9}$

SOLUTION We have $b_n = f(n)$ where $f(x) = \frac{5x-1}{12x+9}$; thus,

$$\lim_{n \rightarrow \infty} \frac{5n-1}{12n+9} = \lim_{x \rightarrow \infty} \frac{5x-1}{12x+9} = \frac{5}{12}.$$

18. $a_n = \frac{4+n-3n^2}{4n^2+1}$

SOLUTION We have $a_n = f(n)$ where $f(x) = \frac{4+x-3x^2}{4x^2+1}$; thus,

$$\lim_{n \rightarrow \infty} \frac{4+n-3n^2}{4n^2+1} = \lim_{x \rightarrow \infty} \frac{4+x-3x^2}{4x^2+1} = -\frac{3}{4}$$

19. $c_n = -2^{-n}$

SOLUTION We have $c_n = f(n)$ where $f(x) = -2^{-x}$; thus,

$$\lim_{n \rightarrow \infty} (-2^{-n}) = \lim_{x \rightarrow \infty} -2^{-x} = \lim_{x \rightarrow \infty} -\frac{1}{2^x} = 0.$$

20. $z_n = \left(\frac{1}{3}\right)^n$

SOLUTION We have $z_n = f(n)$ where $f(x) = \left(\frac{1}{3}\right)^x$; thus,

$$\lim_{n \rightarrow \infty} \left(\frac{1}{3}\right)^n = \lim_{x \rightarrow \infty} \left(\frac{1}{3}\right)^x = 0.$$

21. $c_n = 9^n$

SOLUTION We have $c_n = f(n)$ where $f(x) = 9^x$; thus,

$$\lim_{n \rightarrow \infty} 9^n = \lim_{x \rightarrow \infty} 9^x = \infty$$

Thus, the sequence 9^n diverges.

22. $z_n = 10^{-1/n}$

SOLUTION We have $z_n = f(n)$ where $f(x) = (0.1)^{-1/x}$; thus

$$\lim_{n \rightarrow \infty} (0.1)^{-1/n} = \lim_{x \rightarrow \infty} (0.1)^{-1/x} = (0.1)^{\lim_{x \rightarrow \infty} (-1/x)} = (0.1)^0 = 1.$$

23. $a_n = \frac{n}{\sqrt{n^2+1}}$

SOLUTION We have $a_n = f(n)$ where $f(x) = \frac{x}{\sqrt{x^2+1}}$; thus,

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2+1}} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2+1}} = \lim_{x \rightarrow \infty} \frac{\frac{x}{x}}{\frac{\sqrt{x^2+1}}{x}} = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{\frac{x^2+1}{x^2}}} = \lim_{x \rightarrow \infty} \frac{1}{\sqrt{1+\frac{1}{x^2}}} = \frac{1}{\sqrt{1+0}} = 1.$$

$$24. a_n = \frac{n}{\sqrt{n^3 + 1}}$$

SOLUTION We have $a_n = f(n)$ where $f(x) = \frac{x}{\sqrt{x^3 + 1}}$; thus,

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^3 + 1}} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^3 + 1}} = \lim_{x \rightarrow \infty} \frac{\frac{x}{x^{3/2}}}{\frac{\sqrt{x^3 + 1}}{x^{3/2}}} = \lim_{x \rightarrow \infty} \frac{\frac{1}{\sqrt{x}}}{\sqrt{1 + \frac{1}{x^3}}} = \frac{0}{\sqrt{1 + 0}} = \frac{0}{1} = 0.$$

$$25. a_n = \ln \left(\frac{12n + 2}{-9 + 4n} \right)$$

SOLUTION We have $a_n = f(n)$ where $f(x) = \ln \left(\frac{12x + 2}{-9 + 4x} \right)$; thus,

$$\lim_{n \rightarrow \infty} \ln \left(\frac{12n + 2}{-9 + 4n} \right) = \lim_{x \rightarrow \infty} \ln \left(\frac{12x + 2}{-9 + 4x} \right) = \ln \lim_{x \rightarrow \infty} \left(\frac{12x + 2}{-9 + 4x} \right) = \ln 3$$

$$26. r_n = \ln n - \ln(n^2 + 1)$$

SOLUTION We have $r_n = f(n)$ where $f(x) = \ln x - \ln(x^2 + 1)$; thus,

$$\lim_{n \rightarrow \infty} (\ln n - \ln(n^2 + 1)) = \lim_{x \rightarrow \infty} (\ln x - \ln(x^2 + 1)) = \lim_{x \rightarrow \infty} \ln \frac{x}{x^2 + 1}$$

But this function diverges as $x \rightarrow \infty$, so that r_n diverges as well.

In Exercises 27–30, use Theorem 4 to determine the limit of the sequence.

$$27. a_n = \sqrt{4 + \frac{1}{n}}$$

SOLUTION We have

$$\lim_{n \rightarrow \infty} 4 + \frac{1}{n} = \lim_{x \rightarrow \infty} 4 + \frac{1}{x} = 4$$

Since \sqrt{x} is a continuous function for $x > 0$, Theorem 4 tells us that

$$\lim_{n \rightarrow \infty} \sqrt{4 + \frac{1}{n}} = \sqrt{\lim_{n \rightarrow \infty} 4 + \frac{1}{n}} = \sqrt{4} = 2$$

$$28. a_n = e^{4n/(3n+9)}$$

SOLUTION We have

$$\lim_{n \rightarrow \infty} \frac{4n}{3n + 9} = \frac{4}{3}$$

Since e^x is continuous for all x , Theorem 4 tells us that

$$\lim_{n \rightarrow \infty} e^{4n/(3n+9)} = e^{\lim_{n \rightarrow \infty} 4n/(3n+9)} = e^{4/3}$$

$$29. a_n = \cos^{-1} \left(\frac{n^3}{2n^3 + 1} \right)$$

SOLUTION We have

$$\lim_{n \rightarrow \infty} \frac{n^3}{2n^3 + 1} = \frac{1}{2}$$

Since $\cos^{-1}(x)$ is continuous for all x , Theorem 4 tells us that

$$\lim_{n \rightarrow \infty} \cos^{-1} \left(\frac{n^3}{2n^3 + 1} \right) = \cos^{-1} \left(\lim_{n \rightarrow \infty} \frac{n^3}{2n^3 + 1} \right) = \cos^{-1}(1/2) = \frac{\pi}{3}$$

$$30. a_n = \tan^{-1}(e^{-n})$$

SOLUTION We have

$$\lim_{n \rightarrow \infty} e^{-n} = e^{-\infty} = \lim_{x \rightarrow \infty} e^{-x} = 0$$

Since $\tan^{-1}(x)$ is continuous for all x , Theorem 4 tells us that

$$\lim_{n \rightarrow \infty} \tan^{-1}(e^{-n}) = \tan^{-1} \left(\lim_{n \rightarrow \infty} e^{-n} \right) = \tan^{-1}(0) = 0$$

31. Let $a_n = \frac{n}{n+1}$. Find a number M such that:

(a) $|a_n - 1| \leq 0.001$ for $n \geq M$.

(b) $|a_n - 1| \leq 0.00001$ for $n \geq M$.

Then use the limit definition to prove that $\lim_{n \rightarrow \infty} a_n = 1$.

SOLUTION

(a) We have

$$|a_n - 1| = \left| \frac{n}{n+1} - 1 \right| = \left| \frac{n - (n+1)}{n+1} \right| = \left| \frac{-1}{n+1} \right| = \frac{1}{n+1}.$$

Therefore $|a_n - 1| \leq 0.001$ provided $\frac{1}{n+1} \leq 0.001$, that is, $n \geq 999$. It follows that we can take $M = 999$.

(b) By part (a), $|a_n - 1| \leq 0.00001$ provided $\frac{1}{n+1} \leq 0.00001$, that is, $n \geq 99999$. It follows that we can take $M = 99999$.

We now prove formally that $\lim_{n \rightarrow \infty} a_n = 1$. Using part (a), we know that

$$|a_n - 1| = \frac{1}{n+1} < \epsilon,$$

provided $n > \frac{1}{\epsilon} - 1$. Thus, Let $\epsilon > 0$ and take $M = \frac{1}{\epsilon} - 1$. Then, for $n > M$, we have

$$|a_n - 1| = \frac{1}{n+1} < \frac{1}{M+1} = \epsilon.$$

32. Let $b_n = \left(\frac{1}{3}\right)^n$.

(a) Find a value of M such that $|b_n| \leq 10^{-5}$ for $n \geq M$.

(b) Use the limit definition to prove that $\lim_{n \rightarrow \infty} b_n = 0$.

SOLUTION

(a) Solving $\left(\frac{1}{3}\right)^n \leq 10^{-5}$ for n , we find

$$n \geq 5 \log_3 10 = 5 \frac{\ln 10}{\ln 3} \approx 10.48.$$

It follows that we can take $M = 10.5$.

(b) We see that

$$\left| \left(\frac{1}{3}\right)^n - 0 \right| = \frac{1}{3^n} < \epsilon$$

provided

$$n > \log_3 \frac{1}{\epsilon}.$$

Thus, let $\epsilon > 0$ and take $M = \log_3 \frac{1}{\epsilon}$. Then, for $n > M$, we have

$$\left| \left(\frac{1}{3}\right)^n - 0 \right| = \frac{1}{3^n} < \frac{1}{3^M} = \epsilon.$$

33. Use the limit definition to prove that $\lim_{n \rightarrow \infty} n^{-2} = 0$.

SOLUTION We see that

$$|n^{-2} - 0| = \left| \frac{1}{n^2} \right| = \frac{1}{n^2} < \epsilon$$

provided

$$n > \frac{1}{\sqrt{\epsilon}}.$$

Thus, let $\epsilon > 0$ and take $M = \frac{1}{\sqrt{\epsilon}}$. Then, for $n > M$, we have

$$|n^{-2} - 0| = \left| \frac{1}{n^2} \right| = \frac{1}{n^2} < \frac{1}{M^2} = \epsilon.$$

34. Use the limit definition to prove that $\lim_{n \rightarrow \infty} \frac{n}{n + n^{-1}} = 1$.

SOLUTION Since

$$\frac{n}{n + n^{-1}} = \frac{n^2}{n(n + n^{-1})} = \frac{n^2}{n^2 + 1}$$

we see that

$$\left| \frac{n^2}{n^2 + 1} - 1 \right| = \left| \frac{-1}{n^2 + 1} \right| = \frac{1}{n^2 + 1} < \epsilon$$

provided

$$n > \sqrt{\frac{1}{\epsilon} - 1}$$

So choose $\epsilon > 0$, and let $M = \sqrt{\frac{1}{\epsilon} - 1}$. Then, for $n > M$, we have

$$\left| \frac{n}{n + n^{-1}} - 1 \right| = \left| \frac{-1}{n^2 + 1} \right| = \frac{1}{n^2 + 1} < \frac{1}{(\frac{1}{\epsilon} - 1) + 1} = \epsilon$$

In Exercises 35–62, use the appropriate limit laws and theorems to determine the limit of the sequence or show that it diverges.

35. $a_n = 10 + \left(-\frac{1}{9}\right)^n$

SOLUTION By the Limit Laws for Sequences we have:

$$\lim_{n \rightarrow \infty} \left(10 + \left(-\frac{1}{9}\right)^n\right) = \lim_{n \rightarrow \infty} 10 + \lim_{n \rightarrow \infty} \left(-\frac{1}{9}\right)^n = 10 + \lim_{n \rightarrow \infty} \left(-\frac{1}{9}\right)^n.$$

Now,

$$-\left(\frac{1}{9}\right)^n \leq \left(-\frac{1}{9}\right)^n \leq \left(\frac{1}{9}\right)^n.$$

Because

$$\lim_{n \rightarrow \infty} \left(\frac{1}{9}\right)^n = 0,$$

by the Limit Laws for Sequences,

$$\lim_{n \rightarrow \infty} -\left(\frac{1}{9}\right)^n = -\lim_{n \rightarrow \infty} \left(\frac{1}{9}\right)^n = 0.$$

Thus, we have

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{9}\right)^n = 0,$$

and

$$\lim_{n \rightarrow \infty} \left(10 + \left(-\frac{1}{9}\right)^n\right) = 10 + 0 = 10.$$

36. $d_n = \sqrt{n+3} - \sqrt{n}$

SOLUTION We multiply and divide d_n by the conjugate expression $\sqrt{n+3} + \sqrt{n}$ and use the identity $(a-b)(a+b) = a^2 - b^2$ to obtain:

$$d_n = \frac{(\sqrt{n+3} - \sqrt{n})(\sqrt{n+3} + \sqrt{n})}{\sqrt{n+3} + \sqrt{n}} = \frac{(n+3) - n}{\sqrt{n+3} + \sqrt{n}} = \frac{3}{\sqrt{n+3} + \sqrt{n}}.$$

Thus,

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} \frac{3}{\sqrt{n+3} + \sqrt{n}} = \lim_{x \rightarrow \infty} \frac{3}{\sqrt{x+3} + \sqrt{x}} = 0.$$

37. $c_n = 1.01^n$

SOLUTION Since $c_n = f(n)$ where $f(x) = 1.01^x$, we have

$$\lim_{n \rightarrow \infty} 1.01^n = \lim_{x \rightarrow \infty} 1.01^x = \infty$$

so that the sequence diverges.

38. $b_n = e^{1-n^2}$

SOLUTION Since $b_n = f(n)$ where $f(x) = e^{1-x^2}$, we have

$$\lim_{n \rightarrow \infty} e^{1-n^2} = \lim_{x \rightarrow \infty} e^{1-x^2} = \lim_{x \rightarrow \infty} \frac{e}{e^{x^2}} = 0$$

39. $a_n = 2^{1/n}$

SOLUTION Because 2^x is a continuous function,

$$\lim_{n \rightarrow \infty} 2^{1/n} = \lim_{x \rightarrow \infty} 2^{1/x} = 2^{\lim_{x \rightarrow \infty} (1/x)} = 2^0 = 1.$$

40. $b_n = n^{1/n}$

SOLUTION Let $b_n = n^{1/n}$. Take the natural logarithm of both sides of this expression to obtain

$$\ln b_n = \ln n^{1/n} = \frac{\ln n}{n}.$$

Thus,

$$\lim_{n \rightarrow \infty} (\ln b_n) = \lim_{n \rightarrow \infty} \frac{\ln n}{n} = \lim_{x \rightarrow \infty} \frac{\ln x}{x} = \lim_{x \rightarrow \infty} \frac{1}{x} = 0.$$

Because $f(x) = e^x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} e^{\ln b_n} = e^{\lim_{n \rightarrow \infty} (\ln b_n)} = e^0 = 1.$$

That is,

$$\lim_{n \rightarrow \infty} n^{1/n} = 1.$$

41. $c_n = \frac{9^n}{n!}$

SOLUTION For $n \geq 9$, write

$$c_n = \frac{9^n}{n!} = \frac{9}{1} \cdot \frac{9}{2} \cdots \frac{9}{9} \cdot \frac{9}{10} \cdot \frac{9}{11} \cdots \frac{9}{n-1} \cdot \frac{9}{n}$$

call this C Each factor is less than 1

Then clearly

$$0 \leq \frac{9^n}{n!} \leq C \frac{9}{n}$$

since each factor after the first nine is < 1 . The squeeze theorem tells us that

$$\lim_{n \rightarrow \infty} 0 \leq \lim_{n \rightarrow \infty} \frac{9^n}{n!} \leq \lim_{n \rightarrow \infty} C \frac{9}{n} = C \lim_{n \rightarrow \infty} \frac{9}{n} = C \cdot 0 = 0$$

so that $\lim_{n \rightarrow \infty} c_n = 0$ as well.

42. $a_n = \frac{8^{2n}}{n!}$

SOLUTION Note that

$$a_n = \frac{8^{2n}}{n!} = \frac{64^n}{n!}$$

Now apply the same method as in the Exercise 41. For $n \geq 64$, write

$$c_n = \frac{64^n}{n!} = \frac{64}{1} \cdot \frac{64}{2} \cdots \frac{64}{64} \cdot \frac{64}{65} \cdot \frac{64}{66} \cdots \frac{64}{n-1} \cdot \frac{64}{n}$$

call this C Each factor is less than 1

Then clearly

$$0 \leq \frac{64^n}{n!} \leq C \frac{64}{n}$$

since each factor after the first 64 is < 1 . The squeeze theorem tells us that

$$\lim_{n \rightarrow \infty} 0 \leq \lim_{n \rightarrow \infty} \frac{64^n}{n!} \leq \lim_{n \rightarrow \infty} C \frac{64}{n} = C \lim_{n \rightarrow \infty} \frac{64}{n} = C \cdot 0 = 0$$

so that $\lim_{n \rightarrow \infty} a_n = 0$ as well.

$$43. a_n = \frac{3n^2 + n + 2}{2n^2 - 3}$$

SOLUTION

$$\lim_{n \rightarrow \infty} \frac{3n^2 + n + 2}{2n^2 - 3} = \lim_{x \rightarrow \infty} \frac{3x^2 + x + 2}{2x^2 - 3} = \frac{3}{2}.$$

$$44. a_n = \frac{\sqrt{n}}{\sqrt{n} + 4}$$

SOLUTION

$$\lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n} + 4} = \lim_{x \rightarrow \infty} \frac{\sqrt{x}}{\sqrt{x} + 4} = \lim_{x \rightarrow \infty} \frac{\frac{\sqrt{x}}{\sqrt{x}}}{\frac{\sqrt{x}}{\sqrt{x}} + \frac{4}{\sqrt{x}}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{4}{\sqrt{x}}} = \frac{1}{1 + 0} = 1.$$

$$45. a_n = \frac{\cos n}{n}$$

SOLUTION Since $-1 \leq \cos n \leq 1$ the following holds:

$$-\frac{1}{n} \leq \frac{\cos n}{n} \leq \frac{1}{n}.$$

We now apply the Squeeze Theorem for Sequences and the limits

$$\lim_{n \rightarrow \infty} -\frac{1}{n} = \lim_{n \rightarrow \infty} \frac{1}{n} = 0$$

to conclude that $\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0$.

$$46. c_n = \frac{(-1)^n}{\sqrt{n}}$$

SOLUTION Clearly

$$-\frac{1}{\sqrt{n}} \leq \frac{(-1)^n}{\sqrt{n}} \leq \frac{1}{\sqrt{n}}$$

Since

$$\lim_{n \rightarrow \infty} \frac{-1}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0,$$

the Squeeze Theorem tells us that $\lim_{n \rightarrow \infty} \frac{(-1)^n}{\sqrt{n}} = 0$.

$$47. d_n = \ln 5^n - \ln n!$$

SOLUTION Note that

$$d_n = \ln \frac{5^n}{n!}$$

so that

$$e^{d_n} = \frac{5^n}{n!} \quad \text{so} \quad \lim_{n \rightarrow \infty} e^{d_n} = \lim_{n \rightarrow \infty} \frac{5^n}{n!} = 0$$

by the method of Exercise 41. If d_n converged, we could, since $f(x) = e^x$ is continuous, then write

$$\lim_{n \rightarrow \infty} e^{d_n} = e^{\lim_{n \rightarrow \infty} d_n} = 0$$

which is impossible. Thus $\{d_n\}$ diverges.

$$48. d_n = \ln(n^2 + 4) - \ln(n^2 - 1)$$

SOLUTION Note that

$$d_n = \ln \frac{n^2 + 4}{n^2 - 1}$$

so exponentiating both sides of this expression gives

$$e^{d_n} = \frac{n^2 + 4}{n^2 - 1} = \frac{1 + (4/n^2)}{1 - (1/n^2)}$$

Thus,

$$\lim_{n \rightarrow \infty} e^{d_n} = \lim_{n \rightarrow \infty} \frac{1 + (4/n^2)}{1 - (1/n^2)} = 1$$

Because $f(x) = \ln x$ is continuous for $x > 0$, it follows that

$$\lim_{n \rightarrow \infty} d_n = \lim_{n \rightarrow \infty} \ln(e^{d_n}) = \ln(\lim_{n \rightarrow \infty} e^{d_n}) = \ln 1 = 0$$

$$49. a_n = \left(2 + \frac{4}{n^2}\right)^{1/3}$$

SOLUTION Let $a_n = \left(2 + \frac{4}{n^2}\right)^{1/3}$. Taking the natural logarithm of both sides of this expression yields

$$\ln a_n = \ln \left(2 + \frac{4}{n^2}\right)^{1/3} = \frac{1}{3} \ln \left(2 + \frac{4}{n^2}\right).$$

Thus,

$$\begin{aligned} \lim_{n \rightarrow \infty} \ln a_n &= \lim_{n \rightarrow \infty} \frac{1}{3} \ln \left(2 + \frac{4}{n^2}\right) = \frac{1}{3} \lim_{x \rightarrow \infty} \ln \left(2 + \frac{4}{x^2}\right) = \frac{1}{3} \ln \left(\lim_{x \rightarrow \infty} \left(2 + \frac{4}{x^2}\right)\right) \\ &= \frac{1}{3} \ln(2 + 0) = \frac{1}{3} \ln 2 = \ln 2^{1/3}. \end{aligned}$$

Because $f(x) = e^x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} e^{\ln a_n} = e^{\lim_{n \rightarrow \infty} (\ln a_n)} = e^{\ln 2^{1/3}} = 2^{1/3}.$$

$$50. b_n = \tan^{-1} \left(1 - \frac{2}{n}\right)$$

SOLUTION Because $f(x) = \tan^{-1} x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} b_n = \lim_{x \rightarrow \infty} \tan^{-1} \left(1 - \frac{2}{x}\right) = \tan^{-1} \left(\lim_{x \rightarrow \infty} \left(1 - \frac{2}{x}\right)\right) = \tan^{-1} 1 = \frac{\pi}{4}.$$

$$51. c_n = \ln \left(\frac{2n+1}{3n+4}\right)$$

SOLUTION Because $f(x) = \ln x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} c_n = \lim_{x \rightarrow \infty} \ln \left(\frac{2x+1}{3x+4}\right) = \ln \left(\lim_{x \rightarrow \infty} \frac{2x+1}{3x+4}\right) = \ln \frac{2}{3}.$$

$$52. c_n = \frac{n}{n + n^{1/n}}$$

SOLUTION We rewrite $\frac{n}{n + n^{1/n}}$ as follows:

$$\frac{n}{n + n^{1/n}} = \frac{\frac{n}{n}}{\frac{n}{n} + \frac{n^{1/n}}{n}} = \frac{1}{1 + \frac{n^{1/n}}{n}}.$$

Now,

$$\frac{n^{1/n}}{n} = n^{\frac{1}{n}-1} = \frac{1}{n^{1-1/n}},$$

and

$$\lim_{n \rightarrow \infty} \frac{n^{1/n}}{n} = \lim_{n \rightarrow \infty} \frac{1}{n^{1-1/n}} = \lim_{x \rightarrow \infty} \frac{1}{x^{1-1/x}} = 0.$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{n}{n + n^{1/n}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{x^{1/x}}{x}} = \frac{\lim_{x \rightarrow \infty} 1}{\lim_{x \rightarrow \infty} \left(1 + \frac{x^{1/x}}{x}\right)} = \frac{\lim_{x \rightarrow \infty} 1}{\lim_{x \rightarrow \infty} 1 + \lim_{x \rightarrow \infty} \frac{x^{1/x}}{x}} = \frac{1}{1 + 0} = 1.$$

$$53. y_n = \frac{e^n}{2^n}$$

SOLUTION $\frac{e^n}{2^n} = \left(\frac{e}{2}\right)^n$ and $\frac{e}{2} > 1$. By the Limit of Geometric Sequences, we conclude that $\lim_{n \rightarrow \infty} \left(\frac{e}{2}\right)^n = \infty$. Thus, the given sequence diverges.

$$54. a_n = \frac{n}{2^n}$$

SOLUTION

$$\lim_{n \rightarrow \infty} \frac{n}{2^n} = \lim_{x \rightarrow \infty} \frac{x}{2^x} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(x)}{\frac{d}{dx}(2^x)} = \lim_{x \rightarrow \infty} \frac{1}{(\ln 2) 2^x} = \frac{1}{\ln 2} \lim_{x \rightarrow \infty} \frac{1}{2^x} = \frac{1}{\ln 2} \cdot 0 = 0.$$

$$55. y_n = \frac{e^n + (-3)^n}{5^n}$$

SOLUTION

$$\lim_{n \rightarrow \infty} \frac{e^n + (-3)^n}{5^n} = \lim_{n \rightarrow \infty} \left(\frac{e}{5}\right)^n + \lim_{n \rightarrow \infty} \left(\frac{-3}{5}\right)^n$$

assuming both limits on the right-hand side exist. But by the Limit of Geometric Sequences, since

$$-1 < \frac{-3}{5} < 0 < \frac{e}{5} < 1$$

both limits on the right-hand side are 0, so that y_n converges to 0.

$$56. b_n = \frac{(-1)^n n^3 + 2^{-n}}{3n^3 + 4^{-n}}$$

SOLUTION Assuming both limits on the right-hand side exist, we have

$$\lim_{n \rightarrow \infty} \frac{(-1)^n n^3 + 2^{-n}}{3n^3 + 4^{-n}} = \lim_{n \rightarrow \infty} \frac{(-1)^n n^3}{3n^3 + 4^{-n}} + \lim_{n \rightarrow \infty} \frac{2^{-n}}{3n^3 + 4^{-n}}$$

For the first limit, let us consider instead the limit of its reciprocal:

$$\begin{aligned} \lim_{n \rightarrow \infty} (-1)^n \frac{3n^3 + 4^{-n}}{n^3} &= \lim_{n \rightarrow \infty} (-1)^n \frac{3n^3}{n^3} + \lim_{n \rightarrow \infty} (-1)^n \frac{4^{-n}}{n^3} \\ &= \lim_{n \rightarrow \infty} (-1)^n \cdot 3 + \lim_{n \rightarrow \infty} (-1)^n \frac{1}{4^n n^3} \\ &= \lim_{n \rightarrow \infty} ((-1)^n \cdot 3) + 0 \end{aligned}$$

so that one limit on the right-hand side exists and the other does not; thus the left-hand side diverges as well.

$$57. a_n = n \sin \frac{\pi}{n}$$

SOLUTION By the Theorem on Sequences Defined by a Function, we have

$$\lim_{n \rightarrow \infty} n \sin \frac{\pi}{n} = \lim_{x \rightarrow \infty} x \sin \frac{\pi}{x}.$$

Now,

$$\begin{aligned} \lim_{x \rightarrow \infty} x \sin \frac{\pi}{x} &= \lim_{x \rightarrow \infty} \frac{\sin \frac{\pi}{x}}{\frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{\left(\cos \frac{\pi}{x}\right) \left(-\frac{\pi}{x^2}\right)}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \left(\pi \cos \frac{\pi}{x}\right) \\ &= \pi \lim_{x \rightarrow \infty} \cos \frac{\pi}{x} = \pi \cos 0 = \pi \cdot 1 = \pi. \end{aligned}$$

Thus,

$$\lim_{n \rightarrow \infty} n \sin \frac{\pi}{n} = \pi.$$

$$58. b_n = \frac{n!}{\pi^n}$$

SOLUTION By the method of Exercise 41, we can see that $\lim_{n \rightarrow \infty} \frac{4^n}{n!} = 0$ so that $c_n = \frac{n!}{4^n}$ diverges. But $\pi < 4$ so that $c_n < b_n$ and thus b_n diverges as well.

$$59. b_n = \frac{3 - 4^n}{2 + 7 \cdot 4^n}$$

SOLUTION Divide the numerator and denominator by 4^n to obtain

$$a_n = \frac{3 - 4^n}{2 + 7 \cdot 4^n} = \frac{\frac{3}{4^n} - \frac{4^n}{4^n}}{\frac{2}{4^n} + \frac{7 \cdot 4^n}{4^n}} = \frac{\frac{3}{4^n} - 1}{\frac{2}{4^n} + 7}.$$

Thus,

$$\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} \frac{\frac{3}{4^x} - 1}{\frac{2}{4^x} + 7} = \frac{\lim_{x \rightarrow \infty} \left(\frac{3}{4^x} - 1 \right)}{\lim_{x \rightarrow \infty} \left(\frac{2}{4^x} + 7 \right)} = \frac{3 \lim_{x \rightarrow \infty} \frac{1}{4^x} - \lim_{x \rightarrow \infty} 1}{2 \lim_{x \rightarrow \infty} \frac{1}{4^x} - \lim_{x \rightarrow \infty} 7} = \frac{3 \cdot 0 - 1}{2 \cdot 0 + 7} = -\frac{1}{7}.$$

$$60. a_n = \frac{3 - 4^n}{2 + 7 \cdot 3^n}$$

SOLUTION Divide the numerator and denominator by 3^n to obtain

$$a_n = \frac{3 - 4^n}{2 + 7 \cdot 3^n} = \frac{\frac{3}{3^n} - \frac{4^n}{3^n}}{\frac{2}{3^n} + \frac{7 \cdot 3^n}{3^n}} = \frac{\frac{3}{3^n} - \left(\frac{4}{3} \right)^n}{\frac{2}{3^n} + 7}.$$

We examine the limits of the numerator and the denominator:

$$\lim_{n \rightarrow \infty} \left(\frac{3}{3^n} - \left(\frac{4}{3} \right)^n \right) = 3 \lim_{n \rightarrow \infty} \left(\frac{1}{3} \right)^n - 3 \lim_{n \rightarrow \infty} \left(\frac{4}{3} \right)^n = 3 \cdot 0 - \infty = -\infty,$$

whereas

$$\lim_{n \rightarrow \infty} \left(\frac{2}{3^n} + 7 \right) = \lim_{n \rightarrow \infty} \frac{2}{3^n} + \lim_{n \rightarrow \infty} 7 = 2 \lim_{n \rightarrow \infty} \left(\frac{1}{3} \right)^n + \lim_{n \rightarrow \infty} 7 = 2 \cdot 0 + 7 = 7.$$

Thus, $\lim_{n \rightarrow \infty} a_n = -\infty$; that is, the sequence diverges.

$$61. a_n = \left(1 + \frac{1}{n} \right)^n$$

SOLUTION Taking the natural logarithm of both sides of this expression yields

$$\ln a_n = \ln \left(1 + \frac{1}{n} \right)^n = n \ln \left(1 + \frac{1}{n} \right) = \frac{\ln \left(1 + \frac{1}{n} \right)}{\frac{1}{n}}.$$

Thus,

$$\lim_{n \rightarrow \infty} (\ln a_n) = \lim_{x \rightarrow \infty} \frac{\ln \left(1 + \frac{1}{x} \right)}{\frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} \left(\ln \left(1 + \frac{1}{x} \right) \right)}{\frac{d}{dx} \left(\frac{1}{x} \right)} = \lim_{x \rightarrow \infty} \frac{\frac{1}{1 + \frac{1}{x}} \cdot \left(-\frac{1}{x^2} \right)}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{x}} = \frac{1}{1 + 0} = 1.$$

Because $f(x) = e^x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} e^{\ln a_n} = e^{\lim_{n \rightarrow \infty} (\ln a_n)} = e^1 = e.$$

$$62. a_n = \left(1 + \frac{1}{n^2}\right)^n$$

SOLUTION Taking the natural logarithm of both sides of this expression yields

$$\ln a_n = \ln \left(1 + \frac{1}{n^2}\right)^n = n \ln \left(1 + \frac{1}{n^2}\right) = \frac{\ln \left(1 + \frac{1}{n^2}\right)}{\frac{1}{n}}.$$

Thus,

$$\begin{aligned} \lim_{n \rightarrow \infty} (\ln a_n) &= \lim_{x \rightarrow \infty} \frac{\ln(1+x^{-2})}{x^{-1}} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(\ln(1+x^{-2}))}{\frac{d}{dx}(x^{-1})} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{1}{1+x^{-2}}(-2x^{-3})}{-x^{-2}} = \lim_{x \rightarrow \infty} \frac{2x^{-1}}{1+x^{-2}} = \lim_{x \rightarrow \infty} \frac{\frac{2}{x}}{1+\frac{1}{x^2}} = \frac{0}{1+0} = 0. \end{aligned}$$

Because $f(x) = e^x$ is a continuous function, it follows that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} e^{\ln a_n} = e^{\lim_{n \rightarrow \infty} (\ln a_n)} = e^0 = 1.$$

In Exercises 63–66, find the limit of the sequence using L'Hôpital's Rule.

$$63. a_n = \frac{(\ln n)^2}{n}$$

SOLUTION

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{(\ln n)^2}{n} &= \lim_{x \rightarrow \infty} \frac{(\ln x)^2}{x} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(\ln x)^2}{\frac{d}{dx}x} = \lim_{x \rightarrow \infty} \frac{\frac{2 \ln x}{x}}{1} = \lim_{x \rightarrow \infty} \frac{2 \ln x}{x} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} 2 \ln x}{\frac{d}{dx} x} = \lim_{x \rightarrow \infty} \frac{\frac{2}{x}}{1} = \lim_{x \rightarrow \infty} \frac{2}{x} = 0 \end{aligned}$$

$$64. b_n = \sqrt{n} \ln \left(1 + \frac{1}{n}\right)$$

SOLUTION

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt{n} \ln \left(1 + \frac{1}{n}\right) &= \lim_{x \rightarrow \infty} \sqrt{x} \ln \left(1 + \frac{1}{x}\right) = \lim_{x \rightarrow \infty} \frac{\ln \left(1 + \frac{1}{x}\right)}{x^{-1/2}} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} \ln \left(1 + \frac{1}{x}\right)}{\frac{d}{dx} x^{-1/2}} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{1}{1+\frac{1}{x}} \cdot \left(\frac{-1}{x^2}\right)}{\frac{-1}{2} x^{-3/2}} = \lim_{x \rightarrow \infty} \frac{2}{\sqrt{x} \left(1 + \frac{1}{x}\right)} = 0 \end{aligned}$$

$$65. c_n = n(\sqrt{n^2+1} - n)$$

SOLUTION

$$\begin{aligned} \lim_{n \rightarrow \infty} n(\sqrt{n^2+1} - n) &= \lim_{x \rightarrow \infty} x(\sqrt{x^2+1} - x) = \lim_{x \rightarrow \infty} \frac{x(\sqrt{x^2+1} - x)(\sqrt{x^2+1} + x)}{\sqrt{x^2+1} + x} \\ &= \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2+1} + x} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} x}{\frac{d}{dx} \sqrt{x^2+1} + x} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{x}{\sqrt{x^2+1}}} \\ &= \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{\sqrt{\frac{x^2}{x^2+1}}}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{\sqrt{1+(1/x^2)}}} = \frac{1}{2} \end{aligned}$$

$$66. d_n = n^2(\sqrt[3]{n^3+1} - n)$$

SOLUTION We rewrite d_n as follows:

$$\begin{aligned} d_n &= n^2(\sqrt[3]{n^3+1} - n) = n^2 \left(\sqrt[3]{n^3 \left(1 + \frac{1}{n^3}\right)} - n \right) = n^2 \left(n \sqrt[3]{1 + \frac{1}{n^3}} - n \right) \\ &= n^3 \left(\sqrt[3]{1 + \frac{1}{n^3}} - 1 \right) = \frac{\left((1 + n^{-3})^{1/3} - 1 \right)}{n^{-3}}. \end{aligned}$$

Thus,

$$\begin{aligned}\lim_{n \rightarrow \infty} d_n &= \lim_{x \rightarrow \infty} \frac{(1+x^{-3})^{1/3} - 1}{x^{-3}} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} [(1+x^{-3})^{1/3} - 1]}{\frac{d}{dx} [x^{-3}]} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{1}{3}(1+x^{-3})^{-2/3}(-3x^{-4})}{-3x^{-4}} = \lim_{x \rightarrow \infty} \frac{1}{3}(1+x^{-3})^{-2/3} = \lim_{x \rightarrow \infty} \frac{1}{3\left(1+\frac{1}{x^3}\right)^{2/3}} = \frac{1}{3}.\end{aligned}$$

In Exercises 67–70, use the Squeeze Theorem to evaluate $\lim_{n \rightarrow \infty} a_n$ by verifying the given inequality.

$$67. a_n = \frac{1}{\sqrt{n^4 + n^8}}, \quad \frac{1}{\sqrt{2n^4}} \leq a_n \leq \frac{1}{\sqrt{2n^2}}$$

SOLUTION For all $n > 1$ we have $n^4 < n^8$, so the quotient $\frac{1}{\sqrt{n^4 + n^8}}$ is smaller than $\frac{1}{\sqrt{n^4 + n^4}}$ and larger than $\frac{1}{\sqrt{n^8 + n^8}}$. That is,

$$\begin{aligned}a_n &< \frac{1}{\sqrt{n^4 + n^4}} = \frac{1}{\sqrt{n^4 \cdot 2}} = \frac{1}{\sqrt{2n^4}}; \text{ and} \\ a_n &> \frac{1}{\sqrt{n^8 + n^8}} = \frac{1}{\sqrt{2n^8}} = \frac{1}{\sqrt{2n^4}}.\end{aligned}$$

Now, since $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{2n^4}} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{2n^2}} = 0$, the Squeeze Theorem for Sequences implies that $\lim_{n \rightarrow \infty} a_n = 0$.

$$68. c_n = \frac{1}{\sqrt{n^2 + 1}} + \frac{1}{\sqrt{n^2 + 2}} + \cdots + \frac{1}{\sqrt{n^2 + n}},$$

$$\frac{n}{\sqrt{n^2 + n}} \leq c_n \leq \frac{n}{\sqrt{n^2 + 1}}$$

SOLUTION Since each of the n terms in the sum defining c_n is not smaller than $\frac{1}{\sqrt{n^2 + n}}$ and not larger than $\frac{1}{\sqrt{n^2 + 1}}$ we obtain the following inequalities:

$$\begin{aligned}c_n &\geq \underbrace{\frac{1}{\sqrt{n^2 + n}} + \cdots + \frac{1}{\sqrt{n^2 + n}}}_{n \text{ terms}} = n \cdot \frac{1}{\sqrt{n^2 + n}} = \frac{n}{\sqrt{n^2 + n}}; \\ c_n &\leq \underbrace{\frac{1}{\sqrt{n^2 + 1}} + \cdots + \frac{1}{\sqrt{n^2 + 1}}}_{n \text{ terms}} = n \cdot \frac{1}{\sqrt{n^2 + 1}} = \frac{n}{\sqrt{n^2 + 1}}.\end{aligned}$$

Thus,

$$\frac{n}{\sqrt{n^2 + n}} \leq c_n \leq \frac{n}{\sqrt{n^2 + 1}}.$$

We now compute the limits of the two sequences:

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2 + 1}} &= \lim_{n \rightarrow \infty} \frac{\frac{n}{n}}{\frac{\sqrt{n^2 + 1}}{n}} = \lim_{n \rightarrow \infty} \frac{1}{\frac{\sqrt{n^2 + 1}}{\sqrt{n^2}}} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{n^2}}} = 1; \\ \lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2 + n}} &= \lim_{n \rightarrow \infty} \frac{\frac{n}{n}}{\frac{\sqrt{n^2 + n}}{n}} = \lim_{n \rightarrow \infty} \frac{1}{\frac{\sqrt{n^2 + n}}{\sqrt{n^2}}} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{n}}} = 1.\end{aligned}$$

By the Squeeze Theorem we conclude that:

$$\lim_{n \rightarrow \infty} c_n = 1.$$

$$69. a_n = (2^n + 3^n)^{1/n}, \quad 3 \leq a_n \leq (2 \cdot 3^n)^{1/n} = 2^{1/n} \cdot 3$$

SOLUTION Clearly $2^n + 3^n \geq 3^n$ for all $n \geq 1$. Therefore:

$$(2^n + 3^n)^{1/n} \geq (3^n)^{1/n} = 3.$$

Also $2^n + 3^n \leq 3^n + 3^n = 2 \cdot 3^n$, so

$$(2^n + 3^n)^{1/n} \leq (2 \cdot 3^n)^{1/n} = 2^{1/n} \cdot 3.$$

Thus,

$$3 \leq (2^n + 3^n)^{1/n} \leq 2^{1/n} \cdot 3.$$

Because

$$\lim_{n \rightarrow \infty} 2^{1/n} \cdot 3 = 3 \lim_{n \rightarrow \infty} 2^{1/n} = 3 \cdot 1 = 3$$

and $\lim_{n \rightarrow \infty} 3 = 3$, the Squeeze Theorem for Sequences guarantees

$$\lim_{n \rightarrow \infty} (2^n + 3^n)^{1/n} = 3.$$

70. $a_n = (n + 10^n)^{1/n}$, $10 \leq a_n \leq (2 \cdot 10^n)^{1/n}$

SOLUTION Clearly

$$10^n \leq n + 10^n \leq 10^n + 10^n = 2 \cdot 10^n$$

for all $n \geq 0$. Thus


$$10 \leq (n + 10^n)^{1/n} \leq (2 \cdot 10^n)^{1/n}$$

Now,

$$\lim_{n \rightarrow \infty} (2 \cdot 10^n)^{1/n} = \lim_{n \rightarrow \infty} 2^{1/n} \cdot 10 = 10 \lim_{n \rightarrow \infty} 2^{1/n} = 10 \cdot 1 = 10$$

and $\lim_{n \rightarrow \infty} 10 = 10$, so that the Squeeze Theorem for Sequences tells us that

$$\lim_{n \rightarrow \infty} (n + 10^n)^{1/n} = 10$$

71.  Which of the following statements is equivalent to the assertion $\lim_{n \rightarrow \infty} a_n = L$? Explain.

- (a) For every $\epsilon > 0$, the interval $(L - \epsilon, L + \epsilon)$ contains at least one element of the sequence $\{a_n\}$.
 (b) For every $\epsilon > 0$, the interval $(L - \epsilon, L + \epsilon)$ contains all but at most finitely many elements of the sequence $\{a_n\}$.

SOLUTION Statement (b) is equivalent to Definition 1 of the limit, since the assertion “ $|a_n - L| < \epsilon$ for all $n > M$ ” means that $L - \epsilon < a_n < L + \epsilon$ for all $n > M$; that is, the interval $(L - \epsilon, L + \epsilon)$ contains all the elements a_n except (maybe) the finite number of elements a_1, a_2, \dots, a_M .

Statement (a) is not equivalent to the assertion $\lim_{n \rightarrow \infty} a_n = L$. We show this, by considering the following sequence:

$$a_n = \begin{cases} \frac{1}{n} & \text{for odd } n \\ 1 + \frac{1}{n} & \text{for even } n \end{cases}$$

Clearly for every $\epsilon > 0$, the interval $(-\epsilon, \epsilon) = (L - \epsilon, L + \epsilon)$ for $L = 0$ contains at least one element of $\{a_n\}$, but the sequence diverges (rather than converges to $L = 0$). Since the terms in the odd places converge to 0 and the terms in the even places converge to 1. Hence, a_n does not approach one limit.

72. Show that $a_n = \frac{1}{2n+1}$ is decreasing.

SOLUTION Let $f(x) = \frac{1}{2x+1}$. Then

$$f'(x) = -\frac{1}{(2x+1)^2} \cdot 2 = \frac{-2}{(2x+1)^2} < 0 \quad \text{for } x \neq -\frac{1}{2}.$$

Since $f'(x) < 0$ for $x \neq -\frac{1}{2}$, f is decreasing on the interval $x > -\frac{1}{2}$. It follows that $a_n = f(n)$ is also decreasing.

73. Show that $a_n = \frac{3n^2}{n^2 + 2}$ is increasing. Find an upper bound.

SOLUTION Let $f(x) = \frac{3x^2}{x^2 + 2}$. Then

$$f'(x) = \frac{6x(x^2 + 2) - 3x^2 \cdot 2x}{(x^2 + 2)^2} = \frac{12x}{(x^2 + 2)^2}.$$

$f'(x) > 0$ for $x > 0$, hence f is increasing on this interval. It follows that $a_n = f(n)$ is also increasing. We now show that $M = 3$ is an upper bound for a_n , by writing:

$$a_n = \frac{3n^2}{n^2 + 2} \leq \frac{3n^2 + 6}{n^2 + 2} = \frac{3(n^2 + 2)}{n^2 + 2} = 3.$$

That is, $a_n \leq 3$ for all n .

74. Show that $a_n = \sqrt[3]{n+1} - n$ is decreasing.

SOLUTION Let $f(x) = \sqrt[3]{x+1} - x$. Then

$$f'(x) = \frac{d}{dx} \left((x+1)^{1/3} - x \right) = \frac{1}{3}(x+1)^{-2/3} - 1.$$

For $x \geq 1$,

$$\frac{1}{3}(x+1)^{-2/3} - 1 \leq \frac{1}{3}2^{-2/3} - 1 < 0.$$

We conclude that f is decreasing on the interval $x \geq 1$. It follows that $a_n = f(n)$ is also decreasing.

75. Give an example of a divergent sequence $\{a_n\}$ such that $\lim_{n \rightarrow \infty} |a_n|$ converges.

SOLUTION Let $a_n = (-1)^n$. The sequence $\{a_n\}$ diverges because the terms alternate between $+1$ and -1 ; however, the sequence $\{|a_n|\}$ converges because it is a constant sequence, all of whose terms are equal to 1.

76. Give an example of *divergent* sequences $\{a_n\}$ and $\{b_n\}$ such that $\{a_n + b_n\}$ converges.

SOLUTION Let $a_n = 2^n$ and $b_n = -2^n$. Then $\{a_n\}$ and $\{b_n\}$ are divergent geometric sequences. However, since $a_n + b_n = 2^n - 2^n = 0$, the sequence $\{a_n + b_n\}$ is the constant sequence with all the terms equal zero, so it converges to zero.

77. Using the limit definition, prove that if $\{a_n\}$ converges and $\{b_n\}$ diverges, then $\{a_n + b_n\}$ diverges.

SOLUTION We will prove this result by contradiction. Suppose $\lim_{n \rightarrow \infty} a_n = L_1$ and that $\{a_n + b_n\}$ converges to a limit L_2 . Now, let $\epsilon > 0$. Because $\{a_n\}$ converges to L_1 and $\{a_n + b_n\}$ converges to L_2 , it follows that there exist numbers M_1 and M_2 such that:

$$\begin{aligned} |a_n - L_1| &< \frac{\epsilon}{2} && \text{for all } n > M_1, \\ |(a_n + b_n) - L_2| &< \frac{\epsilon}{2} && \text{for all } n > M_2. \end{aligned}$$

Thus, for $n > M = \max\{M_1, M_2\}$,

$$|a_n - L_1| < \frac{\epsilon}{2} \quad \text{and} \quad |(a_n + b_n) - L_2| < \frac{\epsilon}{2}.$$

By the triangle inequality,

$$\begin{aligned} |b_n - (L_2 - L_1)| &= |a_n + b_n - a_n - (L_2 - L_1)| = |(-a_n + L_1) + (a_n + b_n - L_2)| \\ &\leq |L_1 - a_n| + |a_n + b_n - L_2|. \end{aligned}$$

Thus, for $n > M$,

$$|b_n - (L_2 - L_1)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon;$$

that is, $\{b_n\}$ converges to $L_2 - L_1$, in contradiction to the given data. Thus, $\{a_n + b_n\}$ must diverge.

78. Use the limit definition to prove that if $\{a_n\}$ is a convergent sequence of integers with limit L , then there exists a number M such that $a_n = L$ for all $n \geq M$.

SOLUTION Suppose $\{a_n\}$ converges to L , and let $\epsilon = \frac{1}{2}$. Then, there exists a number M such that

$$|a_n - L| < \frac{1}{2}$$

for all $n \geq M$. In other words, for all $n \geq M$,

$$L - \frac{1}{2} < a_n < L + \frac{1}{2}.$$

However, we are given that $\{a_n\}$ is a sequence of integers. Thus, it must be that $a_n = L$ for all $n \geq M$.

79. Theorem 1 states that if $\lim_{x \rightarrow \infty} f(x) = L$, then the sequence $a_n = f(n)$ converges and $\lim_{n \rightarrow \infty} a_n = L$. Show that the converse is false. In other words, find a function $f(x)$ such that $a_n = f(n)$ converges but $\lim_{x \rightarrow \infty} f(x)$ does not exist.

SOLUTION Let $f(x) = \sin \pi x$ and $a_n = \sin \pi n$. Then $a_n = f(n)$. Since $\sin \pi x$ is oscillating between -1 and 1 the limit $\lim_{x \rightarrow \infty} f(x)$ does not exist. However, the sequence $\{a_n\}$ is the constant sequence in which $a_n = \sin \pi n = 0$ for all n , hence it converges to zero.

80. Use the limit definition to prove that the limit does not change if a finite number of terms are added or removed from a convergent sequence.

SOLUTION Suppose that $\{a_n\}$ is a sequence such that $\lim_{n \rightarrow \infty} a_n = L$. For every $\epsilon > 0$, there is a number M such that $|a_n - L| < \epsilon$ for all $n > M$. That is, the inequality $|a_n - L| < \epsilon$ holds for all the terms of $\{a_n\}$ except possibly a finite number of terms. If we add a finite number of terms, these terms may not satisfy the inequality $|a_n - L| < \epsilon$, but there are still only a finite number of terms that do not satisfy this inequality. By removing terms from the sequence, the number of terms in the new sequence that do not satisfy $|a_n - L| < \epsilon$ are no more than in the original sequence. Hence the new sequence also converges to L .

81. Let $b_n = a_{n+1}$. Use the limit definition to prove that if $\{a_n\}$ converges, then $\{b_n\}$ also converges and $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$.

SOLUTION Suppose $\{a_n\}$ converges to L . Let $b_n = a_{n+1}$, and let $\epsilon > 0$. Because $\{a_n\}$ converges to L , there exists an M' such that $|a_n - L| < \epsilon$ for $n > M'$. Now, let $M = M' - 1$. Then, whenever $n > M$, $n + 1 > M + 1 = M'$. Thus, for $n > M$,

$$|b_n - L| = |a_{n+1} - L| < \epsilon.$$

Hence, $\{b_n\}$ converges to L .

82. Let $\{a_n\}$ be a sequence such that $\lim_{n \rightarrow \infty} |a_n|$ exists and is nonzero. Show that $\lim_{n \rightarrow \infty} a_n$ exists if and only if there exists an integer M such that the sign of a_n does not change for $n > M$.

SOLUTION Let $\{a_n\}$ be a sequence such that $\lim_{n \rightarrow \infty} |a_n|$ exists and is nonzero. Suppose $\lim_{n \rightarrow \infty} a_n$ exists and let $L = \lim_{n \rightarrow \infty} a_n$. Note that L cannot be zero for then $\lim_{n \rightarrow \infty} |a_n|$ would also be zero. Now, choose $\epsilon < |L|$. Then there exists an integer M such that $|a_n - L| < \epsilon$, or $L - \epsilon < a_n < L + \epsilon$, for all $n > M$. If $L < 0$, then $-2L < a_n < 0$, whereas if $L > 0$, then $0 < a_n < 2L$; that is, a_n does not change for $n > M$.

Now suppose that there exists an integer M such that a_n does not change for $n > M$. If $a_n > 0$ for $n > M$, then $a_n = |a_n|$ for $n > M$ and

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} |a_n|.$$

On the other hand, if $a_n < 0$ for $n > M$, then $a_n = -|a_n|$ for $n > M$ and

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} -|a_n| = -\lim_{n \rightarrow \infty} |a_n|.$$

In either case, $\lim_{n \rightarrow \infty} a_n$ exists. Thus, $\lim_{n \rightarrow \infty} a_n$ exists if and only if there exists an integer M such that the sign of a_n does not change for $n > M$.

83. Proceed as in Example 12 to show that the sequence $\sqrt{3}, \sqrt{3\sqrt{3}}, \sqrt{3\sqrt{3\sqrt{3}}}, \dots$ is increasing and bounded above by $M = 3$. Then prove that the limit exists and find its value.

SOLUTION This sequence is defined recursively by the formula:

$$a_{n+1} = \sqrt{3a_n}, \quad a_1 = \sqrt{3}.$$

Consider the following inequalities:

$$\begin{aligned} a_2 &= \sqrt{3a_1} = \sqrt{3\sqrt{3}} > \sqrt{3} = a_1 && \Rightarrow a_2 > a_1; \\ a_3 &= \sqrt{3a_2} > \sqrt{3a_1} = a_2 && \Rightarrow a_3 > a_2; \\ a_4 &= \sqrt{3a_3} > \sqrt{3a_2} = a_3 && \Rightarrow a_4 > a_3. \end{aligned}$$

In general, if we assume that $a_k > a_{k-1}$, then

$$a_{k+1} = \sqrt{3a_k} > \sqrt{3a_{k-1}} = a_k.$$

Hence, by mathematical induction, $a_{n+1} > a_n$ for all n ; that is, the sequence $\{a_n\}$ is increasing.

Because $a_{n+1} = \sqrt{3a_n}$, it follows that $a_n \geq 0$ for all n . Now, $a_1 = \sqrt{3} < 3$. If $a_k \leq 3$, then

$$a_{k+1} = \sqrt{3a_k} \leq \sqrt{3 \cdot 3} = 3.$$

Thus, by mathematical induction, $a_n \leq 3$ for all n .

Since $\{a_n\}$ is increasing and bounded, it follows by the Theorem on Bounded Monotonic Sequences that this sequence is converging. Denote the limit by $L = \lim_{n \rightarrow \infty} a_n$. Using Exercise 81, it follows that

$$L = \lim_{n \rightarrow \infty} a_{n+1} = \lim_{n \rightarrow \infty} \sqrt{3a_n} = \sqrt{3 \lim_{n \rightarrow \infty} a_n} = \sqrt{3L}.$$

Thus, $L^2 = 3L$, so $L = 0$ or $L = 3$. Because the sequence is increasing, we have $a_n \geq a_1 = \sqrt{3}$ for all n . Hence, the limit also satisfies $L \geq \sqrt{3}$. We conclude that the appropriate solution is $L = 3$; that is, $\lim_{n \rightarrow \infty} a_n = 3$.

84. Let $\{a_n\}$ be the sequence defined recursively by

$$a_0 = 0, \quad a_{n+1} = \sqrt{2 + a_n}$$

Thus, $a_1 = \sqrt{2}$, $a_2 = \sqrt{2 + \sqrt{2}}$, $a_3 = \sqrt{2 + \sqrt{2 + \sqrt{2}}}$, \dots

(a) Show that if $a_n < 2$, then $a_{n+1} < 2$. Conclude by induction that $a_n < 2$ for all n .

(b) Show that if $a_n < 2$, then $a_n \leq a_{n+1}$. Conclude by induction that $\{a_n\}$ is increasing.

(c) Use (a) and (b) to conclude that $L = \lim_{n \rightarrow \infty} a_n$ exists. Then compute L by showing that $L = \sqrt{2 + L}$.

SOLUTION

(a) Assume $a_n < 2$. Then

$$a_{n+1} = \sqrt{2 + a_n} < \sqrt{2 + 2} = 2$$

so that $a_{n+1} < 2$. So by induction, $a_n < 2$ for all n and $\{a_n\}$ is bounded above by 2.

(b) Assume $a_n < 2$. Then

$$a_{n+1} = \sqrt{2 + a_n} > \sqrt{a_n + a_n} = \sqrt{2a_n} > \sqrt{a_n^2} = a_n$$

so that $a_n < a_{n+1}$. It follows by induction that $\{a_n\}$ is increasing.

(c) Since $\{a_n\}$ is increasing and bounded above, the Theorem on Bounded Monotone Sequences tells us that $L = \lim_{n \rightarrow \infty} a_n$ exists. We have

$$L = \lim_{n \rightarrow \infty} a_{n+1} = \lim_{n \rightarrow \infty} \sqrt{2 + a_n} = \sqrt{2 + \lim_{n \rightarrow \infty} a_n} = \sqrt{2 + L}$$

by Exercise 81. It follows that $L = \sqrt{2 + L}$, so that $L^2 - L - 2 = 0$. Thus $L = 2$ or $L = -1$. But all terms of $\{a_n\}$ are positive, so we must have $L = 2$.

Further Insights and Challenges

85. Show that $\lim_{n \rightarrow \infty} \sqrt[n]{n!} = \infty$. *Hint:* Verify that $n! \geq (n/2)^{n/2}$ by observing that half of the factors of $n!$ are greater than or equal to $n/2$.

SOLUTION We show that $n! \geq (\frac{n}{2})^{n/2}$. For $n \geq 4$ even, we have:

$$n! = \underbrace{1 \cdots \frac{n}{2}}_{\frac{n}{2} \text{ factors}} \cdot \underbrace{\left(\frac{n}{2} + 1\right) \cdots n}_{\frac{n}{2} \text{ factors}} \geq \underbrace{\left(\frac{n}{2} + 1\right) \cdots n}_{\frac{n}{2} \text{ factors}}$$

Since each one of the $\frac{n}{2}$ factors is greater than $\frac{n}{2}$, we have:

$$n! \geq \underbrace{\left(\frac{n}{2} + 1\right) \cdots n}_{\frac{n}{2} \text{ factors}} \geq \underbrace{\frac{n}{2} \cdots \frac{n}{2}}_{\frac{n}{2} \text{ factors}} = \left(\frac{n}{2}\right)^{n/2}.$$

For $n \geq 3$ odd, we have:

$$n! = 1 \cdots \underbrace{\frac{n-1}{2}}_{\frac{n-1}{2} \text{ factors}} \cdot \underbrace{\frac{n+1}{2} \cdots n}_{\frac{n+1}{2} \text{ factors}} \geq \underbrace{\frac{n+1}{2} \cdots n}_{\frac{n+1}{2} \text{ factors}}.$$

Since each one of the $\frac{n+1}{2}$ factors is greater than $\frac{n}{2}$, we have:

$$n! \geq \underbrace{\frac{n+1}{2} \cdots n}_{\frac{n+1}{2} \text{ factors}} \geq \underbrace{\frac{n}{2} \cdots \frac{n}{2}}_{\frac{n+1}{2} \text{ factors}} = \left(\frac{n}{2}\right)^{(n+1)/2} = \left(\frac{n}{2}\right)^{n/2} \sqrt{\frac{n}{2}} \geq \left(\frac{n}{2}\right)^{n/2}.$$

In either case we have $n! \geq \left(\frac{n}{2}\right)^{n/2}$. Thus,

$$\sqrt[n]{n!} \geq \sqrt{\frac{n}{2}}.$$

Since $\lim_{n \rightarrow \infty} \sqrt{\frac{n}{2}} = \infty$, it follows that $\lim_{n \rightarrow \infty} \sqrt[n]{n!} = \infty$. Thus, the sequence $a_n = \sqrt[n]{n!}$ diverges.

86. Let $b_n = \frac{\sqrt[n]{n!}}{n}$.

(a) Show that $\ln b_n = \frac{1}{n} \sum_{k=1}^n \ln \frac{k}{n}$.

(b) Show that $\ln b_n$ converges to $\int_0^1 \ln x \, dx$, and conclude that $b_n \rightarrow e^{-1}$.

SOLUTION

(a) Let $b_n = \frac{(n!)^{1/n}}{n}$. Then

$$\begin{aligned} \ln b_n &= \ln (n!)^{1/n} - \ln n = \frac{1}{n} \ln (n!) - \ln n = \frac{\ln (n!) - n \ln n}{n} = \frac{1}{n} [\ln (n!) - \ln n^n] = \frac{1}{n} \ln \frac{n!}{n^n} \\ &= \frac{1}{n} \ln \left(\frac{1}{n} \cdot \frac{2}{n} \cdot \frac{3}{n} \cdots \frac{n}{n} \right) = \frac{1}{n} \left(\ln \frac{1}{n} + \ln \frac{2}{n} + \ln \frac{3}{n} + \cdots + \ln \frac{n}{n} \right) = \frac{1}{n} \sum_{k=1}^n \ln \frac{k}{n}. \end{aligned}$$

(b) By part (a) we have,

$$\lim_{n \rightarrow \infty} (\ln b_n) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \ln \frac{k}{n}.$$

Notice that $\frac{1}{n} \sum_{k=1}^n \ln \frac{k}{n}$ is the n th right-endpoint approximation of the integral of $\ln x$ over the interval $[0, 1]$. Hence,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \ln \frac{k}{n} = \int_0^1 \ln x \, dx.$$

We compute the improper integral using integration by parts, with $u = \ln x$ and $v' = 1$. Then $u' = \frac{1}{x}$, $v = x$ and

$$\begin{aligned} \int_0^1 \ln x \, dx &= x \ln x \Big|_0^1 - \int_0^1 \frac{1}{x} x \, dx = 1 \cdot \ln 1 - \lim_{x \rightarrow 0^+} (x \ln x) - \int_0^1 dx \\ &= 0 - \lim_{x \rightarrow 0^+} (x \ln x) - x \Big|_0^1 = -1 - \lim_{x \rightarrow 0^+} (x \ln x). \end{aligned}$$

We compute the remaining limit using L'Hôpital's Rule. This gives:

$$\lim_{x \rightarrow 0^+} (x \cdot \ln x) = \lim_{x \rightarrow 0^+} \frac{\ln x}{\frac{1}{x}} = \lim_{x \rightarrow 0^+} \frac{\frac{1}{x}}{-\frac{1}{x^2}} = \lim_{x \rightarrow 0^+} (-x) = 0.$$

Thus,

$$\lim_{n \rightarrow \infty} \ln b_n = \int_0^1 \ln x \, dx = -1,$$

and

$$\lim_{n \rightarrow \infty} b_n = e^{-1}.$$

87. Given positive numbers $a_1 < b_1$, define two sequences recursively by

$$a_{n+1} = \sqrt{a_n b_n}, \quad b_{n+1} = \frac{a_n + b_n}{2}$$

- (a) Show that $a_n \leq b_n$ for all n (Figure 13).
 (b) Show that $\{a_n\}$ is increasing and $\{b_n\}$ is decreasing.
 (c) Show that $b_{n+1} - a_{n+1} \leq \frac{b_n - a_n}{2}$.
 (d) Prove that both $\{a_n\}$ and $\{b_n\}$ converge and have the same limit. This limit, denoted $\text{AGM}(a_1, b_1)$, is called the **arithmetic-geometric mean** of a_1 and b_1 .
 (e) Estimate $\text{AGM}(1, \sqrt{2})$ to three decimal places.




FIGURE 13

SOLUTION

(a) Examine the following:

$$\begin{aligned} b_{n+1} - a_{n+1} &= \frac{a_n + b_n}{2} - \sqrt{a_n b_n} = \frac{a_n + b_n - 2\sqrt{a_n b_n}}{2} = \frac{(\sqrt{a_n})^2 - 2\sqrt{a_n}\sqrt{b_n} + (\sqrt{b_n})^2}{2} \\ &= \frac{(\sqrt{a_n} - \sqrt{b_n})^2}{2} \geq 0. \end{aligned}$$

We conclude that $b_{n+1} \geq a_{n+1}$ for all $n > 1$. By the given information $b_1 > a_1$; hence, $b_n \geq a_n$ for all n .

(b) By part (a), $b_n \geq a_n$ for all n , so

$$a_{n+1} = \sqrt{a_n b_n} \geq \sqrt{a_n \cdot a_n} = \sqrt{a_n^2} = a_n$$

for all n . Hence, the sequence $\{a_n\}$ is increasing. Moreover, since $a_n \leq b_n$ for all n ,

$$b_{n+1} = \frac{a_n + b_n}{2} \leq \frac{b_n + b_n}{2} = \frac{2b_n}{2} = b_n$$

for all n ; that is, the sequence $\{b_n\}$ is decreasing.

(c) Since $\{a_n\}$ is increasing, $a_{n+1} \geq a_n$. Thus,

$$b_{n+1} - a_{n+1} \leq b_{n+1} - a_n = \frac{a_n + b_n}{2} - a_n = \frac{a_n + b_n - 2a_n}{2} = \frac{b_n - a_n}{2}.$$

Now, by part (a), $a_n \leq b_n$ for all n . By part (b), $\{b_n\}$ is decreasing. Hence $b_n \leq b_1$ for all n . Combining the two inequalities we conclude that $a_n \leq b_1$ for all n . That is, the sequence $\{a_n\}$ is increasing and bounded ($0 \leq a_n \leq b_1$). By the Theorem on Bounded Monotonic Sequences we conclude that $\{a_n\}$ converges. Similarly, since $\{a_n\}$ is increasing, $a_n \geq a_1$ for all n . We combine this inequality with $b_n \geq a_n$ to conclude that $b_n \geq a_1$ for all n . Thus, $\{b_n\}$ is decreasing and bounded ($a_1 \leq b_n \leq b_1$); hence this sequence converges.

To show that $\{a_n\}$ and $\{b_n\}$ converge to the same limit, note that

$$b_n - a_n \leq \frac{b_{n-1} - a_{n-1}}{2} \leq \frac{b_{n-2} - a_{n-2}}{2^2} \leq \dots \leq \frac{b_1 - a_1}{2^{n-1}}.$$

Thus,

$$\lim_{n \rightarrow \infty} (b_n - a_n) = (b_1 - a_1) \lim_{n \rightarrow \infty} \frac{1}{2^{n-1}} = 0.$$

(d) We have

$$a_{n+1} = \sqrt{a_n b_n}, \quad a_1 = 1; \quad b_{n+1} = \frac{a_n + b_n}{2}, \quad b_1 = \sqrt{2}$$

Computing the values of a_n and b_n until the first three decimal digits are equal in successive terms, we obtain:

$$\begin{aligned} a_2 &= \sqrt{a_1 b_1} = \sqrt{1 \cdot \sqrt{2}} = 1.1892 \\ b_2 &= \frac{a_1 + b_1}{2} = \frac{1 + \sqrt{2}}{2} = 1.2071 \\ a_3 &= \sqrt{a_2 b_2} = \sqrt{1.1892 \cdot 1.2071} = 1.1981 \\ b_3 &= \frac{a_2 + b_2}{2} = \frac{1.1892 + 1.2071}{2} = 1.1981 \\ a_4 &= \sqrt{a_3 b_3} = 1.1981 \\ b_4 &= \frac{a_3 + b_3}{2} = 1.1981 \end{aligned}$$

Thus,

$$AGM(1, \sqrt{2}) \approx 1.198.$$

88. Let $c_n = \frac{1}{n} + \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n}$.

(a) Calculate c_1, c_2, c_3, c_4 .

(b) Use a comparison of rectangles with the area under $y = x^{-1}$ over the interval $[n, 2n]$ to prove that

$$\int_n^{2n} \frac{dx}{x} + \frac{1}{2n} \leq c_n \leq \int_n^{2n} \frac{dx}{x} + \frac{1}{n}$$

(c) Use the Squeeze Theorem to determine $\lim_{n \rightarrow \infty} c_n$.


SOLUTION

(a)

$$\begin{aligned} c_1 &= 1 + \frac{1}{2} = \frac{3}{2}; \\ c_2 &= \frac{1}{2} + \frac{1}{3} + \frac{1}{4} = \frac{13}{12}; \\ c_3 &= \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} = \frac{19}{20}; \\ c_4 &= \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} = \frac{743}{840}; \end{aligned}$$

(b) We consider the left endpoint approximation to the integral of $y = \frac{1}{x}$ over the interval $[n, 2n]$. Since the function $y = \frac{1}{x}$ is decreasing, the left endpoint approximation is greater than $\int_n^{2n} \frac{dx}{x}$; that is,

$$\int_n^{2n} \frac{dx}{x} \leq \frac{1}{n} \cdot 1 + \frac{1}{n+1} \cdot 1 + \frac{1}{n+2} \cdot 1 + \cdots + \frac{1}{2n-1} \cdot 1.$$




We express the right hand-side of the inequality in terms of c_n , obtaining:

$$\int_n^{2n} \frac{dx}{x} \leq c_n - \frac{1}{2n}.$$

We now consider the right endpoint approximation to the integral $\int_n^{2n} \frac{dx}{x}$; that is,

$$\frac{1}{n+1} \cdot 1 + \frac{1}{n+2} \cdot 1 + \cdots + \frac{1}{2n} \cdot 1 \leq \int_n^{2n} \frac{dx}{x}.$$



We express the left hand-side of the inequality in terms of c_n , obtaining:

$$c_n - \frac{1}{n} \leq \int_n^{2n} \frac{dx}{x}.$$

Thus,

$$\int_n^{2n} \frac{dx}{x} + \frac{1}{2n} \leq c_n \leq \int_n^{2n} \frac{dx}{x} + \frac{1}{n}.$$

(c) With

$$\int_n^{2n} \frac{dx}{x} = \ln x \Big|_n^{2n} = \ln 2n - \ln n = \ln \frac{2n}{n} = \ln 2,$$

the result from part (b) becomes


$$\ln 2 + \frac{1}{2n} \leq c_n \leq \ln 2 + \frac{1}{n}.$$

Because

$$\lim_{n \rightarrow \infty} \frac{1}{2n} = \lim_{n \rightarrow \infty} \frac{1}{n} = 0,$$

it follows from the Squeeze Theorem that

$$\lim_{n \rightarrow \infty} c_n = \ln 2.$$

89.  Let $a_n = H_n - \ln n$, where H_n is the n th harmonic number

$$H_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$$

- (a) Show that $a_n \geq 0$ for $n \geq 1$. *Hint:* Show that $H_n \geq \int_1^{n+1} \frac{dx}{x}$.
 (b) Show that $\{a_n\}$ is decreasing by interpreting $a_n - a_{n+1}$ as an area.
 (c) Prove that $\lim_{n \rightarrow \infty} a_n$ exists.

This limit, denoted γ , is known as *Euler's Constant*. It appears in many areas of mathematics, including analysis and number theory, and has been calculated to more than 100 million decimal places, but it is still not known whether γ is an irrational number. The first 10 digits are $\gamma \approx 0.5772156649$.


SOLUTION

(a) Since the function $y = \frac{1}{x}$ is decreasing, the left endpoint approximation to the integral $\int_1^{n+1} \frac{dx}{x}$ is greater than this integral; that is,

$$1 \cdot 1 + \frac{1}{2} \cdot 1 + \frac{1}{3} \cdot 1 + \cdots + \frac{1}{n} \cdot 1 \geq \int_1^{n+1} \frac{dx}{x}$$

or

$$H_n \geq \int_1^{n+1} \frac{dx}{x}.$$



Moreover, since the function $y = \frac{1}{x}$ is positive for $x > 0$, we have:

$$\int_1^{n+1} \frac{dx}{x} \geq \int_1^n \frac{dx}{x}.$$

Thus,

$$H_n \geq \int_1^n \frac{dx}{x} = \ln x \Big|_1^n = \ln n - \ln 1 = \ln n,$$

and


$$a_n = H_n - \ln n \geq 0 \quad \text{for all } n \geq 1.$$

(b) To show that $\{a_n\}$ is decreasing, we consider the difference $a_n - a_{n+1}$:

$$\begin{aligned} a_n - a_{n+1} &= H_n - \ln n - (H_{n+1} - \ln(n+1)) = H_n - H_{n+1} + \ln(n+1) - \ln n \\ &= 1 + \frac{1}{2} + \cdots + \frac{1}{n} - \left(1 + \frac{1}{2} + \cdots + \frac{1}{n} + \frac{1}{n+1}\right) + \ln(n+1) - \ln n \\ &= -\frac{1}{n+1} + \ln(n+1) - \ln n. \end{aligned}$$

Now, $\ln(n+1) - \ln n = \int_n^{n+1} \frac{dx}{x}$, whereas $\frac{1}{n+1}$ is the right endpoint approximation to the integral $\int_n^{n+1} \frac{dx}{x}$. Recalling $y = \frac{1}{x}$ is decreasing, it follows that

$$\int_n^{n+1} \frac{dx}{x} \geq \frac{1}{n+1}$$



so

$$a_n - a_{n+1} \geq 0.$$

(c) By parts (a) and (b), $\{a_n\}$ is decreasing and 0 is a lower bound for this sequence. Hence $0 \leq a_n \leq a_1$ for all n . A monotonic and bounded sequence is convergent, so $\lim_{n \rightarrow \infty} a_n$ exists.

10.2 Summing an Infinite Series

Preliminary Questions

1. What role do partial sums play in defining the sum of an infinite series?

SOLUTION The sum of an infinite series is defined as the limit of the sequence of partial sums. If the limit of this sequence does not exist, the series is said to diverge.

2. What is the sum of the following infinite series?

$$\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \frac{1}{64} + \cdots$$

SOLUTION This is a geometric series with $c = \frac{1}{4}$ and $r = \frac{1}{2}$. The sum of the series is therefore

$$\frac{\frac{1}{4}}{1 - \frac{1}{2}} = \frac{\frac{1}{4}}{\frac{1}{2}} = \frac{1}{2}.$$

3. What happens if you apply the formula for the sum of a geometric series to the following series? Is the formula valid?

$$1 + 3 + 3^2 + 3^3 + 3^4 + \cdots$$

SOLUTION This is a geometric series with $c = 1$ and $r = 3$. Applying the formula for the sum of a geometric series then gives

$$\sum_{n=0}^{\infty} 3^n = \frac{1}{1-3} = -\frac{1}{2}.$$

Clearly, this is not valid: a series with all positive terms cannot have a negative sum. The formula is not valid in this case because a geometric series with $r = 3$ diverges.

4. Arvind asserts that $\sum_{n=1}^{\infty} \frac{1}{n^2} = 0$ because $\frac{1}{n^2}$ tends to zero. Is this valid reasoning?

SOLUTION Arvind's reasoning is not valid. Though the terms in the series do tend to zero, the general term in the sequence of partial sums,

$$S_n = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \cdots + \frac{1}{n^2},$$

is clearly larger than 1. The sum of the series therefore cannot be zero.

5. Colleen claims that $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ converges because

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$$

Is this valid reasoning?

SOLUTION Colleen's reasoning is not valid. Although the general term of a convergent series must tend to zero, a series whose general term tends to zero need not converge. In the case of $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$, the series diverges even though its general term tends to zero.

6. Find an N such that $S_N > 25$ for the series $\sum_{n=1}^{\infty} 2$.

SOLUTION The N th partial sum of the series is:

$$S_N = \sum_{n=1}^N 2 = \underbrace{2 + \cdots + 2}_N = 2N.$$

7. Does there exist an N such that $S_N > 25$ for the series $\sum_{n=1}^{\infty} 2^{-n}$? Explain.

SOLUTION The series $\sum_{n=1}^{\infty} 2^{-n}$ is a convergent geometric series with the common ratio $r = \frac{1}{2}$. The sum of the series is:

$$S = \frac{\frac{1}{2}}{1 - \frac{1}{2}} = 1.$$

Notice that the sequence of partial sums $\{S_N\}$ is increasing and converges to 1; therefore $S_N \leq 1$ for all N . Thus, there does not exist an N such that $S_N > 25$.

8. Give an example of a divergent infinite series whose general term tends to zero.

SOLUTION Consider the series $\sum_{n=1}^{\infty} \frac{1}{n^{\frac{9}{10}}}$. The general term tends to zero, since $\lim_{n \rightarrow \infty} \frac{1}{n^{\frac{9}{10}}} = 0$. However, the N th partial sum satisfies the following inequality:

$$S_N = \frac{1}{1^{\frac{9}{10}}} + \frac{1}{2^{\frac{9}{10}}} + \cdots + \frac{1}{N^{\frac{9}{10}}} \geq \frac{N}{N^{\frac{9}{10}}} = N^{1-\frac{9}{10}} = N^{\frac{1}{10}}.$$

That is, $S_N \geq N^{\frac{1}{10}}$ for all N . Since $\lim_{N \rightarrow \infty} N^{\frac{1}{10}} = \infty$, the sequence of partial sums S_n diverges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{n^{\frac{9}{10}}}$ diverges.

Exercises

1. Find a formula for the general term a_n (not the partial sum) of the infinite series.

(a) $\frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \frac{1}{81} + \cdots$

(b) $\frac{1}{1} + \frac{5}{2} + \frac{25}{4} + \frac{125}{8} + \cdots$

(c) $\frac{1}{1} - \frac{2^2}{2 \cdot 1} + \frac{3^3}{3 \cdot 2 \cdot 1} - \frac{4^4}{4 \cdot 3 \cdot 2 \cdot 1} + \cdots$

(d) $\frac{2}{1^2+1} + \frac{1}{2^2+1} + \frac{2}{3^2+1} + \frac{1}{4^2+1} + \cdots$

SOLUTION

(a) The denominators of the terms are powers of 3, starting with the first power. Hence, the general term is:

$$a_n = \frac{1}{3^n}.$$

(b) The numerators are powers of 5, and the denominators are the same powers of 2. The first term is $a_1 = 1$ so,

$$a_n = \left(\frac{5}{2}\right)^{n-1}.$$

(c) The general term of this series is,

$$a_n = (-1)^{n+1} \frac{n^n}{n!}.$$

(d) Notice that the numerators of a_n equal 2 for odd values of n and 1 for even values of n . Thus,

$$a_n = \begin{cases} \frac{2}{n^2+1} & \text{odd } n \\ \frac{1}{n^2+1} & \text{even } n \end{cases}$$

The formula can also be rewritten as follows:

$$a_n = \frac{1 + \frac{(-1)^{n+1}+1}{2}}{n^2+1}.$$

2. Write in summation notation:

(a) $1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \cdots$

(b) $\frac{1}{9} + \frac{1}{16} + \frac{1}{25} + \frac{1}{36} + \cdots$

(c) $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots$

(d) $\frac{125}{9} + \frac{625}{16} + \frac{3125}{25} + \frac{15,625}{36} + \cdots$

SOLUTION

(a) The general term is $a_n = \frac{1}{n^2}$, $n = 1, 2, 3, \dots$; hence, the series is $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

(b) The general term is $a_n = \frac{1}{n^2}$, $n = 3, 4, 5, \dots$ or $a_n = \frac{1}{(n+2)^2}$, $n = 1, 2, 3, \dots$; hence, the series is $\sum_{n=3}^{\infty} \frac{1}{n^2} = \sum_{n=1}^{\infty} \frac{1}{(n+2)^2}$.

(c) The general term is $a_n = \frac{(-1)^{n+1}}{2n-1}$, $n = 1, 2, 3, \dots$; hence, the series is $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1}$.

(d) The general term is $a_n = \frac{5^n}{n^2}$, $n = 3, 4, 5, \dots$ or $a_n = \frac{5^{n+2}}{(n+2)^2}$, $n = 1, 2, 3, \dots$; hence, the series is $\sum_{n=3}^{\infty} \frac{5^n}{n^2} = \sum_{n=1}^{\infty} \frac{5^{n+2}}{(n+2)^2}$.

In Exercises 3–6, compute the partial sums S_2 , S_4 , and S_6 .

$$3. 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

SOLUTION

$$S_2 = 1 + \frac{1}{2^2} = \frac{5}{4};$$

$$S_4 = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} = \frac{205}{144};$$

$$S_6 = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} = \frac{5369}{3600}.$$

$$4. \sum_{k=1}^{\infty} (-1)^k k^{-1}$$

SOLUTION

$$S_2 = (-1)^1 \cdot 1^{-1} + (-1)^2 \cdot 2^{-1} = -1 + \frac{1}{2} = -\frac{1}{2};$$

$$S_4 = (-1)^1 \cdot 1^{-1} + (-1)^2 \cdot 2^{-1} + (-1)^3 \cdot 3^{-1} + (-1)^4 \cdot 4^{-1} = S_2 - \frac{1}{3} + \frac{1}{4} = -\frac{1}{2} - \frac{1}{3} + \frac{1}{4} = -\frac{7}{12};$$

$$S_6 = -\frac{7}{12} + (-1)^5 \cdot 5^{-1} + (-1)^6 \cdot 6^{-1} = -\frac{7}{12} - \frac{1}{5} + \frac{1}{6} = -\frac{37}{60}.$$

$$5. \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots$$

SOLUTION

$$S_2 = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} = \frac{1}{2} + \frac{1}{6} = \frac{4}{6} = \frac{2}{3};$$

$$S_4 = S_2 + a_3 + a_4 = \frac{2}{3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} = \frac{2}{3} + \frac{1}{12} + \frac{1}{20} = \frac{4}{5};$$

$$S_6 = S_4 + a_5 + a_6 = \frac{4}{5} + \frac{1}{5 \cdot 6} + \frac{1}{6 \cdot 7} = \frac{4}{5} + \frac{1}{30} + \frac{1}{42} = \frac{6}{7}.$$

$$6. \sum_{j=1}^{\infty} \frac{1}{j!}$$

SOLUTION

$$S_2 = \frac{1}{1!} + \frac{1}{2!} = 1 + \frac{1}{2} = \frac{3}{2};$$

$$S_4 = S_2 + \frac{1}{3!} + \frac{1}{4!} = \frac{3}{2} + \frac{1}{6} + \frac{1}{24} = \frac{41}{24};$$

$$S_6 = S_4 + \frac{1}{5!} + \frac{1}{6!} = \frac{41}{24} + \frac{1}{120} + \frac{1}{720} = \frac{1237}{720}.$$

7. The series $S = 1 + \left(\frac{1}{5}\right) + \left(\frac{1}{5}\right)^2 + \left(\frac{1}{5}\right)^3 + \cdots$ converges to $\frac{5}{4}$. Calculate S_N for $N = 1, 2, \dots$ until you find an S_N that approximates $\frac{5}{4}$ with an error less than 0.0001.

SOLUTION

$$S_1 = 1$$

$$S_2 = 1 + \frac{1}{5} = \frac{6}{5} = 1.2$$

$$S_3 = 1 + \frac{1}{5} + \frac{1}{25} = \frac{31}{25} = 1.24$$

$$S_3 = 1 + \frac{1}{5} + \frac{1}{25} + \frac{1}{125} = \frac{156}{125} = 1.248$$

$$S_4 = 1 + \frac{1}{5} + \frac{1}{25} + \frac{1}{125} + \frac{1}{625} = \frac{781}{625} = 1.2496$$

$$S_5 = 1 + \frac{1}{5} + \frac{1}{25} + \frac{1}{125} + \frac{1}{625} + \frac{1}{3125} = \frac{3906}{3125} = 1.24992$$

Note that

$$1.25 - S_5 = 1.25 - 1.24992 = 0.00008 < 0.0001$$

8. The series $S = \frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \cdots$ is known to converge to e^{-1} (recall that $0! = 1$). Calculate S_N for $N = 1, 2, \dots$ until you find an S_N that approximates e^{-1} with an error less than 0.001.

SOLUTION The general term of the series is

$$a_n = \frac{(-1)^{n-1}}{(n-1)!};$$

thus, the N th partial sum of the series is

$$S_N = \sum_{n=1}^N a_n = \sum_{n=1}^N \frac{(-1)^{n-1}}{(n-1)!} = \frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \cdots + \frac{(-1)^{N-1}}{(N-1)!}.$$

Using a calculator we find $e^{-1} = 0.367879$. Working sequentially, we find

$$S_1 = \frac{1}{0!} = 1$$

$$S_2 = S_1 + a_2 = 1 - \frac{1}{1!} = 0$$

$$S_3 = S_2 + a_3 = 0 + \frac{1}{2!} = \frac{1}{2} = 0.5$$

$$S_4 = S_3 + a_4 = 0.5 - \frac{1}{3!} = 0.333333$$

$$S_5 = S_4 + a_5 = 0.333333 + \frac{1}{4!} = 0.375$$

$$S_6 = S_5 + a_6 = 0.375 - \frac{1}{5!} = 0.366667$$

$$S_7 = S_6 + a_7 = 0.366667 + \frac{1}{6!} = 0.368056$$

Note that

$$|S_7 - e^{-1}| = 1.76 \times 10^{-4} < 10^{-3}.$$

In Exercises 9 and 10, use a computer algebra system to compute S_{10} , S_{100} , S_{500} , and S_{1000} for the series. Do these values suggest convergence to the given value?

9. $\zeta(3)$

$$\frac{\pi - 3}{4} = \frac{1}{2 \cdot 3 \cdot 4} - \frac{1}{4 \cdot 5 \cdot 6} + \frac{1}{6 \cdot 7 \cdot 8} - \frac{1}{8 \cdot 9 \cdot 10} + \cdots$$

SOLUTION Write

$$a_n = \frac{(-1)^{n+1}}{2n \cdot (2n+1) \cdot (2n+2)}$$

Then

$$S_N = \sum_{i=1}^N a_n$$

Computing, we find

$$\begin{aligned} \frac{\pi-3}{4} &\approx 0.0353981635 \\ S_{10} &\approx 0.03535167962 \\ S_{100} &\approx 0.03539810274 \\ S_{500} &\approx 0.03539816290 \\ S_{1000} &\approx 0.03539816334 \end{aligned}$$

It appears that $S_N \rightarrow \frac{\pi-3}{4}$.

10. CAS

$$\frac{\pi^4}{90} = 1 + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \cdots$$

SOLUTION Write

$$S_N = \sum_{i=1}^N \frac{1}{i^4}$$

Computing, we find

$$\begin{aligned} \frac{\pi^4}{90} &\approx 1.082323234 \\ S(10) &\approx 1.082036583 \\ S(100) &\approx 1.082322905 \\ S(500) &\approx 1.082323231 \\ S(1000) &\approx 1.082323233 \end{aligned}$$

It appears that $S_N \rightarrow \frac{\pi^4}{90}$.

11. Calculate S_3 , S_4 , and S_5 and then find the sum of the telescoping series

$$S = \sum_{n=1}^{\infty} \left(\frac{1}{n+1} - \frac{1}{n+2} \right)$$

SOLUTION

$$\begin{aligned} S_3 &= \left(\frac{1}{2} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \left(\frac{1}{4} - \frac{1}{5} \right) = \frac{1}{2} - \frac{1}{5} = \frac{3}{10}; \\ S_4 &= S_3 + \left(\frac{1}{5} - \frac{1}{6} \right) = \frac{1}{2} - \frac{1}{6} = \frac{1}{3}; \\ S_5 &= S_4 + \left(\frac{1}{6} - \frac{1}{7} \right) = \frac{1}{2} - \frac{1}{7} = \frac{5}{14}. \end{aligned}$$

The general term in the sequence of partial sums is

$$S_N = \left(\frac{1}{2} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \left(\frac{1}{4} - \frac{1}{5} \right) + \cdots + \left(\frac{1}{N+1} - \frac{1}{N+2} \right) = \frac{1}{2} - \frac{1}{N+2};$$

thus,

$$S = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{N+2} \right) = \frac{1}{2}.$$

The sum of the telescoping series is therefore $\frac{1}{2}$.

12. Write $\sum_{n=3}^{\infty} \frac{1}{n(n-1)}$ as a telescoping series and find its sum.

SOLUTION By partial fraction decomposition

$$\frac{1}{n(n-1)} = \frac{1}{n-1} - \frac{1}{n},$$

so

$$\sum_{n=3}^{\infty} \frac{1}{n(n-1)} = \sum_{n=3}^{\infty} \left(\frac{1}{n-1} - \frac{1}{n} \right).$$

The general term in the sequence of partial sums for this series is

$$S_N = \left(\frac{1}{2} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \left(\frac{1}{4} - \frac{1}{5} \right) + \cdots + \left(\frac{1}{N-1} - \frac{1}{N} \right) = \frac{1}{2} - \frac{1}{N};$$

thus,

$$S = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{N} \right) = \frac{1}{2}.$$

13. Calculate S_3 , S_4 , and S_5 and then find the sum $S = \sum_{n=1}^{\infty} \frac{1}{4n^2-1}$ using the identity

$$\frac{1}{4n^2-1} = \frac{1}{2} \left(\frac{1}{2n-1} - \frac{1}{2n+1} \right)$$

SOLUTION

$$S_3 = \frac{1}{2} \left(\frac{1}{1} - \frac{1}{3} \right) + \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} \right) + \frac{1}{2} \left(\frac{1}{5} - \frac{1}{7} \right) = \frac{1}{2} \left(1 - \frac{1}{7} \right) = \frac{3}{7};$$

$$S_4 = S_3 + \frac{1}{2} \left(\frac{1}{7} - \frac{1}{9} \right) = \frac{1}{2} \left(1 - \frac{1}{9} \right) = \frac{4}{9};$$

$$S_5 = S_4 + \frac{1}{2} \left(\frac{1}{9} - \frac{1}{11} \right) = \frac{1}{2} \left(1 - \frac{1}{11} \right) = \frac{5}{11}.$$

The general term in the sequence of partial sums is

$$S_N = \frac{1}{2} \left(\frac{1}{1} - \frac{1}{3} \right) + \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} \right) + \frac{1}{2} \left(\frac{1}{5} - \frac{1}{7} \right) + \cdots + \frac{1}{2} \left(\frac{1}{2N-1} - \frac{1}{2N+1} \right) = \frac{1}{2} \left(1 - \frac{1}{2N+1} \right);$$

thus,

$$S = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \frac{1}{2} \left(1 - \frac{1}{2N+1} \right) = \frac{1}{2}.$$

14. Use partial fractions to rewrite $\sum_{n=1}^{\infty} \frac{1}{n(n+3)}$ as a telescoping series and find its sum.

SOLUTION By partial fraction decomposition

$$\frac{1}{n(n+3)} = \frac{A}{n} + \frac{B}{n+3};$$

clearing denominators gives

$$1 = A(n+3) + Bn.$$

Setting $n = 0$ yields $A = \frac{1}{3}$, while setting $n = -3$ yields $B = -\frac{1}{3}$. Thus,

$$\frac{1}{n(n+3)} = \frac{1}{3} \left(\frac{1}{n} - \frac{1}{n+3} \right),$$

and

$$\sum_{n=1}^{\infty} \frac{1}{n(n+3)} = \sum_{n=1}^{\infty} \frac{1}{3} \left(\frac{1}{n} - \frac{1}{n+3} \right).$$

The general term in the sequence of partial sums for the series on the right-hand side is

$$\begin{aligned} S_N &= \frac{1}{3} \left(1 - \frac{1}{4} \right) + \frac{1}{3} \left(\frac{1}{2} - \frac{1}{5} \right) + \frac{1}{3} \left(\frac{1}{3} - \frac{1}{6} \right) + \frac{1}{3} \left(\frac{1}{4} - \frac{1}{7} \right) + \frac{1}{3} \left(\frac{1}{5} - \frac{1}{8} \right) + \frac{1}{3} \left(\frac{1}{6} - \frac{1}{9} \right) \\ &\quad + \cdots + \frac{1}{3} \left(\frac{1}{N-2} - \frac{1}{N+1} \right) + \frac{1}{3} \left(\frac{1}{N-1} - \frac{1}{N+2} \right) + \frac{1}{3} \left(\frac{1}{N} - \frac{1}{N+3} \right) \\ &= \frac{1}{3} \left(1 + \frac{1}{2} + \frac{1}{3} \right) - \frac{1}{3} \left(\frac{1}{N+1} + \frac{1}{N+2} + \frac{1}{N+3} \right) = \frac{11}{18} - \frac{1}{3} \left(\frac{1}{N+1} + \frac{1}{N+2} + \frac{1}{N+3} \right). \end{aligned}$$

Thus,

$$\lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \left[\frac{11}{18} - \frac{1}{3} \left(\frac{1}{N+1} + \frac{1}{N+2} + \frac{1}{N+3} \right) \right] = \frac{11}{18},$$

and

$$\sum_{n=1}^{\infty} \frac{1}{n(n+3)} = \frac{11}{18}.$$

15. Find the sum of $\frac{1}{1 \cdot 3} + \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} + \cdots$.

SOLUTION We may write this sum as

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)(2n+1)} = \sum_{n=1}^{\infty} \frac{1}{2} \left(\frac{1}{2n-1} - \frac{1}{2n+1} \right).$$

The general term in the sequence of partial sums is

$$S_N = \frac{1}{2} \left(\frac{1}{1} - \frac{1}{3} \right) + \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} \right) + \frac{1}{2} \left(\frac{1}{5} - \frac{1}{7} \right) + \cdots + \frac{1}{2} \left(\frac{1}{2N-1} - \frac{1}{2N+1} \right) = \frac{1}{2} \left(1 - \frac{1}{2N+1} \right);$$

thus,

$$\lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \frac{1}{2} \left(1 - \frac{1}{2N+1} \right) = \frac{1}{2},$$

and

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)(2n+1)} = \frac{1}{2}.$$

16. Find a formula for the partial sum S_N of $\sum_{n=1}^{\infty} (-1)^{n-1}$ and show that the series diverges.

SOLUTION The partial sums of the series are:

$$\begin{aligned} S_1 &= (-1)^{1-1} = 1; \\ S_2 &= (-1)^0 + (-1)^1 = 1 - 1 = 0; \\ S_3 &= (-1)^0 + (-1)^1 + (-1)^2 = 1; \\ S_4 &= (-1)^0 + (-1)^1 + (-1)^2 + (-1)^3 = 0; \cdots \end{aligned}$$

In general,

$$S_N = \begin{cases} 1 & \text{if } N \text{ odd} \\ 0 & \text{if } N \text{ even} \end{cases}$$

Because the values of S_N alternate between 0 and 1, the sequence of partial sums diverges; this, in turn, implies that the series $\sum_{n=1}^{\infty} (-1)^{n-1}$ diverges.

In Exercises 17–22, use Theorem 3 to prove that the following series diverge.

$$17. \sum_{n=1}^{\infty} \frac{n}{10n+12}$$

SOLUTION The general term, $\frac{n}{10n+12}$, has limit

$$\lim_{n \rightarrow \infty} \frac{n}{10n+12} = \lim_{n \rightarrow \infty} \frac{1}{10+(12/n)} = \frac{1}{10}$$

Since the general term does not tend to zero, the series diverges.

$$18. \sum_{n=1}^{\infty} \frac{n}{\sqrt{n^2+1}}$$

SOLUTION The general term, $\frac{n}{\sqrt{n^2+1}}$, has limit

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2+1}} = \lim_{n \rightarrow \infty} \sqrt{\frac{n^2}{n^2+1}} = \lim_{n \rightarrow \infty} \sqrt{\frac{1}{1+(1/n^2)}} = 1$$

Since the general term does not tend to zero, the series diverges.

$$19. \frac{0}{1} - \frac{1}{2} + \frac{2}{3} - \frac{3}{4} + \cdots$$

SOLUTION The general term $a_n = (-1)^{n-1} \frac{n-1}{n}$ does not tend to zero. In fact, because $\lim_{n \rightarrow \infty} \frac{n-1}{n} = 1$, $\lim_{n \rightarrow \infty} a_n$ does not exist. By Theorem 3, we conclude that the given series diverges.

$$20. \sum_{n=1}^{\infty} (-1)^n n^2$$

SOLUTION The general term $a_n = (-1)^n n^2$ does not tend to zero. In fact, because $\lim_{n \rightarrow \infty} n^2 = \infty$, $\lim_{n \rightarrow \infty} a_n$ does not exist. By Theorem 3, we conclude that the given series diverges.

$$21. \cos \frac{1}{2} + \cos \frac{1}{3} + \cos \frac{1}{4} + \cdots$$

SOLUTION The general term $a_n = \cos \frac{1}{n+1}$ tends to 1, not zero. By Theorem 3, we conclude that the given series diverges.

$$22. \sum_{n=0}^{\infty} (\sqrt{4n^2+1} - n)$$

SOLUTION The general term of the series satisfies

$$\sqrt{4n^2+1} - n > \sqrt{4n^2} - n = n$$

Thus the general term tends to infinity. The series diverges by Theorem 2.

In Exercises 23–36, use the formula for the sum of a geometric series to find the sum or state that the series diverges.

$$23. \frac{1}{1} + \frac{1}{8} + \frac{1}{8^2} + \cdots$$

SOLUTION This is a geometric series with $c = 1$ and $r = \frac{1}{8}$, so its sum is

$$\frac{1}{1 - \frac{1}{8}} = \frac{1}{7/8} = \frac{8}{7}$$

$$24. \frac{4^3}{5^3} + \frac{4^4}{5^4} + \frac{4^5}{5^5} + \cdots$$

SOLUTION This is a geometric series with

$$c = \frac{4^3}{5^3} \quad \text{and} \quad r = \frac{4}{5}$$

so its sum is

$$\frac{c}{1-r} = \frac{4^3/5^3}{1 - \frac{4}{5}} = \frac{4^3}{5^3 - 4 \cdot 5^2} = \frac{64}{25}$$

$$25. \sum_{n=3}^{\infty} \left(\frac{3}{11}\right)^{-n}$$

SOLUTION Rewrite this series as

$$\sum_{n=3}^{\infty} \left(\frac{11}{3}\right)^n$$

This is a geometric series with $r = \frac{11}{3} > 1$, so it is divergent.

$$26. \sum_{n=2}^{\infty} \frac{7 \cdot (-3)^n}{5^n}$$

SOLUTION This is a geometric series with $c = 7$ and $r = -\frac{3}{5}$, starting at $n = 2$. Its sum is thus

$$\frac{cr^2}{1-r} = \frac{7 \cdot (9/25)}{1 - \frac{3}{5}} = \frac{63}{25} \cdot \frac{5}{8} = \frac{63}{40}$$

$$27. \sum_{n=-4}^{\infty} \left(-\frac{4}{9}\right)^n$$

SOLUTION This is a geometric series with $c = 1$ and $r = -\frac{4}{9}$, starting at $n = -4$. Its sum is thus

$$\frac{cr^{-4}}{1-r} = \frac{c}{r^4 - r^5} = \frac{1}{\frac{4^4}{9^4} + \frac{4^5}{9^5}} = \frac{9^5}{9 \cdot 4^4 + 4^5} = \frac{59,049}{3328}$$

$$28. \sum_{n=0}^{\infty} \left(\frac{\pi}{e}\right)^n$$

SOLUTION Since $\pi > e$, this is a geometric series with $r > 1$, so it diverges.

$$29. \sum_{n=1}^{\infty} e^{-n}$$

SOLUTION Rewrite the series as

$$\sum_{n=1}^{\infty} \left(\frac{1}{e}\right)^n$$

to recognize it as a geometric series with $c = \frac{1}{e}$ and $r = \frac{1}{e}$. Thus,

$$\sum_{n=1}^{\infty} e^{-n} = \frac{\frac{1}{e}}{1 - \frac{1}{e}} = \frac{1}{e-1}.$$

$$30. \sum_{n=2}^{\infty} e^{3-2n}$$

SOLUTION Rewrite the series as

$$\sum_{n=2}^{\infty} e^3 e^{-2n} = \sum_{n=2}^{\infty} e^3 \left(\frac{1}{e^2}\right)^n$$

to recognize it as a geometric series with $c = e^3 \left(\frac{1}{e^2}\right)^2 = \frac{1}{e}$ and $r = \frac{1}{e^2}$. Thus,

$$\sum_{n=2}^{\infty} e^{3-2n} = \frac{\frac{1}{e}}{1 - \frac{1}{e^2}} = \frac{e}{e^2 - 1}.$$

$$31. \sum_{n=0}^{\infty} \frac{8 + 2^n}{5^n}$$

SOLUTION Rewrite the series as

$$\sum_{n=0}^{\infty} \frac{8}{5^n} + \sum_{n=0}^{\infty} \frac{2^n}{5^n} = \sum_{n=0}^{\infty} 8 \cdot \left(\frac{1}{5}\right)^n + \sum_{n=0}^{\infty} \left(\frac{2}{5}\right)^n,$$

which is a sum of two geometric series. The first series has $c = 8 \left(\frac{1}{5}\right)^0 = 8$ and $r = \frac{1}{5}$; the second has $c = \left(\frac{2}{5}\right)^0 = 1$ and $r = \frac{2}{5}$. Thus,

$$\sum_{n=0}^{\infty} 8 \cdot \left(\frac{1}{5}\right)^n = \frac{8}{1 - \frac{1}{5}} = \frac{8}{\frac{4}{5}} = 10,$$

$$\sum_{n=0}^{\infty} \left(\frac{2}{5}\right)^n = \frac{1}{1 - \frac{2}{5}} = \frac{1}{\frac{3}{5}} = \frac{5}{3},$$

and

$$\sum_{n=0}^{\infty} \frac{8 + 2^n}{5^n} = 10 + \frac{5}{3} = \frac{35}{3}.$$

32.
$$\sum_{n=0}^{\infty} \frac{3(-2)^n - 5^n}{8^n}$$

SOLUTION Rewrite the series as

$$\sum_{n=0}^{\infty} \frac{3(-2)^n - 5^n}{8^n} = \sum_{n=0}^{\infty} \frac{3(-2)^n}{8^n} - \sum_{n=0}^{\infty} \frac{5^n}{8^n}$$

which is a difference of two geometric series. The first has $c = 3$ and $r = -\frac{1}{4}$; the second has $c = 1$ and $r = \frac{5}{8}$. Thus

$$\sum_{n=0}^{\infty} \frac{3(-2)^n}{8^n} = \frac{3}{1 + \frac{1}{4}} = \frac{12}{5}$$

$$\sum_{n=0}^{\infty} \frac{5^n}{8^n} = \frac{1}{1 - \frac{5}{8}} = \frac{8}{3}$$

so that

$$\sum_{n=0}^{\infty} \frac{3(-2)^n - 5^n}{8^n} = \frac{12}{5} - \frac{8}{3} = -\frac{4}{15}$$

33.
$$5 - \frac{5}{4} + \frac{5}{4^2} - \frac{5}{4^3} + \cdots$$

SOLUTION This is a geometric series with $c = 5$ and $r = -\frac{1}{4}$. Thus,

$$\sum_{n=0}^{\infty} 5 \cdot \left(-\frac{1}{4}\right)^n = \frac{5}{1 - \left(-\frac{1}{4}\right)} = \frac{5}{1 + \frac{1}{4}} = \frac{5}{\frac{5}{4}} = 4.$$

34.
$$\frac{2^3}{7} + \frac{2^4}{7^2} + \frac{2^5}{7^3} + \frac{2^6}{7^4} + \cdots$$

SOLUTION This is a geometric series with $c = \frac{8}{7}$ and $r = \frac{2}{7}$. Thus,

$$\sum_{n=0}^{\infty} \frac{8}{7} \cdot \left(\frac{2}{7}\right)^n = \frac{\frac{8}{7}}{1 - \frac{2}{7}} = \frac{\frac{8}{7}}{\frac{5}{7}} = \frac{8}{5}.$$

35.
$$\frac{7}{8} - \frac{49}{64} + \frac{343}{512} - \frac{2401}{4096} + \cdots$$

SOLUTION This is a geometric series with $c = \frac{7}{8}$ and $r = -\frac{7}{8}$. Thus,

$$\sum_{n=0}^{\infty} \frac{7}{8} \cdot \left(-\frac{7}{8}\right)^n = \frac{\frac{7}{8}}{1 - \left(-\frac{7}{8}\right)} = \frac{\frac{7}{8}}{\frac{15}{8}} = \frac{7}{15}.$$

$$36. \frac{25}{9} + \frac{5}{3} + 1 + \frac{3}{5} + \frac{9}{25} + \frac{27}{125} + \cdots$$

SOLUTION This appears to be a geometric series with

$$c = \frac{25}{9} \quad \text{and} \quad r = \frac{3}{5}$$

so its sum is

$$\frac{c}{1-r} = \frac{25/9}{1-3/5} = \frac{25}{9} \cdot \frac{5}{2} = \frac{125}{18}$$

37. Which of the following are *not* geometric series?

$$(a) \sum_{n=0}^{\infty} \frac{7^n}{29^n}$$

$$(b) \sum_{n=3}^{\infty} \frac{1}{n^4}$$

$$(c) \sum_{n=0}^{\infty} \frac{n^2}{2^n}$$

$$(d) \sum_{n=5}^{\infty} \pi^{-n}$$

SOLUTION

$$(a) \sum_{n=0}^{\infty} \frac{7^n}{29^n} = \sum_{n=0}^{\infty} \left(\frac{7}{29}\right)^n : \text{this is a geometric series with common ratio } r = \frac{7}{29}.$$

(b) The ratio between two successive terms is

$$\frac{a_{n+1}}{a_n} = \frac{\frac{1}{(n+1)^4}}{\frac{1}{n^4}} = \frac{n^4}{(n+1)^4} = \left(\frac{n}{n+1}\right)^4.$$

This ratio is not constant since it depends on n . Hence, the series $\sum_{n=3}^{\infty} \frac{1}{n^4}$ is not a geometric series.

(c) The ratio between two successive terms is

$$\frac{a_{n+1}}{a_n} = \frac{\frac{(n+1)^2}{2^{n+1}}}{\frac{n^2}{2^n}} = \frac{(n+1)^2}{n^2} \cdot \frac{2^n}{2^{n+1}} = \left(1 + \frac{1}{n}\right)^2 \cdot \frac{1}{2}.$$

This ratio is not constant since it depends on n . Hence, the series $\sum_{n=0}^{\infty} \frac{n^2}{2^n}$ is not a geometric series.

$$(d) \sum_{n=5}^{\infty} \pi^{-n} = \sum_{n=5}^{\infty} \left(\frac{1}{\pi}\right)^n : \text{this is a geometric series with common ratio } r = \frac{1}{\pi}.$$

38. Use the method of Example 8 to show that $\sum_{k=1}^{\infty} \frac{1}{k^{1/3}}$ diverges.

SOLUTION Each term in the N th partial sum is greater than or equal to $\frac{1}{N^{1/3}}$, hence:

$$S_N = \frac{1}{1^{1/3}} + \frac{1}{2^{1/3}} + \frac{1}{3^{1/3}} + \cdots + \frac{1}{N^{1/3}} \geq \frac{1}{N^{1/3}} + \frac{1}{N^{1/3}} + \frac{1}{N^{1/3}} + \cdots + \frac{1}{N^{1/3}} = N \cdot \frac{1}{N^{1/3}} = N^{2/3}.$$

Since $\lim_{N \rightarrow \infty} N^{2/3} = \infty$, it follows that

$$\lim_{N \rightarrow \infty} S_N = \infty.$$

Thus, the series $\sum_{k=1}^{\infty} \frac{1}{k^{1/3}}$ diverges.

39. Prove that if $\sum_{n=1}^{\infty} a_n$ converges and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} (a_n + b_n)$ diverges. *Hint:* If not, derive a contradiction by writing

$$\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} (a_n + b_n) - \sum_{n=1}^{\infty} a_n$$

SOLUTION Suppose to the contrary that $\sum_{n=1}^{\infty} a_n$ converges, $\sum_{n=1}^{\infty} b_n$ diverges, but $\sum_{n=1}^{\infty} (a_n + b_n)$ converges. Then by the Linearity of Infinite Series, we have

$$\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} (a_n + b_n) - \sum_{n=1}^{\infty} a_n$$


so that $\sum_{n=1}^{\infty} b_n$ converges, a contradiction.

40. Prove the divergence of $\sum_{n=0}^{\infty} \frac{9^n + 2^n}{5^n}$.

SOLUTION Note that this is the sum of two infinite series:

$$\sum_{n=0}^{\infty} \frac{9^n + 2^n}{5^n} = \sum_{n=0}^{\infty} \frac{9^n}{5^n} + \sum_{n=0}^{\infty} \frac{2^n}{5^n}$$

The first of these is a geometric series with $r = \frac{9}{5} > 1$, so diverges, while the second is a geometric series with $r = \frac{2}{5} < 1$, so converges. By the previous exercise, the sum of the two also diverges.

41.  Give a counterexample to show that each of the following statements is false.

(a) If the general term a_n tends to zero, then $\sum_{n=1}^{\infty} a_n = 0$.

(b) The N th partial sum of the infinite series defined by $\{a_n\}$ is a_N .

(c) If a_n tends to zero, then $\sum_{n=1}^{\infty} a_n$ converges.

(d) If a_n tends to L , then $\sum_{n=1}^{\infty} a_n = L$.

SOLUTION

(a) Let $a_n = 2^{-n}$. Then $\lim_{n \rightarrow \infty} a_n = 0$, but a_n is a geometric series with $c = 2^0 = 1$ and $r = 1/2$, so its sum is $\frac{1}{1 - (1/2)} = 2$.

(b) Let $a_n = 1$. Then the n th partial sum is $a_1 + a_2 + \cdots + a_n = n$ while $a_n = 1$.

(c) Let $a_n = \frac{1}{\sqrt{n}}$. An example in the text shows that while a_n tends to zero, the sum $\sum_{n=1}^{\infty} a_n$ does not converge.

(d) Let $a_n = 1$. Then clearly a_n tends to $L = 1$, while the series $\sum_{n=1}^{\infty} a_n$ obviously diverges.

42. Suppose that $S = \sum_{n=1}^{\infty} a_n$ is an infinite series with partial sum $S_N = 5 - \frac{2}{N^2}$.

(a) What are the values of $\sum_{n=1}^{10} a_n$ and $\sum_{n=5}^{16} a_n$?

(b) What is the value of a_3 ?

(c) Find a general formula for a_n .

(d) Find the sum $\sum_{n=1}^{\infty} a_n$.

SOLUTION

(a)

$$\sum_{n=1}^{10} a_n = S_{10} = 5 - \frac{2}{10^2} = \frac{249}{50};$$

$$\sum_{n=5}^{16} a_n = (a_1 + \cdots + a_{16}) - (a_1 + a_2 + a_3 + a_4) = S_{16} - S_4 = \left(5 - \frac{2}{16^2}\right) - \left(5 - \frac{2}{4^2}\right) = \frac{2}{16} - \frac{2}{256} = \frac{15}{128}.$$

(b)

$$a_3 = (a_1 + a_2 + a_3) - (a_1 + a_2) = S_3 - S_2 = \left(5 - \frac{2}{3^2}\right) - \left(5 - \frac{2}{2^2}\right) = \frac{1}{2} - \frac{2}{9} = \frac{5}{18}.$$

(c) Since $a_n = S_n - S_{n-1}$, we have:

$$\begin{aligned} a_n &= S_n - S_{n-1} = \left(5 - \frac{2}{n^2}\right) - \left(5 - \frac{2}{(n-1)^2}\right) = \frac{2}{(n-1)^2} - \frac{2}{n^2} \\ &= \frac{2(n^2 - (n-1)^2)}{(n(n-1))^2} = \frac{2(n^2 - n^2 + 2n - 1)}{(n(n-1))^2} = \frac{2(2n-1)}{n^2(n-1)^2}. \end{aligned}$$

(d) The sum $\sum_{n=1}^{\infty} a_n$ is the limit of the sequence of partial sums $\{S_N\}$. Hence:

$$\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \left(5 - \frac{2}{N^2}\right) = 5.$$

43. Compute the total area of the (infinitely many) triangles in Figure 4.




FIGURE 4

SOLUTION The area of a triangle with base B and height H is $A = \frac{1}{2}BH$. Because all of the triangles in Figure 4 have height $\frac{1}{2}$, the area of each triangle equals one-quarter of the base. Now, for $n \geq 0$, the n th triangle has a base which extends from $x = \frac{1}{2^{n+1}}$ to $x = \frac{1}{2^n}$. Thus,

$$B = \frac{1}{2^n} - \frac{1}{2^{n+1}} = \frac{1}{2^{n+1}} \quad \text{and} \quad A = \frac{1}{4}B = \frac{1}{2^{n+3}}.$$

The total area of the triangles is then given by the geometric series

$$\sum_{n=0}^{\infty} \frac{1}{2^{n+3}} = \sum_{n=0}^{\infty} \frac{1}{8} \left(\frac{1}{2}\right)^n = \frac{\frac{1}{8}}{1 - \frac{1}{2}} = \frac{1}{4}.$$

44. The winner of a lottery receives m dollars at the end of each year for N years. The present value (PV) of this prize in today's dollars is $PV = \sum_{i=1}^N m(1+r)^{-i}$, where r is the interest rate. Calculate PV if $m = \$50,000$, $r = 0.06$, and $N = 20$. What is PV if $N = \infty$?

SOLUTION For the given values r , m and N , we have

$$PV = \sum_{i=1}^{20} 50,000(1+0.06)^{-i} = \sum_{i=1}^{20} 50,000 \left(\frac{50}{53}\right)^i = 50,000 \frac{1 - \left(\frac{50}{53}\right)^{21}}{1 - \frac{50}{53}} = \$623,496.06.$$

If we extend the payments forever, then $N = \infty$ and

$$PV = \sum_{i=1}^{\infty} 50,000(1+0.06)^{-i} = \sum_{i=1}^{\infty} 50,000 \left(\frac{50}{53}\right)^i = \frac{50,000 \left(\frac{50}{53}\right)}{1 - \frac{50}{53}} = \$833,333.33.$$

45. Find the total length of the infinite zigzag path in Figure 5 (each zag occurs at an angle of $\frac{\pi}{4}$).




FIGURE 5

SOLUTION Because the angle at the lower left in Figure 5 has measure $\frac{\pi}{4}$ and each zag in the path occurs at an angle of $\frac{\pi}{4}$, every triangle in the figure is an isosceles right triangle. Accordingly, the length of each new segment in the path is $\frac{1}{\sqrt{2}}$ times the length of the previous segment. Since the first segment has length 1, the total length of the path is

$$\sum_{n=0}^{\infty} \left(\frac{1}{\sqrt{2}}\right)^n = \frac{1}{1 - \frac{1}{\sqrt{2}}} = \frac{\sqrt{2}}{\sqrt{2} - 1} = 2 + \sqrt{2}.$$

46. Evaluate $\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)}$. *Hint:* Find constants A , B , and C such that

$$\frac{1}{n(n+1)(n+2)} = \frac{A}{n} + \frac{B}{n+1} + \frac{C}{n+2}$$

SOLUTION By partial fraction decomposition

$$\frac{1}{n(n+1)(n+2)} = \frac{A}{n} + \frac{B}{n+1} + \frac{C}{n+2};$$

clearing denominators then gives

$$1 = A(n+1)(n+2) + Bn(n+2) + Cn(n+1).$$

Setting $n = 0$ now yields $A = \frac{1}{2}$, while setting $n = -1$ yields $B = -1$ and setting $n = -2$ yields $C = \frac{1}{2}$. Thus,

$$\frac{1}{n(n+1)(n+2)} = \frac{\frac{1}{2}}{n} - \frac{1}{n+1} + \frac{\frac{1}{2}}{n+2} = \frac{1}{2} \left(\frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2} \right),$$

and

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)} = \sum_{n=1}^{\infty} \frac{1}{2} \left(\frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2} \right).$$

The general term of the sequence of partial sums for the series on the right-hand side is

$$\begin{aligned} S_N &= \frac{1}{2} \left(1 - \frac{2}{2} + \frac{1}{3} \right) + \frac{1}{2} \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) + \frac{1}{2} \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5} \right) + \frac{1}{2} \left(\frac{1}{4} - \frac{2}{5} + \frac{1}{6} \right) + \frac{1}{2} \left(\frac{1}{5} - \frac{2}{6} + \frac{1}{7} \right) \\ &\quad + \cdots + \frac{1}{2} \left(\frac{1}{N-2} - \frac{2}{N-1} + \frac{1}{N} \right) + \frac{1}{2} \left(\frac{1}{N-1} - \frac{2}{N} + \frac{1}{N+1} \right) + \frac{1}{2} \left(\frac{1}{N} - \frac{2}{N+1} + \frac{1}{N+2} \right) \\ &= \frac{1}{2} \left(\frac{1}{2} - \frac{1}{N+1} + \frac{1}{N+2} \right). \end{aligned}$$

Thus,

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)} = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \frac{1}{2} \left(\frac{1}{2} - \frac{1}{N+1} + \frac{1}{N+2} \right) = \frac{1}{4}.$$

47. Show that if a is a positive integer, then

$$\sum_{n=1}^{\infty} \frac{1}{n(n+a)} = \frac{1}{a} \left(1 + \frac{1}{2} + \cdots + \frac{1}{a} \right)$$

SOLUTION By partial fraction decomposition

$$\frac{1}{n(n+a)} = \frac{A}{n} + \frac{B}{n+a};$$

clearing the denominators gives

$$1 = A(n+a) + Bn.$$

Setting $n = 0$ then yields $A = \frac{1}{a}$, while setting $n = -a$ yields $B = -\frac{1}{a}$. Thus,

$$\frac{1}{n(n+a)} = \frac{\frac{1}{a}}{n} - \frac{\frac{1}{a}}{n+a} = \frac{1}{a} \left(\frac{1}{n} - \frac{1}{n+a} \right),$$

and

$$\sum_{n=1}^{\infty} \frac{1}{n(n+a)} = \sum_{n=1}^{\infty} \frac{1}{a} \left(\frac{1}{n} - \frac{1}{n+a} \right).$$

For $N > a$, the N th partial sum is


$$S_N = \frac{1}{a} \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{a} \right) - \frac{1}{a} \left(\frac{1}{N+1} + \frac{1}{N+2} + \frac{1}{N+3} + \cdots + \frac{1}{N+a} \right).$$

Thus,

$$\sum_{n=1}^{\infty} \frac{1}{n(n+a)} = \lim_{N \rightarrow \infty} S_N = \frac{1}{a} \left(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{a} \right).$$

48. A ball dropped from a height of 10 ft begins to bounce. Each time it strikes the ground, it returns to two-thirds of its previous height. What is the total distance traveled by the ball if it bounces infinitely many times?

SOLUTION The distance traveled by the ball is shown in the accompanying figure:



The total distance d traveled by the ball is given by the following infinite sum:

$$d = h + 2 \cdot \frac{2}{3}h + 2 \cdot \left(\frac{2}{3}\right)^2 h + 2 \cdot \left(\frac{2}{3}\right)^3 h + \cdots = h + 2h \left(\frac{2}{3} + \left(\frac{2}{3}\right)^2 + \left(\frac{2}{3}\right)^3 + \cdots \right) = h + 2h \sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n.$$

We use the formula for the sum of a geometric series to compute the sum of the resulting series:

$$d = h + 2h \cdot \frac{\left(\frac{2}{3}\right)^1}{1 - \frac{2}{3}} = h + 2h(2) = 5h.$$

With $h = 10$ feet, it follows that the total distance traveled by the ball is 50 feet.

49. Let $\{b_n\}$ be a sequence and let $a_n = b_n - b_{n-1}$. Show that $\sum_{n=1}^{\infty} a_n$ converges if and only if $\lim_{n \rightarrow \infty} b_n$ exists.

SOLUTION Let $a_n = b_n - b_{n-1}$. The general term in the sequence of partial sums for the series $\sum_{n=1}^{\infty} a_n$ is then

$$S_N = (b_1 - b_0) + (b_2 - b_1) + (b_3 - b_2) + \cdots + (b_N - b_{N-1}) = b_N - b_0.$$

Now, if $\lim_{N \rightarrow \infty} b_N$ exists, then so does $\lim_{N \rightarrow \infty} S_N$ and $\sum_{n=1}^{\infty} a_n$ converges. On the other hand, if $\sum_{n=1}^{\infty} a_n$ converges, then

$\lim_{N \rightarrow \infty} S_N$ exists, which implies that $\lim_{N \rightarrow \infty} b_N$ also exists. Thus, $\sum_{n=1}^{\infty} a_n$ converges if and only if $\lim_{n \rightarrow \infty} b_n$ exists.

50. Assumptions Matter Show, by giving counterexamples, that the assertions of Theorem 1 are not valid if the series

$\sum_{n=0}^{\infty} a_n$ and $\sum_{n=0}^{\infty} b_n$ are not convergent.

SOLUTION Let $a_n = 2^{-n} - 2^n$ and $b_n = 2^n$. Then, both

$$\sum_{n=0}^{\infty} a_n \quad \text{and} \quad \sum_{n=0}^{\infty} b_n$$

diverge, so the sum

$$\sum_{n=0}^{\infty} a_n + \sum_{n=0}^{\infty} b_n$$

is not defined. However,

$$\sum_{n=0}^{\infty} (a_n + b_n) = \sum_{n=0}^{\infty} ((2^{-n} - 2^n) + 2^n) = \sum_{n=0}^{\infty} 2^{-n} = 1.$$

Further Insights and Challenges

Exercises 51–53 use the formula

$$1 + r + r^2 + \cdots + r^{N-1} = \frac{1 - r^N}{1 - r} \quad \boxed{7}$$

51. Professor George Andrews of Pennsylvania State University observed that we can use Eq. (7) to calculate the derivative of $f(x) = x^N$ (for $N \geq 0$). Assume that $a \neq 0$ and let $x = ra$. Show that

$$f'(a) = \lim_{x \rightarrow a} \frac{x^N - a^N}{x - a} = a^{N-1} \lim_{r \rightarrow 1} \frac{r^N - 1}{r - 1}$$

and evaluate the limit.

SOLUTION According to the definition of derivative of $f(x)$ at $x = a$

$$f'(a) = \lim_{x \rightarrow a} \frac{x^N - a^N}{x - a}.$$

Now, let $x = ra$. Then $x \rightarrow a$ if and only if $r \rightarrow 1$, and

$$f'(a) = \lim_{x \rightarrow a} \frac{x^N - a^N}{x - a} = \lim_{r \rightarrow 1} \frac{(ra)^N - a^N}{ra - a} = \lim_{r \rightarrow 1} \frac{a^N (r^N - 1)}{a(r - 1)} = a^{N-1} \lim_{r \rightarrow 1} \frac{r^N - 1}{r - 1}.$$

By Eq. (7) for a geometric sum,

$$\frac{1 - r^N}{1 - r} = \frac{r^N - 1}{r - 1} = 1 + r + r^2 + \cdots + r^{N-1},$$

so

$$\lim_{r \rightarrow 1} \frac{r^N - 1}{r - 1} = \lim_{r \rightarrow 1} (1 + r + r^2 + \cdots + r^{N-1}) = 1 + 1 + 1^2 + \cdots + 1^{N-1} = N.$$

Therefore, $f'(a) = a^{N-1} \cdot N = Na^{N-1}$

52. Pierre de Fermat used geometric series to compute the area under the graph of $f(x) = x^N$ over $[0, A]$. For $0 < r < 1$, let $F(r)$ be the sum of the areas of the infinitely many right-endpoint rectangles with endpoints Ar^n , as in Figure 6. As r tends to 1, the rectangles become narrower and $F(r)$ tends to the area under the graph.

(a) Show that $F(r) = A^{N+1} \frac{1 - r}{1 - r^{N+1}}$.

(b) Use Eq. (7) to evaluate $\int_0^A x^N dx = \lim_{r \rightarrow 1} F(r)$.




FIGURE 6

SOLUTION

(a) The area of the rectangle whose base extends from $x = r^n A$ to $x = r^{n-1} A$ is

$$(r^{n-1} A)^N (r^{n-1} A - r^n A).$$

Hence, $F(r)$ is the sum

$$\begin{aligned} F(r) &= \sum_{n=1}^{\infty} (r^{n-1}A)^N (r^{n-1}A - r^nA) = \sum_{n=1}^{\infty} r^{(n-1)N} r^{n-1} (1-r)A^{N+1} = A^{N+1}(1-r) \sum_{n=1}^{\infty} r^{nN-N+n-1} \\ &= \frac{A^{N+1}(1-r)}{r^{N+1}} \sum_{n=1}^{\infty} (r^{N+1})^n = \frac{A^{N+1}(1-r)}{r^{N+1}} \cdot \frac{r^{N+1}}{1-r^{N+1}} = A^{N+1} \frac{1-r}{1-r^{N+1}}. \end{aligned}$$

(b) Using the result from part (a) and Eq. (7) from Exercise 51,

$$\int_0^A x^N dx = \lim_{r \rightarrow 1} F(r) = A^{N+1} \lim_{r \rightarrow 1} \frac{1-r}{1-r^{N+1}} = A^{N+1} \lim_{r \rightarrow 1} \frac{1}{1+r+r^2+\dots+r^N} = A^{N+1} \cdot \frac{1}{N+1} = \frac{A^{N+1}}{N+1}.$$

53. Verify the Gregory–Leibniz formula as follows.

(a) Set $r = -x^2$ in Eq. (7) and rearrange to show that

$$\frac{1}{1+x^2} = 1 - x^2 + x^4 - \dots + (-1)^{N-1} x^{2N-2} + \frac{(-1)^N x^{2N}}{1+x^2}$$

(b) Show, by integrating over $[0, 1]$, that

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots + \frac{(-1)^{N-1}}{2N-1} + (-1)^N \int_0^1 \frac{x^{2N}}{1+x^2} dx$$

(c) Use the Comparison Theorem for integrals to prove that

$$0 \leq \int_0^1 \frac{x^{2N}}{1+x^2} dx \leq \frac{1}{2N+1}$$

Hint: Observe that the integrand is $\leq x^{2N}$.

(d) Prove that

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \dots$$

Hint: Use (b) and (c) to show that the partial sums S_N of satisfy $|S_N - \frac{\pi}{4}| \leq \frac{1}{2N+1}$, and thereby conclude that $\lim_{N \rightarrow \infty} S_N = \frac{\pi}{4}$.

SOLUTION

(a) Start with Eq. (7), and substitute $-x^2$ for r :

$$\begin{aligned} 1 + r + r^2 + \dots + r^{N-1} &= \frac{1-r^N}{1-r} \\ 1 - x^2 + x^4 + \dots + (-1)^{N-1} x^{2N-2} &= \frac{1 - (-1)^N x^{2N}}{1 - (-x^2)} \\ 1 - x^2 + x^4 + \dots + (-1)^{N-1} x^{2N-2} &= \frac{1}{1+x^2} - \frac{(-1)^N x^{2N}}{1+x^2} \\ \frac{1}{1+x^2} &= 1 - x^2 + x^4 + \dots + (-1)^{N-1} x^{2N-2} + \frac{(-1)^N x^{2N}}{1+x^2} \end{aligned}$$

(b) The integrals of both sides must be equal. Now,

$$\int_0^1 \frac{1}{1+x^2} dx = \tan^{-1} x \Big|_0^1 = \tan^{-1} 1 - \tan^{-1} 0 = \frac{\pi}{4}$$

while

$$\begin{aligned} &\int_0^1 \left(1 - x^2 + x^4 + \dots + (-1)^{N-1} x^{2N-2} + \frac{(-1)^N x^{2N}}{1+x^2} \right) dx \\ &= \left(x - \frac{1}{3} x^3 + \frac{1}{5} x^5 + \dots + (-1)^{N-1} \frac{1}{2N-1} x^{2N-1} \right) + (-1)^N \int_0^1 \frac{x^{2N}}{1+x^2} dx \\ &= 1 - \frac{1}{3} + \frac{1}{5} + \dots + (-1)^{N-1} \frac{1}{2N-1} + (-1)^N \int_0^1 \frac{x^{2N}}{1+x^2} dx \end{aligned}$$

(c) Note that for $x \in [0, 1]$, we have $1 + x^2 \geq 1$, so that

$$0 \leq \frac{x^{2N}}{1+x^2} \leq x^{2N}$$

By the Comparison Theorem for integrals, we then see that

$$0 \leq \int_0^1 \frac{x^{2N} dx}{1+x^2} \leq \int_0^1 x^{2N} dx = \frac{1}{2N+1} x^{2N+1} \Big|_0^1 = \frac{1}{2N+1}$$

(d) Write

$$a_n = (-1)^n \frac{1}{2n-1}, \quad n \geq 1$$

and let S_N be the partial sums. Then

$$\left| S_N - \frac{\pi}{4} \right| = \left| (-1)^N \int_0^1 \frac{x^{2N} dx}{1+x^2} \right| = \int_0^1 \frac{x^{2N} dx}{1+x^2} \leq \frac{1}{2N+1}$$

Thus $\lim_{N \rightarrow \infty} S_N = \frac{\pi}{4}$ so that

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \dots$$

54. Cantor's Disappearing Table (following Larry Knop of Hamilton College) Take a table of length L (Figure 7). At stage 1, remove the section of length $L/4$ centered at the midpoint. Two sections remain, each with length less than $L/2$. At stage 2, remove sections of length $L/4^2$ from each of these two sections (this stage removes $L/8$ of the table). Now four sections remain, each of length less than $L/4$. At stage 3, remove the four central sections of length $L/4^3$, etc.

(a) Show that at the N th stage, each remaining section has length less than $L/2^N$ and that the total amount of table removed is

$$L \left(\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots + \frac{1}{2^{N+1}} \right)$$

(b) Show that in the limit as $N \rightarrow \infty$, precisely one-half of the table remains.

This result is curious, because there are no nonzero intervals of table left (at each stage, the remaining sections have a length less than $L/2^N$). So the table has “disappeared.” However, we can place any object longer than $L/4$ on the table. It will not fall through because it will not fit through any of the removed sections.




FIGURE 7

SOLUTION

(a) After the N th stage, the total amount of table that has been removed is

$$\frac{L}{4} + \frac{2L}{4^2} + \frac{4L}{4^3} + \dots + \frac{2^{N-1}L}{4^N} = L \left(\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots + \frac{2^{N-1}}{2^{2N}} \right) = L \left(\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots + \frac{1}{2^{N+1}} \right)$$

At the first stage ($N = 1$), there are two remaining sections each of length

$$\frac{L - \frac{L}{4}}{2} = \frac{3L}{8} < \frac{L}{2}.$$

Suppose that at the K th stage, each of the 2^K remaining sections has length less than $\frac{L}{2^K}$. The $(K+1)$ st stage is obtained

by removing the section of length $\frac{L}{4^{K+1}}$ centered at the midpoint of each segment in the K th stage. Let a_k and a_{K+1} , respectively, denote the length of each segment in the K th and $(K+1)$ st stage. Then,

$$a_{K+1} = \frac{a_K - \frac{L}{4^{K+1}}}{2} < \frac{\frac{L}{2^K} - \frac{L}{4^{K+1}}}{2} = \frac{L}{2^K} \left(\frac{1 - \frac{1}{2^{K+2}}}{2} \right) < \frac{L}{2^K} \cdot \frac{1}{2} = \frac{L}{2^{K+1}}.$$

Thus, by mathematical induction, each remaining section at the N th stage has length less than $\frac{L}{2^N}$.

(b) From part (a), we know that after N stages, the amount of the table that has been removed is

$$L \left(\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \cdots + \frac{1}{2^{N+1}} \right) = \sum_{n=1}^N \frac{1}{2^{n+1}}.$$

As $N \rightarrow \infty$, the amount of the table that has been removed becomes a geometric series whose sum is

$$L \sum_{n=1}^{\infty} \frac{1}{2} \left(\frac{1}{2} \right)^n = L \frac{\frac{1}{4}}{1 - \frac{1}{2}} = \frac{1}{2}L.$$

Thus, the amount of table that remains is $L - \frac{1}{2}L = \frac{1}{2}L$.

55. The Koch snowflake (described in 1904 by Swedish mathematician Helge von Koch) is an infinitely jagged “fractal” curve obtained as a limit of polygonal curves (it is continuous but has no tangent line at any point). Begin with an equilateral triangle (stage 0) and produce stage 1 by replacing each edge with four edges of one-third the length, arranged as in Figure 8. Continue the process: At the n th stage, replace each edge with four edges of one-third the length.

(a) Show that the perimeter P_n of the polygon at the n th stage satisfies $P_n = \frac{4}{3}P_{n-1}$. Prove that $\lim_{n \rightarrow \infty} P_n = \infty$. The snowflake has infinite length.

(b) Let A_0 be the area of the original equilateral triangle. Show that $(3)4^{n-1}$ new triangles are added at the n th stage, each with area $A_0/9^n$ (for $n \geq 1$). Show that the total area of the Koch snowflake is $\frac{8}{5}A_0$.




FIGURE 8

SOLUTION

(a) Each edge of the polygon at the $(n-1)$ st stage is replaced by four edges of one-third the length; hence the perimeter of the polygon at the n th stage is $\frac{4}{3}$ times the perimeter of the polygon at the $(n-1)$ th stage. That is, $P_n = \frac{4}{3}P_{n-1}$. Thus,

$$P_1 = \frac{4}{3}P_0; \quad P_2 = \frac{4}{3}P_1 = \left(\frac{4}{3}\right)^2 P_0, \quad P_3 = \frac{4}{3}P_2 = \left(\frac{4}{3}\right)^3 P_0,$$

and, in general, $P_n = \left(\frac{4}{3}\right)^n P_0$. As $n \rightarrow \infty$, it follows that

$$\lim_{n \rightarrow \infty} P_n = P_0 \lim_{n \rightarrow \infty} \left(\frac{4}{3}\right)^n = \infty.$$

(b) When each edge is replaced by four edges of one-third the length, one new triangle is created. At the $(n-1)$ st stage, there are $3 \cdot 4^{n-1}$ edges in the snowflake, so $3 \cdot 4^{n-1}$ new triangles are generated at the n th stage. Because the area of an equilateral triangle is proportional to the square of its side length and the side length for each new triangle is one-third the side length of triangles from the previous stage, it follows that the area of the triangles added at each stage is reduced by a factor of $\frac{1}{9}$ from the area of the triangles added at the previous stage. Thus, each triangle added at the n th stage has an area of $A_0/9^n$. This means that the n th stage contributes

$$3 \cdot 4^{n-1} \cdot \frac{A_0}{9^n} = \frac{3}{4}A_0 \left(\frac{4}{9}\right)^n$$

to the area of the snowflake. The total area is therefore

$$A = A_0 + \frac{3}{4}A_0 \sum_{n=1}^{\infty} \left(\frac{4}{9}\right)^n = A_0 + \frac{3}{4}A_0 \frac{\frac{4}{9}}{1 - \frac{4}{9}} = A_0 + \frac{3}{4}A_0 \cdot \frac{4}{5} = \frac{8}{5}A_0.$$

10.3 Convergence of Series with Positive Terms

Preliminary Questions

1. Let $S = \sum_{n=1}^{\infty} a_n$. If the partial sums S_N are increasing, then (choose the correct conclusion):

- (a) $\{a_n\}$ is an increasing sequence.
 (b) $\{a_n\}$ is a positive sequence.

SOLUTION The correct response is **(b)**. Recall that $S_N = a_1 + a_2 + a_3 + \cdots + a_N$; thus, $S_N - S_{N-1} = a_N$. If S_N is increasing, then $S_N - S_{N-1} \geq 0$. It then follows that $a_N \geq 0$; that is, $\{a_n\}$ is a positive sequence.

2. What are the hypotheses of the Integral Test?

SOLUTION The hypotheses for the Integral Test are: A function $f(x)$ such that $a_n = f(n)$ must be positive, decreasing, and continuous for $x \geq 1$.

3. Which test would you use to determine whether $\sum_{n=1}^{\infty} n^{-3.2}$ converges?

SOLUTION Because $n^{-3.2} = \frac{1}{n^{3.2}}$, we see that the indicated series is a p -series with $p = 3.2 > 1$. Therefore, the series converges.

4. Which test would you use to determine whether $\sum_{n=1}^{\infty} \frac{1}{2^n + \sqrt{n}}$ converges?

SOLUTION Because

$$\frac{1}{2^n + \sqrt{n}} < \frac{1}{2^n} = \left(\frac{1}{2}\right)^n,$$

and

$$\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n$$

is a convergent geometric series, the comparison test would be an appropriate choice to establish that the given series converges.

5. Ralph hopes to investigate the convergence of $\sum_{n=1}^{\infty} \frac{e^{-n}}{n}$ by comparing it with $\sum_{n=1}^{\infty} \frac{1}{n}$. Is Ralph on the right track?

SOLUTION No, Ralph is not on the right track. For $n \geq 1$,

$$\frac{e^{-n}}{n} < \frac{1}{n};$$

however, $\sum_{n=1}^{\infty} \frac{1}{n}$ is a divergent series. The Comparison Test therefore does not allow us to draw a conclusion about the

convergence or divergence of the series $\sum_{n=1}^{\infty} \frac{e^{-n}}{n}$.

Exercises

In Exercises 1–14, use the Integral Test to determine whether the infinite series is convergent.

1. $\sum_{n=1}^{\infty} \frac{1}{n^4}$

SOLUTION Let $f(x) = \frac{1}{x^4}$. This function is continuous, positive and decreasing on the interval $x \geq 1$, so the Integral Test applies. Moreover,

$$\int_1^{\infty} \frac{dx}{x^4} = \lim_{R \rightarrow \infty} \int_1^R x^{-4} dx = -\frac{1}{3} \lim_{R \rightarrow \infty} \left(\frac{1}{R^3} - 1\right) = \frac{1}{3}.$$

The integral converges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{n^4}$ also converges.

2. $\sum_{n=1}^{\infty} \frac{1}{n+3}$

SOLUTION Let $f(x) = \frac{1}{x+3}$. This function is continuous, positive and decreasing on the interval $x \geq 1$, so the Integral Test applies. Moreover,

$$\int_1^{\infty} \frac{dx}{x+3} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x+3} = \lim_{R \rightarrow \infty} (\ln(R+3) - \ln 4) = \infty.$$

The integral diverges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{n+3}$ also diverges.

$$3. \sum_{n=1}^{\infty} n^{-1/3}$$

SOLUTION Let $f(x) = x^{-1/3} = \frac{1}{\sqrt[3]{x}}$. This function is continuous, positive and decreasing on the interval $x \geq 1$, so the Integral Test applies. Moreover,

$$\int_1^{\infty} x^{-1/3} dx = \lim_{R \rightarrow \infty} \int_1^R x^{-1/3} dx = \frac{3}{2} \lim_{R \rightarrow \infty} (R^{2/3} - 1) = \infty.$$

The integral diverges; hence, the series $\sum_{n=1}^{\infty} n^{-1/3}$ also diverges.

$$4. \sum_{n=5}^{\infty} \frac{1}{\sqrt{n-4}}$$

SOLUTION Let $f(x) = \frac{1}{\sqrt{x-4}}$. This function is continuous, positive and decreasing on the interval $x \geq 5$, so the Integral Test applies. Moreover,

$$\int_5^{\infty} \frac{dx}{\sqrt{x-4}} = \lim_{R \rightarrow \infty} \int_5^R \frac{dx}{\sqrt{x-4}} = 2 \lim_{R \rightarrow \infty} (\sqrt{R-4} - 1) = \infty.$$

The integral diverges; hence, the series $\sum_{n=5}^{\infty} \frac{1}{\sqrt{n-4}}$ also diverges.

$$5. \sum_{n=25}^{\infty} \frac{n^2}{(n^3+9)^{5/2}}$$

SOLUTION Let $f(x) = \frac{x^2}{(x^3+9)^{5/2}}$. This function is positive and continuous for $x \geq 25$. Moreover, because

$$f'(x) = \frac{2x(x^3+9)^{5/2} - x^2 \cdot \frac{5}{2}(x^3+9)^{3/2} \cdot 3x^2}{(x^3+9)^5} = \frac{x(36-11x^3)}{2(x^3+9)^{7/2}},$$

we see that $f'(x) < 0$ for $x \geq 25$, so f is decreasing on the interval $x \geq 25$. The Integral Test therefore applies. To evaluate the improper integral, we use the substitution $u = x^3 + 9$, $du = 3x^2 dx$. We then find

$$\begin{aligned} \int_{25}^{\infty} \frac{x^2}{(x^3+9)^{5/2}} dx &= \lim_{R \rightarrow \infty} \int_{25}^R \frac{x^2}{(x^3+9)^{5/2}} dx = \frac{1}{3} \lim_{R \rightarrow \infty} \int_{15634}^{R^3+9} \frac{du}{u^{5/2}} \\ &= -\frac{2}{9} \lim_{R \rightarrow \infty} \left(\frac{1}{(R^3+9)^{3/2}} - \frac{1}{15634^{3/2}} \right) = \frac{2}{9 \cdot 15634^{3/2}}. \end{aligned}$$

The integral converges; hence, the series $\sum_{n=25}^{\infty} \frac{n^2}{(n^3+9)^{5/2}}$ also converges.

$$6. \sum_{n=1}^{\infty} \frac{n}{(n^2+1)^{3/5}}$$

SOLUTION Let $f(x) = \frac{x}{(x^2+1)^{3/5}}$. Because

$$f'(x) = \frac{(x^2+1)^{3/5} - x \cdot \frac{6}{5}x(x^2+1)^{-2/5}}{(x^2+1)^6} = \frac{1 - \frac{1}{5}x^2}{(x^2+1)^{8/5}},$$

we see that $f'(x) < 0$ for $x > \sqrt{5} \approx 2.236$. We conclude that f is decreasing on the interval $x \geq 3$. Since f is also positive and continuous on this interval, the Integral Test can be applied. To evaluate the improper integral, we make the substitution $u = x^2 + 1$, $du = 2x dx$. This gives

$$\int_3^{\infty} \frac{x}{(x^2+1)^{3/5}} dx = \lim_{R \rightarrow \infty} \int_3^R \frac{x}{(x^2+1)^{3/5}} dx = \frac{1}{2} \lim_{R \rightarrow \infty} \int_{10}^{R^2+1} \frac{du}{u^{3/5}} = \frac{5}{4} \lim_{R \rightarrow \infty} ((R^2+1)^{2/5} - 10^{2/5}) = \infty.$$

The integral diverges; therefore, the series $\sum_{n=3}^{\infty} \frac{n}{(n^2+1)^{3/5}}$ also diverges. Since the divergence of the series is not affected by adding the finite sum $\sum_{n=1}^2 \frac{n}{(n^2+1)^{3/5}}$, the series $\sum_{n=1}^{\infty} \frac{n}{(n^2+1)^{3/5}}$ also diverges.

$$7. \sum_{n=1}^{\infty} \frac{1}{n^2+1}$$

SOLUTION Let $f(x) = \frac{1}{x^2+1}$. This function is positive, decreasing and continuous on the interval $x \geq 1$, hence the Integral Test applies. Moreover,

$$\int_1^{\infty} \frac{dx}{x^2+1} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x^2+1} = \lim_{R \rightarrow \infty} \left(\tan^{-1} R - \frac{\pi}{4} \right) = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}.$$

The integral converges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{n^2+1}$ also converges.

$$8. \sum_{n=4}^{\infty} \frac{1}{n^2-1}$$

SOLUTION Let $f(x) = \frac{1}{x^2-1}$. This function is continuous, positive and decreasing on the interval $x \geq 4$; therefore, the Integral Test applies. We compute the improper integral using partial fractions:

$$\begin{aligned} \int_4^{\infty} \frac{dx}{x^2-1} &= \lim_{R \rightarrow \infty} \int_4^R \left(\frac{\frac{1}{2}}{x-1} - \frac{\frac{1}{2}}{x+1} \right) dx = \frac{1}{2} \lim_{R \rightarrow \infty} \ln \frac{x-1}{x+1} \Big|_4^R = \frac{1}{2} \lim_{R \rightarrow \infty} \left(\ln \frac{R-1}{R+1} - \ln \frac{3}{5} \right) \\ &= \frac{1}{2} \left(\ln 1 - \ln \frac{3}{5} \right) = -\frac{1}{2} \ln \frac{3}{5}. \end{aligned}$$

The integral converges; hence, the series $\sum_{n=4}^{\infty} \frac{1}{n^2-1}$ also converges.

$$9. \sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

SOLUTION Let $f(x) = \frac{1}{x(x+1)}$. This function is positive, continuous and decreasing on the interval $x \geq 1$, so the Integral Test applies. We compute the improper integral using partial fractions:

$$\int_1^{\infty} \frac{dx}{x(x+1)} = \lim_{R \rightarrow \infty} \int_1^R \left(\frac{1}{x} - \frac{1}{x+1} \right) dx = \lim_{R \rightarrow \infty} \ln \frac{x}{x+1} \Big|_1^R = \lim_{R \rightarrow \infty} \left(\ln \frac{R}{R+1} - \ln \frac{1}{2} \right) = \ln 1 - \ln \frac{1}{2} = \ln 2.$$

The integral converges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$ converges.

$$10. \sum_{n=1}^{\infty} ne^{-n^2}$$

SOLUTION Let $f(x) = xe^{-x^2}$. This function is continuous and positive on the interval $x \geq 1$. Moreover, because

$$f'(x) = 1 \cdot e^{-x^2} + x \cdot e^{-x^2} \cdot (-2x) = e^{-x^2} (1 - 2x^2),$$

we see that $f'(x) < 0$ for $x \geq 1$, so f is decreasing on this interval. To compute the improper integral we make the substitution $u = x^2$, $du = 2x dx$. Then, we find

$$\int_1^{\infty} xe^{-x^2} dx = \lim_{R \rightarrow \infty} \int_1^R xe^{-x^2} dx = \frac{1}{2} \int_1^{R^2} e^{-u} du = -\frac{1}{2} \lim_{R \rightarrow \infty} (e^{-R^2} - e^{-1}) = \frac{1}{2e}.$$

The integral converges; hence, the series $\sum_{n=1}^{\infty} ne^{-n^2}$ also converges.

$$11. \sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$$

SOLUTION Let $f(x) = \frac{1}{x(\ln x)^2}$. This function is positive and continuous for $x \geq 2$. Moreover,

$$f'(x) = -\frac{1}{x^2(\ln x)^4} \left(1 \cdot (\ln x)^2 + x \cdot 2(\ln x) \cdot \frac{1}{x} \right) = -\frac{1}{x^2(\ln x)^4} \left((\ln x)^2 + 2 \ln x \right).$$

Since $\ln x > 0$ for $x > 1$, $f'(x)$ is negative for $x > 1$; hence, f is decreasing for $x \geq 2$. To compute the improper integral, we make the substitution $u = \ln x$, $du = \frac{1}{x} dx$. We obtain:

$$\begin{aligned} \int_2^{\infty} \frac{1}{x(\ln x)^2} dx &= \lim_{R \rightarrow \infty} \int_2^R \frac{1}{x(\ln x)^2} dx = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u^2} \\ &= -\lim_{R \rightarrow \infty} \left(\frac{1}{\ln R} - \frac{1}{\ln 2} \right) = \frac{1}{\ln 2}. \end{aligned}$$

The integral converges; hence, the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}$ also converges.

$$12. \sum_{n=1}^{\infty} \frac{\ln n}{n^2}$$

SOLUTION Let $f(x) = \frac{\ln x}{x^2}$. Because

$$f'(x) = \frac{\frac{1}{x} \cdot x^2 - 2x \ln x}{x^4} = \frac{x(1 - 2 \ln x)}{x^4} = \frac{1 - 2 \ln x}{x^3},$$

we see that $f'(x) < 0$ for $x > \sqrt{e} \approx 1.65$. We conclude that f is decreasing on the interval $x \geq 2$. Since f is also positive and continuous on this interval, the Integral Test can be applied. By Integration by Parts, we find

$$\int \frac{\ln x}{x^2} dx = -\frac{\ln x}{x} + \int x^{-2} dx = -\frac{\ln x}{x} - \frac{1}{x} + C;$$

therefore,

$$\int_2^{\infty} \frac{\ln x}{x^2} dx = \lim_{R \rightarrow \infty} \int_2^R \frac{\ln x}{x^2} dx = \lim_{R \rightarrow \infty} \left(\frac{1}{2} + \frac{\ln 2}{2} - \frac{1}{R} - \frac{\ln R}{R} \right) = \frac{1 + \ln 2}{2} - \lim_{R \rightarrow \infty} \frac{\ln R}{R}.$$

We compute the resulting limit using L'Hôpital's Rule:

$$\lim_{R \rightarrow \infty} \frac{\ln R}{R} = \lim_{R \rightarrow \infty} \frac{1/R}{1} = 0.$$

Hence,

$$\int_2^{\infty} \frac{\ln x}{x^2} dx = \frac{1 + \ln 2}{2}.$$

The integral converges; therefore, the series $\sum_{n=2}^{\infty} \frac{\ln n}{n^2}$ also converges. Since the convergence of the series is not affected

by adding the finite sum $\sum_{n=1}^1 \frac{\ln n}{n^2}$, the series $\sum_{n=1}^{\infty} \frac{\ln n}{n^2}$ also converges.

$$13. \sum_{n=1}^{\infty} \frac{1}{2^{\ln n}}$$

SOLUTION Note that

$$2^{\ln n} = (e^{\ln 2})^{\ln n} = (e^{\ln n})^{\ln 2} = n^{\ln 2}.$$

Thus,

$$\sum_{n=1}^{\infty} \frac{1}{2^{\ln n}} = \sum_{n=1}^{\infty} \frac{1}{n^{\ln 2}}.$$

Now, let $f(x) = \frac{1}{x^{\ln 2}}$. This function is positive, continuous and decreasing on the interval $x \geq 1$; therefore, the Integral Test applies. Moreover,

$$\int_1^{\infty} \frac{dx}{x^{\ln 2}} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x^{\ln 2}} = \frac{1}{1 - \ln 2} \lim_{R \rightarrow \infty} (R^{1 - \ln 2} - 1) = \infty,$$

because $1 - \ln 2 > 0$. The integral diverges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{2^{\ln n}}$ also diverges.

14. $\sum_{n=1}^{\infty} \frac{1}{3^{\ln n}}$

SOLUTION Note that

$$3^{\ln n} = (e^{\ln 3})^{\ln n} = (e^{\ln n})^{\ln 3} = n^{\ln 3}.$$

Thus,

$$\sum_{n=1}^{\infty} \frac{1}{3^{\ln n}} = \sum_{n=1}^{\infty} \frac{1}{n^{\ln 3}}.$$

Now, let $f(x) = \frac{1}{x^{\ln 3}}$. This function is positive, continuous and decreasing on the interval $x \geq 1$; therefore, the Integral Test applies. Moreover,

$$\int_1^{\infty} \frac{dx}{x^{\ln 3}} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x^{\ln 3}} = \frac{1}{1 - \ln 3} \lim_{R \rightarrow \infty} (R^{1 - \ln 3} - 1) = -\frac{1}{1 - \ln 3},$$

because $1 - \ln 3 < 0$. The integral converges; hence, the series $\sum_{n=1}^{\infty} \frac{1}{3^{\ln n}}$ also converges.

15. Show that $\sum_{n=1}^{\infty} \frac{1}{n^3 + 8n}$ converges by using the Comparison Test with $\sum_{n=1}^{\infty} n^{-3}$.

SOLUTION We compare the series with the p -series $\sum_{n=1}^{\infty} n^{-3}$. For $n \geq 1$,

$$\frac{1}{n^3 + 8n} \leq \frac{1}{n^3}.$$

Since $\sum_{n=1}^{\infty} \frac{1}{n^3}$ converges (it is a p -series with $p = 3 > 1$), the series $\sum_{n=1}^{\infty} \frac{1}{n^3 + 8n}$ also converges by the Comparison Test.

16. Show that $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n^2 - 3}}$ diverges by comparing with $\sum_{n=2}^{\infty} n^{-1}$.

SOLUTION For $n \geq 2$,

$$\frac{1}{\sqrt{n^2 - 3}} \geq \frac{1}{\sqrt{n^2}} = \frac{1}{n}.$$

The harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges, and it still diverges if we drop the first term. Thus, the series $\sum_{n=2}^{\infty} \frac{1}{n}$ also diverges.

The Comparison Test now lets us conclude that the larger series $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n^2 - 3}}$ also diverges.

17. Let $S = \sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}}$. Verify that for $n \geq 1$,

$$\frac{1}{n + \sqrt{n}} \leq \frac{1}{n}, \quad \frac{1}{n + \sqrt{n}} \leq \frac{1}{\sqrt{n}}$$

Can either inequality be used to show that S diverges? Show that $\frac{1}{n + \sqrt{n}} \geq \frac{1}{2n}$ and conclude that S diverges.

SOLUTION For $n \geq 1$, $n + \sqrt{n} \geq n$ and $n + \sqrt{n} \geq \sqrt{n}$. Taking the reciprocal of each of these inequalities yields

$$\frac{1}{n + \sqrt{n}} \leq \frac{1}{n} \quad \text{and} \quad \frac{1}{n + \sqrt{n}} \leq \frac{1}{\sqrt{n}}.$$

These inequalities indicate that the series $\sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}}$ is smaller than both $\sum_{n=1}^{\infty} \frac{1}{n}$ and $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$; however, $\sum_{n=1}^{\infty} \frac{1}{n}$ and $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ both diverge so neither inequality allows us to show that S diverges.

On the other hand, for $n \geq 1$, $n \geq \sqrt{n}$, so $2n \geq n + \sqrt{n}$ and

$$\frac{1}{n + \sqrt{n}} \geq \frac{1}{2n}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{2n} = 2 \sum_{n=1}^{\infty} \frac{1}{n}$ diverges, since the harmonic series diverges. The Comparison Test then lets us conclude that the larger series $\sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}}$ also diverges.

18. Which of the following inequalities can be used to study the convergence of $\sum_{n=2}^{\infty} \frac{1}{n^2 + \sqrt{n}}$? Explain.

$$\frac{1}{n^2 + \sqrt{n}} \leq \frac{1}{\sqrt{n}}, \quad \frac{1}{n^2 + \sqrt{n}} \leq \frac{1}{n^2}$$

SOLUTION The series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ is a divergent p -series, hence the series $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n}}$ also diverges. The first inequality given above therefore establishes that $\sum_{n=2}^{\infty} \frac{1}{n^2 + \sqrt{n}}$ is smaller than a divergent series, which does not allow us to conclude whether $\sum_{n=2}^{\infty} \frac{1}{n^2 + \sqrt{n}}$ converges or diverges.

However, the second inequality given above establishes that $\sum_{n=2}^{\infty} \frac{1}{n^2 + \sqrt{n}}$ is smaller than the convergent p -series $\sum_{n=2}^{\infty} \frac{1}{n^2}$. By the Comparison Test, we therefore conclude that $\sum_{n=2}^{\infty} \frac{1}{n^2 + \sqrt{n}}$ also converges.

In Exercises 19–30, use the Comparison Test to determine whether the infinite series is convergent.

19. $\sum_{n=1}^{\infty} \frac{1}{n2^n}$

SOLUTION We compare with the geometric series $\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n$. For $n \geq 1$,

$$\frac{1}{n2^n} \leq \frac{1}{2^n} = \left(\frac{1}{2}\right)^n.$$

Since $\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n$ converges (it is a geometric series with $r = \frac{1}{2}$), we conclude by the Comparison Test that $\sum_{n=1}^{\infty} \frac{1}{n2^n}$ also converges.

20. $\sum_{n=1}^{\infty} \frac{n^3}{n^5 + 4n + 1}$

SOLUTION For $n \geq 1$,

$$\frac{n^3}{n^5 + 4n + 1} \leq \frac{n^3}{n^5} = \frac{1}{n^2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=1}^{\infty} \frac{n^3}{n^5 + 4n + 1}$ also converges.

$$21. \sum_{n=1}^{\infty} \frac{1}{n^{1/3} + 2^n}$$

SOLUTION For $n \geq 1$,

$$\frac{1}{n^{1/3} + 2^n} \leq \frac{1}{2^n}$$

The series $\sum_{n=1}^{\infty} \frac{1}{2^n}$ is a geometric series with $r = \frac{1}{2}$, so it converges. By the Comparison test, so does $\sum_{n=1}^{\infty} \frac{1}{n^{1/3} + 2^n}$.

$$22. \sum_{n=1}^{\infty} \frac{1}{\sqrt{n^3 + 2n - 1}}$$

SOLUTION For $n \geq 1$, we have $2n - 1 \geq 0$ so that

$$\frac{1}{\sqrt{n^3 + 2n - 1}} \leq \frac{1}{\sqrt{n^3}} = \frac{1}{n^{3/2}}.$$

This latter series is a p -series with $p = \frac{3}{2} > 1$, so it converges. By the Comparison Test, so does $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n^3 + 2n - 1}}$.

$$23. \sum_{m=1}^{\infty} \frac{4}{m! + 4^m}$$

SOLUTION For $m \geq 1$,

$$\frac{4}{m! + 4^m} \leq \frac{4}{4^m} = \left(\frac{1}{4}\right)^{m-1}.$$

The series $\sum_{m=1}^{\infty} \left(\frac{1}{4}\right)^{m-1}$ is a geometric series with $r = \frac{1}{4}$, so it converges. By the Comparison Test we can therefore conclude that the series $\sum_{m=1}^{\infty} \frac{4}{m! + 4^m}$ also converges.

$$24. \sum_{n=4}^{\infty} \frac{\sqrt{n}}{n-3}$$

SOLUTION For $n \geq 4$,

$$\frac{\sqrt{n}}{n-3} \geq \frac{\sqrt{n}}{n} = \frac{1}{n^{1/2}}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^{1/2}}$ is a p -series with $p = \frac{1}{2} < 1$, so it diverges, and it continues to diverge if we drop the terms $n = 1, 2, 3$; that is, $\sum_{n=4}^{\infty} \frac{1}{n^{1/2}}$ also diverges. By the Comparison Test we can therefore conclude that series $\sum_{n=4}^{\infty} \frac{\sqrt{n}}{n-3}$ diverges.

$$25. \sum_{k=1}^{\infty} \frac{\sin^2 k}{k^2}$$

SOLUTION For $k \geq 1$, $0 \leq \sin^2 k \leq 1$, so

$$0 \leq \frac{\sin^2 k}{k^2} \leq \frac{1}{k^2}.$$

The series $\sum_{k=1}^{\infty} \frac{1}{k^2}$ is a p -series with $p = 2 > 1$, so it converges. By the Comparison Test we can therefore conclude that the series $\sum_{k=1}^{\infty} \frac{\sin^2 k}{k^2}$ also converges.

$$26. \sum_{k=2}^{\infty} \frac{k^{1/3}}{k^{5/4} - k}$$

SOLUTION For $k \geq 2$, $k^{5/4} - k < k^{5/4}$ so that

$$\frac{k^{1/3}}{k^{5/4} - k} \geq \frac{k^{1/3}}{k^{5/4}} = \frac{1}{k^{11/12}}$$

The series $\sum_{k=2}^{\infty} \frac{1}{k^{11/12}}$ is a p -series with $p = \frac{11}{12} < 1$, so it diverges. By the Comparison Test, so does $\sum_{k=2}^{\infty} \frac{k^{1/3}}{k^{5/4} - k}$.

$$27. \sum_{n=1}^{\infty} \frac{2}{3^n + 3^{-n}}$$

SOLUTION Since $3^{-n} > 0$ for all n ,

$$\frac{2}{3^n + 3^{-n}} \leq \frac{2}{3^n} = 2 \left(\frac{1}{3}\right)^n.$$

The series $\sum_{n=1}^{\infty} 2 \left(\frac{1}{3}\right)^n$ is a geometric series with $r = \frac{1}{3}$, so it converges. By the Comparison Theorem we can therefore

conclude that the series $\sum_{n=1}^{\infty} \frac{2}{3^n + 3^{-n}}$ also converges.

$$28. \sum_{k=1}^{\infty} 2^{-k^2}$$

SOLUTION For $k \geq 1$, $k^2 \geq k$ and

$$\frac{1}{2^{k^2}} \leq \frac{1}{2^k} = \left(\frac{1}{2}\right)^k.$$

The series $\sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^k$ is a geometric series with $r = \frac{1}{2}$, so it converges. By the Comparison Test we can therefore conclude

that the series $\sum_{k=1}^{\infty} \frac{1}{2^{k^2}} = \sum_{k=1}^{\infty} 2^{-k^2}$ also converges.

$$29. \sum_{n=1}^{\infty} \frac{1}{(n+1)!}$$

SOLUTION Note that for $n \geq 2$,

$$(n+1)! = 1 \cdot \underbrace{2 \cdot 3 \cdots n}_{n \text{ factors}} \cdot (n+1) \leq 2^n$$

so that

$$\sum_{n=1}^{\infty} \frac{1}{(n+1)!} = 1 + \sum_{n=2}^{\infty} \frac{1}{(n+1)!} \leq 1 + \sum_{n=2}^{\infty} \frac{1}{2^n}$$

But $\sum_{n=2}^{\infty} \frac{1}{2^n}$ is a geometric series with ratio $r = \frac{1}{2}$, so it converges. By the comparison test, $\sum_{n=1}^{\infty} \frac{1}{(n+1)!}$ converges as well.

$$30. \sum_{n=1}^{\infty} \frac{n!}{n^3}$$

SOLUTION Note that for $n \geq 4$, we have $(n-1)(n-2) > n$ [to see this, solve the equation $(n-1)(n-2) = n$ for n ; the positive root is $2 + \sqrt{2} \approx 3.4$]. Thus

$$\sum_{n=4}^{\infty} \frac{n!}{n^3} = \sum_{n=4}^{\infty} \frac{n(n-1)(n-2)(n-3)!}{n^3} \geq \sum_{n=4}^{\infty} \frac{(n-3)!}{n} \geq \sum_{n=4}^{\infty} \frac{1}{n}$$

But $\sum_{n=4}^{\infty} \frac{1}{n}$ is the harmonic series, which diverges, so that $\sum_{n=4}^{\infty} \frac{n!}{n^3}$ also diverges. Adding back in the terms for $n = 1, 2$, and 3 does not affect this result. Thus the original series diverges.

Exercise 31–36: For all $a > 0$ and $b > 1$, the inequalities

$$\ln n \leq n^a, \quad n^a < b^n$$

are true for n sufficiently large (this can be proved using L'Hopital's Rule). Use this, together with the Comparison Theorem, to determine whether the series converges or diverges.

$$31. \sum_{n=1}^{\infty} \frac{\ln n}{n^3}$$

SOLUTION For n sufficiently large (say $n = k$, although in this case $n = 1$ suffices), we have $\ln n \leq n$, so that

$$\sum_{n=k}^{\infty} \frac{\ln n}{n^3} \leq \sum_{n=k}^{\infty} \frac{n}{n^3} = \sum_{n=k}^{\infty} \frac{1}{n^2}$$

This is a p -series with $p = 2 > 1$, so it converges. Thus $\sum_{n=k}^{\infty} \frac{\ln n}{n^3}$ also converges; adding back in the finite number of terms for $1 \leq n \leq k$ does not affect this result.

$$32. \sum_{m=2}^{\infty} \frac{1}{\ln m}$$

SOLUTION For $m > 1$ sufficiently large (say $m = k$, although in this case $m = 2$ suffices), we have $\ln m \leq m$, so that

$$\sum_{m=k}^{\infty} \frac{1}{\ln m} \geq \sum_{m=k}^{\infty} \frac{1}{m}$$

This is the harmonic series, which diverges (the absence of the finite number of terms for $m = 1, \dots, k - 1$ does not affect convergence). By the comparison theorem, $\sum_{m=2}^{\infty} \frac{1}{\ln m}$ also diverges (again, ignoring the finite number of terms for $m = 1, \dots, k - 1$ does not affect convergence).

$$33. \sum_{n=1}^{\infty} \frac{(\ln n)^{100}}{n^{1.1}}$$

SOLUTION Choose N so that $\ln n \leq n^{0.0005}$ for $n \geq N$. Then also for $n > N$, $(\ln n)^{100} \leq (n^{0.0005})^{100} = n^{0.05}$. Then

$$\sum_{n=N}^{\infty} \frac{(\ln n)^{100}}{n^{1.1}} \leq \sum_{n=N}^{\infty} \frac{n^{0.05}}{n^{1.1}} = \sum_{n=N}^{\infty} \frac{1}{n^{1.05}}$$

But $\sum_{n=N}^{\infty} \frac{1}{n^{1.05}}$ is a p -series with $p = 1.05 > 1$, so is convergent. It follows that $\sum_{n=N}^{\infty} \frac{(\ln n)^{100}}{n^{1.1}}$ is also convergent;

adding back in the finite number of terms for $n = 1, 2, \dots, N - 1$ shows that $\sum_{n=1}^{\infty} \frac{(\ln n)^{100}}{n^{1.1}}$ converges as well.

$$34. \sum_{n=1}^{\infty} \frac{1}{(\ln n)^{10}}$$

SOLUTION Choose N such that $\ln n \leq n^{0.1}$ for $n \geq N$; then also $(\ln n)^{10} \leq n$ for $n \geq N$. So we have

$$\sum_{n=N}^{\infty} \frac{1}{(\ln n)^{10}} \geq \sum_{n=N}^{\infty} \frac{1}{n}$$

The latter sum is the harmonic series, which diverges, so the series on the left diverges as well. Adding back in the finite number of terms for $n < N$ shows that $\sum_{n=1}^{\infty} \frac{1}{(\ln n)^{10}}$ diverges.

$$35. \sum_{n=1}^{\infty} \frac{n}{3^n}$$

SOLUTION Choose N such that $n \leq 2^n$ for $n \geq N$. Then

$$\sum_{n=N}^{\infty} \frac{n}{3^n} \leq \sum_{n=N}^{\infty} \left(\frac{2}{3}\right)^n$$

The latter sum is a geometric series with $r = \frac{2}{3} < 1$, so it converges. Thus the series on the left converges as well. Adding back in the finite number of terms for $n < N$ shows that $\sum_{n=1}^{\infty} \frac{n}{3^n}$ converges.

$$36. \sum_{n=1}^{\infty} \frac{n^5}{2^n}$$

SOLUTION Choose N such that $n^5 \leq 1.5^n$ for $n \geq N$. Then

$$\sum_{n=N}^{\infty} \frac{n^5}{2^n} \leq \sum_{n=N}^{\infty} \left(\frac{1.5}{2}\right)^n$$

The latter sum is a geometric series with $r = \frac{1.5}{2} < 1$, so it converges. Thus the series on the left converges as well. Adding back in the finite number of terms for $n < N$ shows that $\sum_{n=1}^{\infty} \frac{n^5}{2^n}$ converges.

37. Show that $\sum_{n=1}^{\infty} \sin \frac{1}{n^2}$ converges. *Hint:* Use the inequality $\sin x \leq x$ for $x \geq 0$.

SOLUTION For $n \geq 1$,

$$0 \leq \frac{1}{n^2} \leq 1 < \pi;$$

therefore, $\sin \frac{1}{n^2} > 0$ for $n \geq 1$. Moreover, for $n \geq 1$,

$$\sin \frac{1}{n^2} \leq \frac{1}{n^2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=1}^{\infty} \sin \frac{1}{n^2}$ also converges.

38. Does $\sum_{n=2}^{\infty} \frac{\sin(1/n)}{\ln n}$ converge?

SOLUTION No, it diverges. Either the Comparison Theorem or the Limit Comparison Theorem may be used. Using the Comparison Theorem, recall that

$$\frac{\sin x}{x} > \cos x \quad \text{for } x > 0$$

so that $\sin x > x \cos x$. Substituting $1/n$ for x gives

$$\sin\left(\frac{1}{n}\right) > \frac{1}{n} \cos\left(\frac{1}{n}\right) = \frac{\cos(1/n)}{n} \geq \frac{1}{2n}$$

since $\cos\left(\frac{1}{n}\right) \geq \frac{1}{2}$ for $n \geq 2$. Thus

$$\sum_{n=1}^{\infty} \frac{\sin(1/n)}{\ln n} > \sum_{n=1}^{\infty} \frac{1}{2n \ln n}$$

Apply the Integral Test to the latter expression, making the substitution $u = \ln x$:

$$\int_1^{\infty} \frac{1}{2x \ln x} dx = \frac{1}{2} \int_0^{\infty} \frac{1}{u} du = \frac{1}{2} \ln u \Big|_0^{\infty}$$

and the integral diverges. Thus

$$\sum_{n=1}^{\infty} \frac{1}{2n \ln n} \text{ diverges, and thus } \sum_{n=1}^{\infty} \frac{\sin(1/n)}{\ln n} \text{ diverges as well.}$$

Applying the Limit Comparison Test is similar but perhaps simpler: Recall that

$$\lim_{x \rightarrow \infty} \frac{\sin(1/x)}{1/x} = \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

so apply the Limit Comparison Test with $b_n = \frac{1/x}{\ln x}$:

$$L = \lim_{x \rightarrow \infty} \frac{\sin(1/x)}{\ln x} \cdot \frac{\ln x}{1/x} = \lim_{x \rightarrow \infty} \frac{\sin(1/x)}{1/x} = 1$$

so that either both series converge or both diverge. But by the Integral Test as above,

$$\sum_{n=1}^{\infty} \frac{(1/x)}{\ln x} = \sum_{n=1}^{\infty} \frac{1}{x \ln x}$$

diverges.

In Exercises 39–48, use the Limit Comparison Test to prove convergence or divergence of the infinite series.

$$39. \sum_{n=2}^{\infty} \frac{n^2}{n^4 - 1}$$

SOLUTION Let $a_n = \frac{n^2}{n^4 - 1}$. For large n , $\frac{n^2}{n^4 - 1} \approx \frac{n^2}{n^4} = \frac{1}{n^2}$, so we apply the Limit Comparison Test with $b_n = \frac{1}{n^2}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{n^2}{n^4 - 1}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{n^4}{n^4 - 1} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges; hence, $\sum_{n=2}^{\infty} \frac{1}{n^2}$ also converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=2}^{\infty} \frac{n^2}{n^4 - 1}$ converges.

$$40. \sum_{n=2}^{\infty} \frac{1}{n^2 - \sqrt{n}}$$

SOLUTION Let $a_n = \frac{1}{n^2 - \sqrt{n}}$. For large n , $\frac{1}{n^2 - \sqrt{n}} \approx \frac{1}{n^2}$, so we apply the Limit Comparison Test with $b_n = \frac{1}{n^2}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{n^2 - \sqrt{n}}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{n^2}{n^2 - \sqrt{n}} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges; hence, the series $\sum_{n=2}^{\infty} \frac{1}{n^2}$ also converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=2}^{\infty} \frac{1}{n^2 - \sqrt{n}}$ converges.

$$41. \sum_{n=2}^{\infty} \frac{n}{\sqrt{n^3 + 1}}$$

SOLUTION Let $a_n = \frac{n}{\sqrt{n^3 + 1}}$. For large n , $\frac{n}{\sqrt{n^3 + 1}} \approx \frac{n}{\sqrt{n^3}} = \frac{1}{\sqrt{n}}$, so we apply the Limit Comparison test with $b_n = \frac{1}{\sqrt{n}}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{n}{\sqrt{n^3 + 1}}}{\frac{1}{\sqrt{n}}} = \lim_{n \rightarrow \infty} \frac{\sqrt{n^3}}{\sqrt{n^3 + 1}} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ is a p -series with $p = \frac{1}{2} < 1$, so it diverges; hence, $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n}}$ also diverges. Because $L > 0$, by the Limit Comparison Test we can conclude that the series $\sum_{n=2}^{\infty} \frac{n}{\sqrt{n^3 + 1}}$ diverges.

$$42. \sum_{n=2}^{\infty} \frac{n^3}{\sqrt{n^7 + 2n^2 + 1}}$$

SOLUTION Let a_n be the general term of our series. Observe that

$$a_n = \frac{n^3}{\sqrt{n^7 + 2n^2 + 1}} = \frac{n^{-3} \cdot n^3}{n^{-3} \cdot \sqrt{n^7 + 2n^2 + 1}} = \frac{1}{\sqrt{n + 2n^{-4} + n^{-6}}}$$

This suggests that we apply the Limit Comparison Test, comparing our series with

$$\sum_{n=2}^{\infty} b_n = \sum_{n=2}^{\infty} \frac{1}{n^{1/2}}$$

The ratio of the terms is

$$\frac{a_n}{b_n} = \frac{1}{\sqrt{n + 2n^{-4} + n^{-6}}} \cdot \frac{\sqrt{n}}{1} = \frac{1}{\sqrt{1 + 2n^{-5} + n^{-7}}}$$

Hence

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{1 + 2n^{-5} + n^{-7}}} = 1$$

The p -series $\sum_{n=2}^{\infty} \frac{1}{n^{1/2}}$ diverges since $p = 1/2 < 1$. Therefore, our original series diverges.

$$43. \sum_{n=3}^{\infty} \frac{3n + 5}{n(n-1)(n-2)}$$

SOLUTION Let $a_n = \frac{3n + 5}{n(n-1)(n-2)}$. For large n , $\frac{3n + 5}{n(n-1)(n-2)} \approx \frac{3n}{n^3} = \frac{3}{n^2}$, so we apply the Limit Comparison Test with $b_n = \frac{1}{n^2}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{3n+5}{n(n+1)(n+2)}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{3n^3 + 5n^2}{n(n+1)(n+2)} = 3.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges; hence, the series $\sum_{n=3}^{\infty} \frac{1}{n^2}$ also converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=3}^{\infty} \frac{3n + 5}{n(n-1)(n-2)}$ converges.

$$44. \sum_{n=1}^{\infty} \frac{e^n + n}{e^{2n} - n^2}$$

SOLUTION Let

$$a_n = \frac{e^n + n}{e^{2n} - n^2} = \frac{e^n + n}{(e^n - n)(e^n + n)} = \frac{1}{e^n - n}.$$

For large n ,

$$\frac{1}{e^n - n} \approx \frac{1}{e^n} = e^{-n},$$

so we apply the Limit Comparison Test with $b_n = e^{-n}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{e^n - n}}{e^{-n}} = \lim_{n \rightarrow \infty} \frac{e^n}{e^n - n} = 1.$$

The series $\sum_{n=1}^{\infty} e^{-n} = \sum_{n=1}^{\infty} \left(\frac{1}{e}\right)^n$ is a geometric series with $r = \frac{1}{e} < 1$, so it converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \frac{e^n + n}{e^{2n} - n^2}$ also converges.

$$45. \sum_{n=1}^{\infty} \frac{1}{\sqrt{n} + \ln n}$$

SOLUTION Let

$$a_n = \frac{1}{\sqrt{n} + \ln n}$$

For large n , $\sqrt{n} + \ln n \approx \sqrt{n}$, so apply the Comparison Test with $b_n = \frac{1}{\sqrt{n}}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n} + \ln n} \cdot \frac{\sqrt{n}}{1} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{\ln n}{\sqrt{n}}} = 1$$

The series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ is a p -series with $p = \frac{1}{2} < 1$, so it diverges. Because L exists, the Limit Comparison Test tells us the original series also diverges.

$$46. \sum_{n=1}^{\infty} \frac{\ln(n+4)}{n^{5/2}}$$

SOLUTION Let

$$a_n = \frac{\ln(n+4)}{n^{5/2}}$$

For large n , $a_n \approx \frac{\ln n}{n^{5/2}}$, so apply the Comparison Test with $b_n = \frac{\ln n}{n^{5/2}}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\ln(n+4)}{n^{5/2}} \cdot \frac{n^{5/2}}{\ln n} = \lim_{n \rightarrow \infty} \frac{\ln(n+4)}{\ln n}$$

Applying L'Hôpital's rule gives

$$L = \lim_{n \rightarrow \infty} \frac{\ln(n+4)}{\ln n} = \lim_{n \rightarrow \infty} \frac{1/(n+4)}{1/n} = \lim_{n \rightarrow \infty} \frac{n}{n+4} = \lim_{n \rightarrow \infty} \frac{1}{1+4/n} = 1$$

To see that $\sum_{n=1}^{\infty} b_n$ converges, choose N so that $\ln n < n$ for $n \geq N$; then

$$\sum_{n=N}^{\infty} \frac{\ln n}{n^{5/2}} \leq \sum_{n=N}^{\infty} \frac{n}{n^{5/2}} = \sum_{n=N}^{\infty} \frac{1}{n^{3/2}}$$

which is a p -series with $p = \frac{3}{2} > 1$, so it converges. Adding back in the finite number of terms for $n < N$ shows that $\sum b_n$ converges as well. Since L exists and $\sum b_n$ converges, the Limit Comparison Test tells us that $\sum_{n=1}^{\infty} a_n$ converges.

$$47. \sum_{n=1}^{\infty} \left(1 - \cos \frac{1}{n}\right) \text{ Hint: Compare with } \sum_{n=1}^{\infty} n^{-2}.$$

SOLUTION Let $a_n = 1 - \cos \frac{1}{n}$, and apply the Limit Comparison Test with $b_n = \frac{1}{n^2}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1 - \cos \frac{1}{n}}{\frac{1}{n^2}} = \lim_{x \rightarrow \infty} \frac{1 - \cos \frac{1}{x}}{\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{-\frac{1}{x^2} \sin \frac{1}{x}}{-\frac{2}{x^3}} = \frac{1}{2} \lim_{x \rightarrow \infty} \frac{\sin \frac{1}{x}}{\frac{1}{x}}$$

As $x \rightarrow \infty$, $u = \frac{1}{x} \rightarrow 0$, so

$$L = \frac{1}{2} \lim_{x \rightarrow \infty} \frac{\sin \frac{1}{x}}{\frac{1}{x}} = \frac{1}{2} \lim_{u \rightarrow 0} \frac{\sin u}{u} = \frac{1}{2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a p -series with $p = 2 > 1$, so it converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \left(1 - \cos \frac{1}{n}\right)$ also converges.

48. $\sum_{n=1}^{\infty} (1 - 2^{-1/n})$ *Hint:* Compare with the harmonic series.

SOLUTION Let $a_n = 1 - 2^{-1/n}$, and apply the Limit Comparison Test with $b_n = \frac{1}{n}$. We find

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1 - 2^{-1/n}}{\frac{1}{n}} = \lim_{x \rightarrow \infty} \frac{1 - 2^{-1/x}}{\frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{-\frac{1}{x^2}(\ln 2)2^{-1/x}}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} (2^{-1/x} \ln 2) = \ln 2.$$

The harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges; because $L > 0$, we can conclude by the Limit Comparison Test that the series $\sum_{n=1}^{\infty} (1 - 2^{-1/n})$ also diverges.

In Exercises 49–74, determine convergence or divergence using any method covered so far.

49. $\sum_{n=4}^{\infty} \frac{1}{n^2 - 9}$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{1}{n^2 - 9}$ and $b_n = \frac{1}{n^2}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{n^2 - 9}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{n^2}{n^2 - 9} = 1.$$

Since the p -series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, the series $\sum_{n=4}^{\infty} \frac{1}{n^2}$ also converges. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=4}^{\infty} \frac{1}{n^2 - 9}$ converges.

50. $\sum_{n=1}^{\infty} \frac{\cos^2 n}{n^2}$

SOLUTION For all $n \geq 1$, $0 \leq \cos^2 n \leq 1$, so

$$0 \leq \frac{\cos^2 n}{n^2} \leq \frac{1}{n^2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series; hence, by the Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \frac{\cos^2 n}{n^2}$ also converges.

51. $\sum_{n=1}^{\infty} \frac{\sqrt{n}}{4n + 9}$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{\sqrt{n}}{4n + 9}$ and $b_n = \frac{1}{\sqrt{n}}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{\sqrt{n}}{4n + 9}}{\frac{1}{\sqrt{n}}} = \lim_{n \rightarrow \infty} \frac{n}{4n + 9} = \frac{1}{4}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ is a divergent p -series. Because $L > 0$, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \frac{\sqrt{n}}{4n + 9}$ also diverges.

$$52. \sum_{n=1}^{\infty} \frac{n - \cos n}{n^3}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{n - \cos n}{n^3}$ and $b_n = \frac{1}{n^2}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{n - \cos n}{n^3}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \left(1 - \frac{\cos n}{n}\right) = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series. Because L exists, by the Limit Comparison Test we can conclude that the

series $\sum_{n=1}^{\infty} \frac{n - \cos n}{n^3}$ also converges.

$$53. \sum_{n=1}^{\infty} \frac{n^2 - n}{n^5 + n}$$

SOLUTION First rewrite $a_n = \frac{n^2 - n}{n^5 + n} = \frac{n(n-1)}{n(n^4+1)} = \frac{n-1}{n^4+1}$ and observe

$$\frac{n-1}{n^4+1} < \frac{n}{n^4} = \frac{1}{n^3}$$

for $n \geq 1$. The series $\sum_{n=1}^{\infty} \frac{1}{n^3}$ is a convergent p -series, so by the Comparison Test we can conclude that the series

$\sum_{n=1}^{\infty} \frac{n^2 - n}{n^5 + n}$ also converges.

$$54. \sum_{n=1}^{\infty} \frac{1}{n^2 + \sin n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{1}{n^2 + \sin n}$ and $b_n = \frac{1}{n^2}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{n^2 + \sin n}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{\sin n}{n^2}} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series. Because L exists, by the Limit Comparison Test we can conclude that the

series $\sum_{n=1}^{\infty} \frac{1}{n^2 + \sin n}$ also converges.

$$55. \sum_{n=5}^{\infty} (4/5)^{-n}$$

SOLUTION

$$\sum_{n=5}^{\infty} \left(\frac{4}{5}\right)^{-n} = \sum_{n=5}^{\infty} \left(\frac{5}{4}\right)^n$$

which is a geometric series starting at $n = 5$ with ratio $r = \frac{5}{4} > 1$. Thus the series diverges.

$$56. \sum_{n=1}^{\infty} \frac{1}{3^{n^2}}$$

SOLUTION Because $n^2 \geq n$ for $n \geq 1$, $3^{n^2} \geq 3^n$ and

$$\frac{1}{3^{n^2}} \leq \frac{1}{3^n} = \left(\frac{1}{3}\right)^n.$$

The series $\sum_{n=1}^{\infty} \left(\frac{1}{3}\right)^n$ is a geometric series with $r = \frac{1}{3}$, so it converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=1}^{\infty} \frac{1}{3n^2}$ also converges.

$$57. \sum_{n=2}^{\infty} \frac{1}{n^{3/2} \ln n}$$

SOLUTION For $n \geq 3$, $\ln n > 1$, so $n^{3/2} \ln n > n^{3/2}$ and

$$\frac{1}{n^{3/2} \ln n} < \frac{1}{n^{3/2}}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ is a convergent p -series, so the series $\sum_{n=3}^{\infty} \frac{1}{n^{3/2}}$ also converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=3}^{\infty} \frac{1}{n^{3/2} \ln n}$ converges. Hence, the series $\sum_{n=2}^{\infty} \frac{1}{n^{3/2} \ln n}$ also converges.

$$58. \sum_{n=2}^{\infty} \frac{(\ln n)^{12}}{n^{9/8}}$$

SOLUTION By the comment preceding Exercise 31, we can choose N so that for $n \geq N$, we have $\ln n < n^{1/192}$. Then also for $n \geq N$ we have $(\ln n)^{12} < n^{12/192} = n^{1/16}$. Then

$$\sum_{n=N}^{\infty} \frac{(\ln n)^{12}}{n^{9/8}} \leq \sum_{n=N}^{\infty} \frac{n^{1/16}}{n^{9/8}} = \sum_{n=N}^{\infty} \frac{1}{n^{17/16}}$$

which is a convergent p -series. Thus the series on the left converges as well; adding back in the finite number of terms for $n \leq N$ shows that $\sum_{n=2}^{\infty} \frac{(\ln n)^{12}}{n^{9/8}}$ converges.

$$59. \sum_{k=1}^{\infty} 4^{1/k}$$

SOLUTION

$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} 4^{1/k} = 4^0 = 1 \neq 0;$$

therefore, the series $\sum_{k=1}^{\infty} 4^{1/k}$ diverges by the Divergence Test.

$$60. \sum_{n=1}^{\infty} \frac{4^n}{5^n - 2n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{4^n}{5^n - 2n}$ and $b_n = \frac{4^n}{5^n}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{4^n}{5^n - 2n}}{\frac{4^n}{5^n}} = \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{2n}{5^n}}.$$

Now,

$$\lim_{n \rightarrow \infty} \frac{2n}{5^n} = \lim_{x \rightarrow \infty} \frac{2x}{5^x} = \lim_{x \rightarrow \infty} \frac{2}{5^x \ln 5} = 0,$$

so

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{1}{1 - 0} = 1.$$

The series $\sum_{n=1}^{\infty} \left(\frac{4}{5}\right)^n$ is a convergent geometric series. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \frac{4^n}{5^n - 2n}$ also converges.

$$61. \sum_{n=2}^{\infty} \frac{1}{(\ln n)^4}$$

SOLUTION By the comment preceding Exercise 31, we can choose N so that for $n \geq N$, we have $\ln n < n^{1/8}$, so that $(\ln n)^4 < n^{1/2}$. Then

$$\sum_{n=N}^{\infty} \frac{1}{(\ln n)^4} > \sum_{n=N}^{\infty} \frac{1}{n^{1/2}}$$

which is a divergent p -series. Thus the series on the left diverges as well, and adding back in the finite number of terms for $n < N$ does not affect the result. Thus $\sum_{n=2}^{\infty} \frac{1}{(\ln n)^4}$ diverges.

$$62. \sum_{n=1}^{\infty} \frac{2^n}{3^n - n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{2^n}{3^n - n}$ and $b_n = \frac{2^n}{3^n}$:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{2^n}{3^n - n}}{\frac{2^n}{3^n}} = \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{n}{3^n}}$$

Now,

$$\lim_{n \rightarrow \infty} \frac{n}{3^n} = \lim_{x \rightarrow \infty} \frac{x}{3^x} = \lim_{x \rightarrow \infty} \frac{1}{3^x \ln 3} = 0,$$

so

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{1}{1 - 0} = 1.$$

The series $\sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n$ is a convergent geometric series. Because L exists, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \frac{2^n}{3^n - n}$ also converges.

$$63. \sum_{n=1}^{\infty} \frac{1}{n \ln n - n}$$

SOLUTION For $n \geq 2$, $n \ln n - n \leq n \ln n$; therefore,

$$\frac{1}{n \ln n - n} \geq \frac{1}{n \ln n}.$$

Now, let $f(x) = \frac{1}{x \ln x}$. For $x \geq 2$, this function is continuous, positive and decreasing, so the Integral Test applies. Using the substitution $u = \ln x$, $du = \frac{1}{x} dx$, we find

$$\int_2^{\infty} \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_2^R \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u} = \lim_{R \rightarrow \infty} (\ln(\ln R) - \ln(\ln 2)) = \infty.$$

The integral diverges; hence, the series $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ also diverges. By the Comparison Test we can therefore conclude that the series $\sum_{n=2}^{\infty} \frac{1}{n \ln n - n}$ diverges.

$$64. \sum_{n=1}^{\infty} \frac{1}{n(\ln n)^2 - n}$$

SOLUTION Use the Integral Test. Note that $x(\ln x)^2 - x$ has a zero at $x = e$, so restrict the integral to $[4, \infty)$:

$$\int_4^{\infty} \frac{1}{x(\ln x)^2 - x} dx$$

Substitute $u = \ln x$ so that $du = \frac{1}{x} dx$ to get

$$\begin{aligned} \int_{\ln 4}^{\infty} \frac{1}{u^2 - 1} du &= \lim_{R \rightarrow \infty} \left(\frac{1}{2} \ln \left| \frac{x-1}{x+1} \right| \Big|_4^R \right) = \frac{1}{2} \lim_{R \rightarrow \infty} \left(\ln \left(\frac{R-1}{R+1} \right) - \ln \left(\frac{3}{5} \right) \right) \\ &= \frac{1}{2} \left(\ln \lim_{R \rightarrow \infty} \left(\frac{R-1}{R+1} \right) - \ln \left(\frac{3}{5} \right) \right) = \frac{1}{2} \left(\ln 1 - \ln \left(\frac{3}{5} \right) \right) = \frac{1}{2} \ln \left(\frac{5}{3} \right) < \infty \end{aligned}$$

Since the integral converges, the series does as well starting at $n = 4$, using the Integral Test. Adding in the terms for $n = 1, 2, 3$ does not affect this result.

65. $\sum_{n=1}^{\infty} \frac{1}{n^n}$

SOLUTION For $n \geq 2$, $n^n \geq 2^n$; therefore,

$$\frac{1}{n^n} \leq \frac{1}{2^n} = \left(\frac{1}{2} \right)^n.$$

The series $\sum_{n=1}^{\infty} \left(\frac{1}{2} \right)^n$ is a convergent geometric series, so $\sum_{n=2}^{\infty} \left(\frac{1}{2} \right)^n$ also converges. By the Comparison Test we can

therefore conclude that the series $\sum_{n=2}^{\infty} \frac{1}{n^n}$ converges. Hence, the series $\sum_{n=1}^{\infty} \frac{1}{n^n}$ converges.

66. $\sum_{n=1}^{\infty} \frac{n^2 - 4n^{3/2}}{n^3}$

SOLUTION Let $a_n = \frac{1}{n}$ and $b_n = -\frac{4}{n^{3/2}}$. Then

$$\sum_{n=1}^{\infty} (a_n + b_n) = \sum_{n=1}^{\infty} \frac{n^2 - 4n^{3/2}}{n^3}$$

$\sum_{n=1}^{\infty} a_n$ diverges since it is the harmonic series

$\sum_{n=1}^{\infty} b_n$ is a p -series with $p = \frac{3}{2} > 1$, so converges

Since $\sum a_n$ diverges and $\sum b_n$ converges, it follows that $\sum (a_n + b_n)$ diverges.

67. $\sum_{n=1}^{\infty} \frac{1 + (-1)^n}{n}$

SOLUTION Let

$$a_n = \frac{1 + (-1)^n}{n}$$

Then

$$a_n = \begin{cases} 0 & n \text{ odd} \\ \frac{2}{2k} = \frac{1}{k} & n = 2k \text{ even} \end{cases}$$

Therefore, $\{a_n\}$ consists of 0s in the odd places and the harmonic series in the even places, so $\sum_{i=1}^{\infty} a_n$ is just the sum of the harmonic series, which diverges. Thus $\sum_{i=1}^{\infty} a_n$ diverges as well.

68. $\sum_{n=1}^{\infty} \frac{2 + (-1)^n}{n^{3/2}}$

SOLUTION For $n \geq 1$

$$0 < \frac{2 + (-1)^n}{n^{3/2}} \leq \frac{2 + 1}{n^{3/2}} = \frac{3}{n^{3/2}}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ is a convergent p -series; hence, the series $\sum_{n=1}^{\infty} \frac{3}{n^{3/2}}$ also converges. By the Comparison Test we can

therefore conclude that the series $\sum_{n=1}^{\infty} \frac{2 + (-1)^n}{n^{3/2}}$ converges.

$$69. \sum_{n=1}^{\infty} \sin \frac{1}{n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \sin \frac{1}{n}$ and $b_n = \frac{1}{n}$:

$$L = \lim_{n \rightarrow \infty} \frac{\sin \frac{1}{n}}{\frac{1}{n}} = \lim_{u \rightarrow 0} \frac{\sin u}{u} = 1,$$

where $u = \frac{1}{n}$. The harmonic series diverges. Because $L > 0$, by the Limit Comparison Test we can conclude that the series $\sum_{n=1}^{\infty} \sin \frac{1}{n}$ also diverges.

$$70. \sum_{n=1}^{\infty} \frac{\sin(1/n)}{\sqrt{n}}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{\sin(1/n)}{\sqrt{n}}$ and $b_n = \frac{1/n}{\sqrt{n}}$:

$$L = \lim_{n \rightarrow \infty} \frac{\sin(1/n)}{\sqrt{n}} \cdot \frac{\sqrt{n}}{1/n} = \lim_{n \rightarrow \infty} \frac{\sin(1/n)}{1/n} = \lim_{u \rightarrow 0} \frac{\sin u}{u} = 1$$

so that $\sum a_n$ and $\sum b_n$ either both converge or both diverge. But

$$\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} \frac{1/n}{\sqrt{n}} = \sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$$

is a convergent p -series. Thus $\sum_{n=1}^{\infty} \frac{\sin(1/n)}{\sqrt{n}}$ converges as well.

$$71. \sum_{n=1}^{\infty} \frac{2n+1}{4^n}$$

SOLUTION For $n \geq 3$, $2n+1 < 2^n$, so

$$\frac{2n+1}{4^n} < \frac{2^n}{4^n} = \left(\frac{1}{2}\right)^n.$$

The series $\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n$ is a convergent geometric series, so $\sum_{n=3}^{\infty} \left(\frac{1}{2}\right)^n$ also converges. By the Comparison Test we can

therefore conclude that the series $\sum_{n=3}^{\infty} \frac{2n+1}{4^n}$ converges. Finally, the series $\sum_{n=1}^{\infty} \frac{2n+1}{4^n}$ converges.

$$72. \sum_{n=3}^{\infty} \frac{1}{e^{\sqrt{n}}}$$

SOLUTION Apply the integral test, making the substitution $z = \sqrt{x}$ so that $z^2 = x$ and $2z dz = dx$:

$$\int_3^{\infty} \frac{1}{e^{\sqrt{x}}} dx = \int_3^{\infty} e^{-x^{1/2}} dx = \int_{\sqrt{3}}^{\infty} 2ze^{-z} dz$$

Evaluate this integral using integration by parts with $u = 2z$, $dv = e^{-z} dz$:

$$\begin{aligned} \int_{\sqrt{3}}^{\infty} 2ze^{-z} dz &= uv \Big|_{\sqrt{3}}^{\infty} - \int_{\sqrt{3}}^{\infty} v du = (-2ze^{-z}) \Big|_{\sqrt{3}}^{\infty} - \int_{\sqrt{3}}^{\infty} (-2e^{-z}) dz = 2\sqrt{3}e^{-\sqrt{3}} - (2e^{-z}) \Big|_{\sqrt{3}}^{\infty} \\ &= 2\sqrt{3}e^{-\sqrt{3}} + 2e^{-\sqrt{3}} < \infty \end{aligned}$$

Since the integral converges, so does the series $\sum_{n=3}^{\infty} \frac{1}{e^{\sqrt{n}}}$.

$$73. \sum_{n=4}^{\infty} \frac{\ln n}{n^2 - 3n}$$

SOLUTION By the comment preceding Exercise 31, we can choose $N \geq 4$ so that for $n \geq N$, $\ln n < n^{1/2}$. Then

$$\sum_{n=N}^{\infty} \frac{\ln n}{n^2 - 3n} \leq \sum_{n=N}^{\infty} \frac{n^{1/2}}{n^2 - 3n} = \sum_{n=N}^{\infty} \frac{1}{n^{3/2} - 3n^{1/2}}$$

To evaluate convergence of the latter series, let $a_n = \frac{1}{n^{3/2} - 3n^{1/2}}$ and $b_n = \frac{1}{n^{3/2}}$, and apply the Limit Comparison Test:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{n^{3/2} - 3n^{1/2}} \cdot n^{3/2} = \lim_{n \rightarrow \infty} \frac{1}{1 - 3n^{-1}} = 0$$

Thus $\sum a_n$ converges if $\sum b_n$ does. But $\sum b_n$ is a convergent p -series. Thus $\sum a_n$ converges and, by the comparison test, so does the original series. Adding back in the finite number of terms for $n < N$ does not affect convergence.

$$74. \sum_{n=1}^{\infty} \frac{1}{3^{\ln n}}$$

SOLUTION Note that

$$3^{\ln n} = (e^{\ln 3})^{\ln n} = (e^{\ln n})^{\ln 3} = n^{\ln 3}.$$

Thus the sum is a p -series with $p = \ln 3 > 1$, so is convergent.

$$75. \sum_{n=2}^{\infty} \frac{1}{n^{1/2} \ln n}$$

SOLUTION By the comment preceding Exercise 31, we can choose $N \geq 2$ so that for $n \geq N$, $\ln n < n^{1/4}$. Then

$$\sum_{n=N}^{\infty} \frac{1}{n^{1/2} \ln n} > \sum_{n=N}^{\infty} \frac{1}{n^{3/4}}$$

which is a divergent p -series. Thus the original series diverges as well - as usual, adding back in the finite number of terms for $n < N$ does not affect convergence.

$$76. \sum_{n=1}^{\infty} \frac{1}{n^{3/2} - \ln^4 n}$$

SOLUTION Let

$$a_n = \frac{1}{n^{3/2} - \ln^4 n}, \quad b_n = \frac{1}{n^{3/2}},$$

and apply the Limit Comparison Test:

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{n^{3/2}}{n^{3/2} - \ln^4 n} = \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{\ln^4 n}{n^{3/2}}}$$

But by the comment preceding Exercise 31, $\ln n$, and thus $\ln^4 n$, are eventually smaller than any positive power of n , so for n sufficiently large, $\frac{\ln^4 n}{n^{3/2}}$ is arbitrarily small. Thus $L = 1$ and $\sum a_n$ converges if and only if $\sum b_n$ does. But $\sum b_n$ is

a convergent p -series, so $\sum_{n=1}^{\infty} \frac{1}{n^{3/2} - \ln^4 n}$ converges.

$$77. \sum_{n=1}^{\infty} \frac{4n^2 + 15n}{3n^4 - 5n^2 - 17}$$

SOLUTION Apply the Limit Comparison Test with

$$a_n = \frac{4n^2 + 15n}{3n^4 - 5n^2 - 17}, \quad b_n = \frac{4n^2}{3n^4} = \frac{4}{3n^2}$$

We have

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{4n^2 + 15n}{3n^4 - 5n^2 - 17} \cdot \frac{3n^2}{4} = \lim_{n \rightarrow \infty} \frac{12n^4 + 45n^3}{12n^4 - 20n^2 - 68} = \lim_{n \rightarrow \infty} \frac{12 + 45/n}{12 - 20/n^2 - 68/n^4} = 1$$

Now, $\sum_{n=1}^{\infty} b_n$ is a p -series with $p = 2 > 1$, so converges. Since $L = 1$, we see that $\sum_{n=1}^{\infty} \frac{4n^2 + 15n}{3n^4 - 5n^2 - 17}$ converges as well.

$$78. \sum_{n=1}^{\infty} \frac{n}{4^{-n} + 5^{-n}}$$

SOLUTION Note that

$$\lim_{n \rightarrow \infty} \frac{n}{4^{-n} + 5^{-n}} = \lim_{n \rightarrow \infty} \frac{n4^n}{1 + \left(\frac{4}{5}\right)^n}$$

This limit approaches $\infty/1 = \infty$, so the terms of the sequence do not tend to zero. Thus the series is divergent.

79. For which a does $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^a}$ converge?

SOLUTION First consider the case $a > 0$ but $a \neq 1$. Let $f(x) = \frac{1}{x(\ln x)^a}$. This function is continuous, positive and decreasing for $x \geq 2$, so the Integral Test applies. Now,

$$\int_2^{\infty} \frac{dx}{x(\ln x)^a} = \lim_{R \rightarrow \infty} \int_2^R \frac{dx}{x(\ln x)^a} = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u^a} = \frac{1}{1-a} \lim_{R \rightarrow \infty} \left(\frac{1}{(\ln R)^{a-1}} - \frac{1}{(\ln 2)^{a-1}} \right).$$

Because

$$\lim_{R \rightarrow \infty} \frac{1}{(\ln R)^{a-1}} = \begin{cases} \infty, & 0 < a < 1 \\ 0, & a > 1 \end{cases}$$

we conclude the integral diverges when $0 < a < 1$ and converges when $a > 1$. Therefore

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^a} \text{ converges for } a > 1 \text{ and diverges for } 0 < a < 1.$$

Next, consider the case $a = 1$. The series becomes $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$. Let $f(x) = \frac{1}{x \ln x}$. For $x \geq 2$, this function is continuous, positive and decreasing, so the Integral Test applies. Using the substitution $u = \ln x$, $du = \frac{1}{x} dx$, we find

$$\int_2^{\infty} \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_2^R \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u} = \lim_{R \rightarrow \infty} (\ln(\ln R) - \ln(\ln 2)) = \infty.$$

The integral diverges; hence, the series also diverges.

Finally, consider the case $a < 0$. Let $b = -a > 0$ so the series becomes $\sum_{n=2}^{\infty} \frac{(\ln n)^b}{n}$. Since $\ln n > 1$ for all $n \geq 3$, it follows that

$$(\ln n)^b > 1 \quad \text{so} \quad \frac{(\ln n)^b}{n} > \frac{1}{n}.$$

The series $\sum_{n=3}^{\infty} \frac{1}{n}$ diverges, so by the Comparison Test we can conclude that $\sum_{n=3}^{\infty} \frac{(\ln n)^b}{n}$ also diverges. Consequently,

$\sum_{n=2}^{\infty} \frac{(\ln n)^b}{n}$ diverges. Thus,

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^a} \text{ diverges for } a < 0.$$

To summarize:

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^a} \text{ converges if } a > 1 \text{ and diverges if } a \leq 1.$$

80. For which a does $\sum_{n=2}^{\infty} \frac{1}{n^a \ln n}$ converge?

SOLUTION First consider the case $a > 1$. For $n \geq 3$, $\ln n > 1$ and

$$\frac{1}{n^a \ln n} < \frac{1}{n^a}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^a}$ is a p -series with $p = a > 1$, so it converges; hence, $\sum_{n=3}^{\infty} \frac{1}{n^a}$ also converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=3}^{\infty} \frac{1}{n^a \ln n}$ converges, which implies the series $\sum_{n=2}^{\infty} \frac{1}{n^a \ln n}$ also converges.

For $a \leq 1$, $n^a \leq n$ so

$$\frac{1}{n^a \ln n} \geq \frac{1}{n \ln n}$$

for $n \geq 2$. Let $f(x) = \frac{1}{x \ln x}$. For $x \geq 2$, this function is continuous, positive and decreasing, so the Integral Test applies.

Using the substitution $u = \ln x$, $du = \frac{1}{x} dx$, we find

$$\int_2^{\infty} \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_2^R \frac{dx}{x \ln x} = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u} = \lim_{R \rightarrow \infty} (\ln(\ln R) - \ln(\ln 2)) = \infty.$$

The integral diverges; hence, the series $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ also diverges. By the Comparison Test we can therefore conclude that

the series $\sum_{n=2}^{\infty} \frac{1}{n^a \ln n}$ diverges.

To summarize,

$$\sum_{n=2}^{\infty} \frac{1}{n^a \ln n} \text{ converges for } a > 1 \text{ and diverges for } a \leq 1.$$

Approximating Infinite Sums In Exercises 81–83, let $a_n = f(n)$, where $f(x)$ is a continuous, decreasing function such that $f(x) \geq 0$ and $\int_1^{\infty} f(x) dx$ converges.

81. Show that

$$\int_1^{\infty} f(x) dx \leq \sum_{n=1}^{\infty} a_n \leq a_1 + \int_1^{\infty} f(x) dx \quad \boxed{3}$$

SOLUTION From the proof of the Integral Test, we know that

$$a_2 + a_3 + a_4 + \cdots + a_N \leq \int_1^N f(x) dx \leq \int_1^{\infty} f(x) dx;$$

that is,

$$S_N - a_1 \leq \int_1^{\infty} f(x) dx \quad \text{or} \quad S_N \leq a_1 + \int_1^{\infty} f(x) dx.$$

Also from the proof of the Integral test, we know that

$$\int_1^N f(x) dx \leq a_1 + a_2 + a_3 + \cdots + a_{N-1} = S_N - a_N \leq S_N.$$

Thus,

$$\int_1^N f(x) dx \leq S_N \leq a_1 + \int_1^{\infty} f(x) dx.$$

Taking the limit as $N \rightarrow \infty$ yields Eq. (3), as desired.

82. *CRS* Using Eq. (3), show that

$$5 \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 6$$

This series converges slowly. Use a computer algebra system to verify that $S_N < 5$ for $N \leq 43,128$ and $S_{43,129} \approx 5.00000021$.

SOLUTION By Eq. (3), we have

$$\int_1^{\infty} \frac{dx}{x^{1.2}} \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 1 + \int_1^{\infty} \frac{dx}{x^{1.2}}.$$

Since

$$\int_1^{\infty} \frac{dx}{x^{1.2}} = \lim_{R \rightarrow \infty} \int_1^R \frac{dx}{x^{1.2}} = \lim_{R \rightarrow \infty} \left(\frac{1}{0.2} - \frac{R^{-0.2}}{0.2} \right) = 5,$$

it follows that

$$5 \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 6.$$

Because $a_n = n^{-1.2} \geq 0$ for all N , S_N is increasing and it suffices to show that $S_N < 5$ for $N = 43,128$ to conclude that $S_N < 5$ for all $N \leq 43,128$. Using a computer algebra system, we obtain:

$$S_{43,128} = \sum_{n=1}^{43,128} \frac{1}{n^{1.2}} = 4.9999974685$$

and

$$S_{43,129} = \sum_{n=1}^{43,129} \frac{1}{n^{1.2}} = 5.0000002118.$$

83. Let $S = \sum_{n=1}^{\infty} a_n$. Arguing as in Exercise 81, show that

$$\sum_{n=1}^M a_n + \int_{M+1}^{\infty} f(x) dx \leq S \leq \sum_{n=1}^{M+1} a_n + \int_{M+1}^{\infty} f(x) dx \quad \boxed{4}$$

Conclude that

$$0 \leq S - \left(\sum_{n=1}^M a_n + \int_{M+1}^{\infty} f(x) dx \right) \leq a_{M+1} \quad \boxed{5}$$

This provides a method for approximating S with an error of at most a_{M+1} .

SOLUTION Following the proof of the Integral Test and the argument in Exercise 81, but starting with $n = M + 1$ rather than $n = 1$, we obtain

$$\int_{M+1}^{\infty} f(x) dx \leq \sum_{n=M+1}^{\infty} a_n \leq a_{M+1} + \int_{M+1}^{\infty} f(x) dx.$$

Adding $\sum_{n=1}^M a_n$ to each part of this inequality yields

$$\sum_{n=1}^M a_n + \int_{M+1}^{\infty} f(x) dx \leq \sum_{n=1}^{\infty} a_n = S \leq \sum_{n=1}^{M+1} a_n + \int_{M+1}^{\infty} f(x) dx.$$

Subtracting $\sum_{n=1}^M a_n + \int_{M+1}^{\infty} f(x) dx$ from each part of this last inequality then gives us

$$0 \leq S - \left(\sum_{n=1}^M a_n + \int_{M+1}^{\infty} f(x) dx \right) \leq a_{M+1}.$$

84. *CRS* Use Eq. (4) with $M = 43,129$ to prove that

$$5.5915810 \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 5.5915839$$

SOLUTION Using Eq. (4) with $f(x) = \frac{1}{x^{1.2}}$, $a_n = \frac{1}{n^{1.2}}$ and $M = 43129$, we find

$$S_{43129} + \int_{43130}^{\infty} \frac{dx}{x^{1.2}} \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq S_{43130} + \int_{43130}^{\infty} \frac{dx}{x^{1.2}}.$$

Now,

$$S_{43129} = 5.0000002118;$$

$$S_{43130} = S_{43129} + \frac{1}{43130^{1.2}} = 5.0000029551;$$

and

$$\int_{43130}^{\infty} \frac{dx}{x^{1.2}} = \lim_{R \rightarrow \infty} \int_{43130}^R \frac{dx}{x^{1.2}} = -5 \lim_{R \rightarrow \infty} \left(\frac{1}{R^{0.2}} - \frac{1}{43130^{0.2}} \right) = \frac{5}{43130^{0.2}} = 0.5915808577.$$

Thus,

$$5.0000002118 + 0.5915808577 \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 5.0000029551 + 0.5915808577,$$

or

$$5.5915810695 \leq \sum_{n=1}^{\infty} \frac{1}{n^{1.2}} \leq 5.5915838128.$$

85. *CRS* Apply Eq. (4) with $M = 40,000$ to show that

$$1.644934066 \leq \sum_{n=1}^{\infty} \frac{1}{n^2} \leq 1.644934068$$

Is this consistent with Euler's result, according to which this infinite series has sum $\pi^2/6$?

SOLUTION Using Eq. (4) with $f(x) = \frac{1}{x^2}$, $a_n = \frac{1}{n^2}$ and $M = 40,000$, we find

$$S_{40,000} + \int_{40,001}^{\infty} \frac{dx}{x^2} \leq \sum_{n=1}^{\infty} \frac{1}{n^2} \leq S_{40,001} + \int_{40,001}^{\infty} \frac{dx}{x^2}.$$

Now,

$$S_{40,000} = 1.6449090672;$$

$$S_{40,001} = S_{40,000} + \frac{1}{40,001} = 1.6449090678;$$

and

$$\int_{40,001}^{\infty} \frac{dx}{x^2} = \lim_{R \rightarrow \infty} \int_{40,001}^R \frac{dx}{x^2} = - \lim_{R \rightarrow \infty} \left(\frac{1}{R} - \frac{1}{40,001} \right) = \frac{1}{40,001} = 0.0000249994.$$

Thus,

$$1.6449090672 + 0.0000249994 \leq \sum_{n=1}^{\infty} \frac{1}{n^2} \leq 1.6449090678 + 0.0000249994,$$

or

$$1.6449340665 \leq \sum_{n=1}^{\infty} \frac{1}{n^2} \leq 1.6449340672.$$

Since $\frac{\pi^2}{6} \approx 1.6449340668$, our approximation is consistent with Euler's result.

86. CAS Using a CAS and Eq. (5), determine the value of $\sum_{n=1}^{\infty} n^{-6}$ to within an error less than 10^{-4} . Check that your result is consistent with that of Euler, who proved that the sum is equal to $\pi^6/945$.

SOLUTION According to Eq. (5), if we choose M so that $(M+1)^{-6} < 10^{-4}$, we can then approximate the sum to within 10^{-4} . Solving $(M+1)^{-6} = 10^{-4}$ gives $M+1 = 10^{-2/3} \approx 4.641$, so the smallest such integral M is $M = 4$. Denote by S the sum of the series. Then

$$0 \leq S - \left(\sum_{n=1}^4 n^{-6} + \int_5^{\infty} x^{-6} dx \right) \leq (M+1)^{-6} < 10^{-4}$$

We have

$$\begin{aligned} \sum_{n=1}^4 n^{-6} &= \frac{1}{1} + \frac{1}{64} + \frac{1}{729} + \frac{1}{4096} \approx 1.017240883 \\ \int_5^{\infty} x^{-6} dx &= -\frac{1}{5}x^{-5} \Big|_5^{\infty} = \frac{1}{5^6} \approx 0.000064 \end{aligned}$$

The sum of these two is ≈ 1.017304883 , while $\frac{\pi^6}{945} \approx 1.017343063$. These two values differ by approximately $0.000038180 < 10^{-4}$, so the result is consistent with Euler's calculation.

87. CAS Using a CAS and Eq. (5), determine the value of $\sum_{n=1}^{\infty} n^{-5}$ to within an error less than 10^{-4} .

SOLUTION Using Eq. (5) with $f(x) = x^{-5}$ and $a_n = n^{-5}$, we have

$$0 \leq \sum_{n=1}^{\infty} n^{-5} - \left(\sum_{n=1}^{M+1} n^{-5} + \int_{M+1}^{\infty} x^{-5} dx \right) \leq (M+1)^{-5}.$$

To guarantee an error less than 10^{-4} , we need $(M+1)^{-5} \leq 10^{-4}$. This yields $M \geq 10^{4/5} - 1 \approx 5.3$, so we choose $M = 6$. Now,

$$\sum_{n=1}^7 n^{-5} = 1.0368498887,$$

and

$$\int_7^{\infty} x^{-5} dx = \lim_{R \rightarrow \infty} \int_7^R x^{-5} dx = -\frac{1}{4} \lim_{R \rightarrow \infty} (R^{-4} - 7^{-4}) = \frac{1}{4 \cdot 7^4} = 0.0001041233.$$

Thus,

$$\sum_{n=1}^{\infty} n^{-5} \approx \sum_{n=1}^7 n^{-5} + \int_7^{\infty} x^{-5} dx = 1.0368498887 + 0.0001041233 = 1.0369540120.$$

88. How far can a stack of identical books (of mass m and unit length) extend without tipping over? The stack will not tip over if the $(n+1)$ st book is placed at the bottom of the stack with its right edge located at the center of mass of the first n books (Figure 5). Let c_n be the center of mass of the first n books, measured along the x -axis, where we take the positive x -axis to the left of the origin as in Figure 6. Recall that if an object of mass m_1 has center of mass at x_1 and a second object of mass m_2 has center of mass x_2 , then the center of mass of the system has x -coordinate

$$\frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$$

- Show that if the $(n+1)$ st book is placed with its right edge at c_n , then its center of mass is located at $c_n + \frac{1}{2}$.
- Consider the first n books as a single object of mass nm with center of mass at c_n and the $(n+1)$ st book as a second object of mass m . Show that if the $(n+1)$ st book is placed with its right edge at c_n , then $c_{n+1} = c_n + \frac{1}{2(n+1)}$.
- Prove that $\lim_{n \rightarrow \infty} c_n = \infty$. Thus, by using enough books, the stack can be extended as far as desired without tipping over.




FIGURE 5




FIGURE 6

SOLUTION Let $f(x) = \frac{1}{x}$. This function is continuous, positive and decreasing, so following the argument of Exercise 81, we know that

$$\int_1^N f(x) dx \leq S_N \leq a_1 + \int_1^N f(x) dx,$$

or

$$\ln N \leq 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{N} \leq 1 + \ln N.$$

Using the inequality on the right-hand side, we know that

$$S_{8100} \leq 1 + \ln 8100 = 9.999619 < 10;$$

using the inequality on the left-hand side, we can guarantee $S_N \geq 100$ by making $\ln N \geq 100$. Thus, we can take

$$N \geq e^{100} \approx 2.688 \times 10^{43}.$$

89. The following argument proves the divergence of the harmonic series $S = \sum_{n=1}^{\infty} 1/n$ without using the Integral Test.

Let

$$S_1 = 1 + \frac{1}{3} + \frac{1}{5} + \cdots, \quad S_2 = \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \cdots$$

Show that if S converges, then

(a) S_1 and S_2 also converge and $S = S_1 + S_2$.

(b) $S_1 > S_2$ and $S_2 = \frac{1}{2}S$.

Observe that (b) contradicts (a), and conclude that S diverges.

SOLUTION Assume throughout that S converges; we will derive a contradiction. Write

$$a_n = \frac{1}{n}, \quad b_n = \frac{1}{2n-1}, \quad c_n = \frac{1}{2n}$$

for the n^{th} terms in the series S , S_1 , and S_2 . Since $2n-1 \geq n$ for $n \geq 1$, we have $b_n < a_n$. Since $S = \sum a_n$ converges, so does $S_1 = \sum b_n$ by the Comparison Test. Also, $c_n = \frac{1}{2}a_n$, so again by the Comparison Test, the convergence of S implies the convergence of $S_2 = \sum c_n$. Now, define two sequences

$$b'_n = \begin{cases} b_{(n+1)/2} & n \text{ odd} \\ 0 & n \text{ even} \end{cases}$$

$$c'_n = \begin{cases} 0 & n \text{ odd} \\ c_{n/2} & n \text{ even} \end{cases}$$

That is, b'_n and c'_n look like b_n and c_n , but have zeros inserted in the “missing” places compared to a_n . Then $a_n = b'_n + c'_n$; also $S_1 = \sum b_n = \sum b'_n$ and $S_2 = \sum c_n = \sum c'_n$. Finally, since S , S_1 , and S_2 all converge, we have

$$S = \sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (b'_n + c'_n) = \sum_{n=1}^{\infty} b'_n + \sum_{n=1}^{\infty} c'_n = \sum_{n=1}^{\infty} b_n + \sum_{n=1}^{\infty} c_n = S_1 + S_2$$

Now, $b_n > c_n$ for every n , so that $S_1 > S_2$. Also, we showed above that $c_n = \frac{1}{2}a_n$, so that $2S_2 = S$. Putting all this together gives

$$S = S_1 + S_2 > S_2 + S_2 = 2S_2 = S$$

so that $S > S$, a contradiction. Thus S must diverge.

Further Insights and Challenges

90. Let $S = \sum_{n=2}^{\infty} a_n$, where $a_n = (\ln(\ln n))^{-\ln n}$.

- (a) Show, by taking logarithms, that $a_n = n^{-\ln(\ln(\ln n))}$.
 (b) Show that $\ln(\ln(\ln n)) \geq 2$ if $n > C$, where $C = e^{e^{e^2}}$.
 (c) Show that S converges.

SOLUTION

(a) Let $a_n = (\ln(\ln n))^{-\ln n}$. Then

$$\ln a_n = (-\ln n) \ln(\ln(\ln n)),$$

and

$$a_n = e^{(-\ln n) \ln(\ln(\ln n))} = \left(e^{\ln n}\right)^{-\ln(\ln(\ln n))} = n^{-\ln(\ln(\ln n))}.$$

(b) Suppose $n > e^{e^{e^2}}$. Then

$$\ln n > \ln e^{e^{e^2}} = e^{e^2};$$

$$\ln(\ln n) > \ln e^{e^2} = e^2; \text{ and}$$

$$\ln(\ln(\ln n)) > \ln e^2 = 2.$$

(c) Combining the results from parts (a) and (b), we have

$$a_n = \frac{1}{n^{\ln(\ln(\ln n))}} \leq \frac{1}{n^2}$$

for $n > C = e^{e^{e^2}}$. The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series, so $\sum_{n=C+1}^{\infty} \frac{1}{n^2}$ also converges. By the Comparison Test we can therefore conclude that the series $\sum_{n=C+1}^{\infty} a_n$ converges, which means that the series $\sum_{n=2}^{\infty} a_n$ converges.

91. **Kummer's Acceleration Method** Suppose we wish to approximate $S = \sum_{n=1}^{\infty} 1/n^2$. There is a similar telescoping series whose value can be computed exactly (Example 1 in Section 10.2):

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$$

(a) Verify that

$$S = \sum_{n=1}^{\infty} \frac{1}{n(n+1)} + \sum_{n=1}^{\infty} \left(\frac{1}{n^2} - \frac{1}{n(n+1)} \right)$$

Thus for M large,

$$S \approx 1 + \sum_{n=1}^M \frac{1}{n^2(n+1)} \quad \boxed{6}$$

(b) Explain what has been gained. Why is Eq. (6) a better approximation to S than is $\sum_{n=1}^M 1/n^2$?

(c) *CRS* Compute

$$\sum_{n=1}^{1000} \frac{1}{n^2}, \quad 1 + \sum_{n=1}^{100} \frac{1}{n^2(n+1)}$$

Which is a better approximation to S , whose exact value is $\pi^2/6$?

SOLUTION

(a) Because the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ and $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$ both converge,

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} + \sum_{n=1}^{\infty} \left(\frac{1}{n^2} - \frac{1}{n(n+1)} \right) = \sum_{n=1}^{\infty} \frac{1}{n(n+1)} + \sum_{n=1}^{\infty} \frac{1}{n^2} - \sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \sum_{n=1}^{\infty} \frac{1}{n^2} = S.$$

Now,

$$\frac{1}{n^2} - \frac{1}{n(n+1)} = \frac{n+1}{n^2(n+1)} - \frac{n}{n^2(n+1)} = \frac{1}{n^2(n+1)},$$

so, for M large,

$$S \approx 1 + \sum_{n=1}^M \frac{1}{n^2(n+1)}.$$

(b) The series $\sum_{n=1}^{\infty} \frac{1}{n^2(n+1)}$ converges more rapidly than $\sum_{n=1}^{\infty} \frac{1}{n^2}$ since the degree of n in the denominator is larger.

(c) Using a computer algebra system, we find

$$\sum_{n=1}^{1000} \frac{1}{n^2} = 1.6439345667 \quad \text{and} \quad 1 + \sum_{n=1}^{100} \frac{1}{n^2(n+1)} = 1.6448848903.$$

The second sum is more accurate because it is closer to the exact solution $\frac{\pi^2}{6} \approx 1.6449340668$.

92. CAS The series $S = \sum_{k=1}^{\infty} k^{-3}$ has been computed to more than 100 million digits. The first 30 digits are

$$S = 1.202056903159594285399738161511$$

Approximate S using the Acceleration Method of Exercise 91 with $M = 100$ and auxiliary series

$$R = \sum_{n=1}^{\infty} (n(n+1)(n+2))^{-1}.$$

According to Exercise 46 in Section 10.2, R is a telescoping series with the sum $R = \frac{1}{4}$.

SOLUTION We compute the difference between the general term of the given series and the general term of the auxiliary series:

$$\frac{1}{k^3} - \frac{1}{k(k+1)(k+2)} = \frac{(k+1)(k+2) - k^2}{k^3(k+1)(k+2)} = \frac{k^2 + 3k + 2 - k^2}{k^3(k+1)(k+2)} = \frac{3k+2}{k^3(k+1)(k+2)}$$

Hence,

$$\sum_{k=1}^{\infty} \frac{1}{k^3} = \sum_{k=1}^{\infty} \frac{1}{k(k+1)(k+2)} + \sum_{k=1}^{\infty} \frac{3k+2}{k^3(k+1)(k+2)} = \frac{1}{4} + \sum_{k=1}^{\infty} \frac{3k+2}{k^3(k+1)(k+2)}$$

With $M = 100$ and using a computer algebra system, we find

$$\sum_{k=1}^{\infty} \frac{1}{k^3} \approx \frac{1}{4} + \sum_{k=1}^{100} \frac{3k+2}{k^3(k+1)(k+2)} = 1.2020559349.$$

10.4 Absolute and Conditional Convergence

Preliminary Questions

1. Give an example of a series such that $\sum a_n$ converges but $\sum |a_n|$ diverges.

SOLUTION The series $\sum \frac{(-1)^n}{\sqrt[3]{n}}$ converges by the Leibniz Test, but the positive series $\sum \frac{1}{\sqrt[3]{n}}$ is a divergent p -series.

2. Which of the following statements is equivalent to Theorem 1?

- (a) If $\sum_{n=0}^{\infty} |a_n|$ diverges, then $\sum_{n=0}^{\infty} a_n$ also diverges.
 (b) If $\sum_{n=0}^{\infty} a_n$ diverges, then $\sum_{n=0}^{\infty} |a_n|$ also diverges.
 (c) If $\sum_{n=0}^{\infty} a_n$ converges, then $\sum_{n=0}^{\infty} |a_n|$ also converges.

SOLUTION The correct answer is (b): If $\sum_{n=0}^{\infty} a_n$ diverges, then $\sum_{n=0}^{\infty} |a_n|$ also diverges. Take $a_n = (-1)^n \frac{1}{n}$ to see that statements (a) and (c) are not true in general.

3. Lathika argues that $\sum_{n=1}^{\infty} (-1)^n \sqrt{n}$ is an alternating series and therefore converges. Is Lathika right?

SOLUTION No. Although $\sum_{n=1}^{\infty} (-1)^n \sqrt{n}$ is an alternating series, the terms $a_n = \sqrt{n}$ do not form a decreasing sequence that tends to zero. In fact, $a_n = \sqrt{n}$ is an increasing sequence that tends to ∞ , so $\sum_{n=1}^{\infty} (-1)^n \sqrt{n}$ diverges by the Divergence Test.

4. Suppose that a_n is positive, decreasing, and tends to 0, and let $S = \sum_{n=1}^{\infty} (-1)^{n-1} a_n$. What can we say about $|S - S_{100}|$ if $a_{101} = 10^{-3}$? Is S larger or smaller than S_{100} ?

SOLUTION From the text, we know that $|S - S_{100}| < a_{101} = 10^{-3}$. Also, the Leibniz test tells us that $S_{2N} < S < S_{2N+1}$ for any $N \geq 1$, so that $S_{100} < S$.

Exercises

1. Show that

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{2^n}$$

converges absolutely.

SOLUTION The positive series $\sum_{n=0}^{\infty} \frac{1}{2^n}$ is a geometric series with $r = \frac{1}{2}$. Thus, the positive series converges, and the given series converges absolutely.

2. Show that the following series converges conditionally:

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n^{2/3}} = \frac{1}{1^{2/3}} - \frac{1}{2^{2/3}} + \frac{1}{3^{2/3}} - \frac{1}{4^{2/3}} + \cdots$$

SOLUTION Let $a_n = \frac{1}{n^{2/3}}$. Then a_n forms a decreasing sequence that tends to zero; hence, the series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n^{2/3}}$ converges by the Leibniz Test. However, the positive series $\sum_{n=1}^{\infty} \frac{1}{n^{2/3}}$ is a divergent p -series, so the original series converges conditionally.

In Exercises 3–10, determine whether the series converges absolutely, conditionally, or not at all.

$$3. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{1/3}}$$

SOLUTION The sequence $a_n = \frac{1}{n^{1/3}}$ is positive, decreasing, and tends to zero; hence, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{1/3}}$ converges by the Leibniz Test. However, the positive series $\sum_{n=1}^{\infty} \frac{1}{n^{1/3}}$ is a divergent p -series, so the original series converges conditionally.

$$4. \sum_{n=1}^{\infty} \frac{(-1)^n n^4}{n^3 + 1}$$

SOLUTION Because

$$\lim_{n \rightarrow \infty} \frac{n^4}{n^3 + 1} = \infty,$$

the general term $\frac{(-1)^n n^4}{n^3 + 1}$ of the series does not tend to zero; hence, this series diverges by the Divergence Test.

$$5. \sum_{n=0}^{\infty} \frac{(-1)^{n-1}}{(1.1)^n}$$

SOLUTION The positive series $\sum_{n=0}^{\infty} \left(\frac{1}{1.1}\right)^n$ is a convergent geometric series; thus, the original series converges absolutely.

$$6. \sum_{n=1}^{\infty} \frac{\sin(\frac{\pi n}{4})}{n^2}$$

SOLUTION Because

$$\left| \frac{\sin(\frac{\pi n}{4})}{n^2} \right| = \frac{|\sin(\frac{\pi n}{4})|}{n^2} \leq \frac{1}{n^2}$$

the positive series forms a convergent p -series; thus, the original series converges absolutely.

$$7. \sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$$

SOLUTION Let $a_n = \frac{1}{n \ln n}$. Then a_n forms a decreasing sequence (note that n and $\ln n$ are both increasing functions of n) that tends to zero; hence, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$ converges by the Leibniz Test. However, the positive series $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ diverges, so the original series converges conditionally.

$$8. \sum_{n=1}^{\infty} \frac{(-1)^n}{1 + \frac{1}{n}}$$

SOLUTION Because

$$\lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} = \frac{1}{1 + 0} = 1,$$

the general term $\frac{(-1)^n}{1 + \frac{1}{n}}$ of the series does not tend to zero; hence, the series diverges by the Divergent Test.

$$9. \sum_{n=2}^{\infty} \frac{\cos n\pi}{(\ln n)^2}$$

SOLUTION Since $\cos n\pi$ alternates between $+1$ and -1 ,

$$\sum_{n=2}^{\infty} \frac{\cos n\pi}{(\ln n)^2} = \sum_{n=2}^{\infty} \frac{(-1)^n}{(\ln n)^2}$$

This is an alternating series whose general term decreases to zero, so it converges. The associated positive series,

$$\sum_{n=2}^{\infty} \frac{1}{(\ln n)^2}$$

is a divergent series, so the original series converges conditionally.

10.
$$\sum_{n=1}^{\infty} \frac{\cos n}{2^n}$$

SOLUTION The associated positive series is

$$\sum_{n=1}^{\infty} \frac{|\cos n|}{2^n} \leq \sum_{n=1}^{\infty} \frac{1}{2^n}$$

which is a convergent geometric series. Thus the associated positive series converges, so the original series converges absolutely.

11. Let
$$S = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^3}.$$

- (a) Calculate S_n for $1 \leq n \leq 10$.
 (b) Use Eq. (2) to show that $0.9 \leq S \leq 0.902$.

SOLUTION

(a)

$$\begin{array}{ll} S_1 = 1 & S_6 = S_5 - \frac{1}{6^3} = 0.899782407 \\ S_2 = 1 - \frac{1}{2^3} = \frac{7}{8} = 0.875 & S_7 = S_6 + \frac{1}{7^3} = 0.902697859 \\ S_3 = S_2 + \frac{1}{3^3} = 0.912037037 & S_8 = S_7 - \frac{1}{8^3} = 0.900744734 \\ S_4 = S_3 - \frac{1}{4^3} = 0.896412037 & S_9 = S_8 + \frac{1}{9^3} = 0.902116476 \\ S_5 = S_4 + \frac{1}{5^3} = 0.904412037 & S_{10} = S_9 - \frac{1}{10^3} = 0.901116476 \end{array}$$

(b) By Eq. (2),

$$|S_{10} - S| \leq a_{11} = \frac{1}{11^3},$$

so

$$S_{10} - \frac{1}{11^3} \leq S \leq S_{10} + \frac{1}{11^3},$$

or

$$0.900365161 \leq S \leq 0.901867791.$$

12. Use Eq. (2) to approximate

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n!}$$

to four decimal places.

SOLUTION Let
$$S = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n!},$$
 so that $a_n = \frac{1}{n!}$. By Eq. (2),

$$|S_N - S| \leq a_{N+1} = \frac{1}{(N+1)!}.$$

To guarantee accuracy to four decimal places, we must choose N so that

$$\frac{1}{(N+1)!} < 5 \times 10^{-5} \quad \text{or} \quad (N+1)! > 20,000.$$

Because $7! = 5040$ and $8! = 40,320$, the smallest value that satisfies the required inequality is $N = 7$. Thus,

$$S \approx S_7 = 1 - \frac{1}{2!} + \frac{1}{3!} - \frac{1}{4!} + \frac{1}{5!} - \frac{1}{6!} + \frac{1}{7!} = 0.632142857.$$

13. Approximate $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^4}$ to three decimal places.

SOLUTION Let $S = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^4}$, so that $a_n = \frac{1}{n^4}$. By Eq. (2),

$$|S_N - S| \leq a_{N+1} = \frac{1}{(N+1)^4}.$$

To guarantee accuracy to three decimal places, we must choose N so that

$$\frac{1}{(N+1)^4} < 5 \times 10^{-4} \quad \text{or} \quad N > \sqrt[4]{2000} - 1 \approx 5.7.$$

The smallest value that satisfies the required inequality is then $N = 6$. Thus,


$$S \approx S_6 = 1 - \frac{1}{2^4} + \frac{1}{3^4} - \frac{1}{4^4} + \frac{1}{5^4} - \frac{1}{6^4} = 0.946767824.$$

14. *CAS* Let

$$S = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{n^2 + 1}$$

Use a computer algebra system to calculate and plot the partial sums S_n for $1 \leq n \leq 100$. Observe that the partial sums zigzag above and below the limit.

SOLUTION The partial sums associated with the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n}{n^2 + 1}$ are plotted below. As expected, the partial sums alternate between overestimating and underestimating the sum.



In Exercises 15 and 16, find a value of N such that S_N approximates the series with an error of at most 10^{-5} . If you have a CAS, compute this value of S_N .

15. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+2)(n+3)}$

SOLUTION Let $S = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+2)(n+3)}$, so that $a_n = \frac{1}{n(n+2)(n+3)}$. By Eq. (2),

$$|S_N - S| \leq a_{N+1} = \frac{1}{(N+1)(N+3)(N+4)}.$$

We must choose N so that

$$\frac{1}{(N+1)(N+3)(N+4)} \leq 10^{-5} \quad \text{or} \quad (N+1)(N+3)(N+4) \geq 10^5.$$

For $N = 43$, the product on the left hand side is 95,128, while for $N = 44$ the product is 101,520; hence, the smallest value of N which satisfies the required inequality is $N = 44$. Thus,

$$S \approx S_{44} = \sum_{n=1}^{44} \frac{(-1)^{n+1}}{n(n+2)(n+3)} = 0.0656746.$$

$$16. \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \ln n}{n!}$$

SOLUTION Let $S = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \ln n}{n!}$, so that $a_n = \frac{\ln n}{n!}$. By Eq. (2),

$$|S_N - S| \leq a_{N+1} = \frac{\ln(N+1)}{(N+1)!}.$$

To make the error at most 10^{-5} , we must choose N so that

$$\frac{\ln(N+1)}{(N+1)!} \leq 10^{-5}.$$

For $N = 7$, the left-hand side of the above inequality is 5.157×10^{-5} , while for $N = 8$, the left-hand side is 6.055×10^{-6} ; hence, the smallest value for N which satisfies the required inequality is $N = 8$. Thus,

$$S \approx S_8 = \sum_{n=1}^8 \frac{(-1)^{n+1} \ln n}{n!} = -0.209975859.$$

In Exercises 17–32, determine convergence or divergence by any method.

$$17. \sum_{n=0}^{\infty} 7^{-n}$$

SOLUTION This is a (positive) geometric series with $r = \frac{1}{7} < 1$, so it converges.

$$18. \sum_{n=1}^{\infty} \frac{1}{n^{7.5}}$$

SOLUTION This is a p -series with $p = 7.5 > 1$, so it converges.

$$19. \sum_{n=1}^{\infty} \frac{1}{5^n - 3^n}$$

SOLUTION Use the Limit Comparison Test with $\frac{1}{5^n}$:

$$L = \lim_{n \rightarrow \infty} \frac{1/(5^n - 3^n)}{1/5^n} = \lim_{n \rightarrow \infty} \frac{5^n}{5^n - 3^n} = \lim_{n \rightarrow \infty} \frac{1}{1 - (3/5)^n} = 1$$

But $\sum_{n=1}^{\infty} \frac{1}{5^n}$ is a convergent geometric series. Since $L = 1$, the Limit Comparison Test tells us that the original series converges as well.

$$20. \sum_{n=2}^{\infty} \frac{n}{n^2 - n}$$

SOLUTION Apply the Limit Comparison Test and compare with the divergent harmonic series:

$$L = \lim_{n \rightarrow \infty} \frac{\frac{n}{n^2 - n}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n^2}{n^2 - n} = 1.$$

Because $L > 0$, we conclude that the series $\sum_{n=2}^{\infty} \frac{n}{n^2 - n}$ diverges.

$$21. \sum_{n=1}^{\infty} \frac{1}{3n^4 + 12n}$$

SOLUTION Use the Limit Comparison Test with $\frac{1}{3n^4}$:

$$L = \lim_{n \rightarrow \infty} \frac{(1/(3n^4 + 12n))}{1/3n^4} = \lim_{n \rightarrow \infty} \frac{3n^4}{3n^4 + 12n} = \lim_{n \rightarrow \infty} \frac{1}{1 + 4n^{-3}} = 1$$

But $\sum_{n=1}^{\infty} \frac{1}{3n^4} = \frac{1}{3} \sum_{n=1}^{\infty} \frac{1}{n^4}$ is a convergent p -series. Since $L = 1$, the Limit Comparison Test tells us that the original series converges as well.

$$22. \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^2 + 1}}$$

SOLUTION This is an alternating series with $a_n = \frac{1}{\sqrt{n^2 + 1}}$. Because a_n is a decreasing sequence that converges to

zero, the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^2 + 1}}$ converges by the Leibniz Test.

$$23. \sum_{n=1}^{\infty} \frac{1}{\sqrt{n^2 + 1}}$$

SOLUTION Apply the Limit Comparison Test and compare the series with the divergent harmonic series:

$$L = \lim_{n \rightarrow \infty} \frac{\frac{1}{\sqrt{n^2 + 1}}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2 + 1}} = 1.$$

Because $L > 0$, we conclude that the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n^2 + 1}}$ diverges.

$$24. \sum_{n=0}^{\infty} \frac{(-1)^n n}{\sqrt{n^2 + 1}}$$

SOLUTION This series diverges, since the general term of the associated positive series tends to 1, not to 0:

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2 + 1}} = \lim_{n \rightarrow \infty} \sqrt{\frac{n^2}{n^2 + 1}} = \lim_{n \rightarrow \infty} \sqrt{\frac{1}{1 + n^{-2}}} = 1$$

$$25. \sum_{n=1}^{\infty} \frac{3^n + (-2)^n}{5^n}$$

SOLUTION The series

$$\sum_{n=1}^{\infty} \frac{3^n}{5^n} = \sum_{n=1}^{\infty} \left(\frac{3}{5}\right)^n$$

is a convergent geometric series, as is the series

$$\sum_{n=1}^{\infty} \frac{(-1)^n 2^n}{5^n} = \sum_{n=1}^{\infty} \left(-\frac{2}{5}\right)^n.$$

Hence,

$$\sum_{n=1}^{\infty} \frac{3^n + (-1)^n 2^n}{5^n} = \sum_{n=1}^{\infty} \left(\frac{3}{5}\right)^n + \sum_{n=1}^{\infty} \left(-\frac{2}{5}\right)^n$$

also converges.

$$26. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n+1)!}$$

SOLUTION This is an alternating series with $a_n = \frac{1}{(2n+1)!}$. Because a_n is a decreasing sequence which converges to

zero, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n+1)!}$ converges by the Leibniz Test.

$$27. \sum_{n=1}^{\infty} (-1)^n n^2 e^{-n^3/3}$$

SOLUTION Consider the associated positive series $\sum_{n=1}^{\infty} n^2 e^{-n^3/3}$. This series can be seen to converge by the Integral Test:

$$\int_1^{\infty} x^2 e^{-x^3/3} dx = \lim_{R \rightarrow \infty} \int_1^R x^2 e^{-x^3/3} dx = - \lim_{R \rightarrow \infty} e^{-x^3/3} \Big|_1^R = e^{-1/3} + \lim_{R \rightarrow \infty} e^{-R^3/3} = e^{-1/3}.$$

The integral converges, so the original series converges absolutely.

$$28. \sum_{n=1}^{\infty} n e^{-n^3/3}$$

SOLUTION This is a positive series, and by the Comparison Test with the associated positive series in the previous exercise,

$$\sum_{n=1}^{\infty} n e^{-n^3/3} \leq \sum_{n=1}^{\infty} n^2 e^{-n^3/3}$$

Since the series on the right converges, so does the original series.

$$29. \sum_{n=2}^{\infty} \frac{(-1)^n}{n^{1/2} (\ln n)^2}$$

SOLUTION This is an alternating series with $a_n = \frac{1}{n^{1/2} (\ln n)^2}$. Because a_n is a decreasing sequence which converges to zero, the series $\sum_{n=2}^{\infty} \frac{(-1)^n}{n^{1/2} (\ln n)^2}$ converges by the Leibniz Test. (Note that the series converges only conditionally, not absolutely; the associated positive series is eventually greater than $\frac{1}{n^{3/4}}$, which is a divergent p -series).

$$30. \sum_{n=2}^{\infty} \frac{1}{n (\ln n)^{1/4}}$$

SOLUTION Use the Integral Test, with the substitution $u = \ln x$:

$$\begin{aligned} \int_2^{\infty} \frac{1}{x \ln^{1/4} x} dx &= \lim_{R \rightarrow \infty} \int_2^R \frac{1}{x \ln^{1/4} x} dx = \lim_{R \rightarrow \infty} \int_{\ln 2}^R u^{-1/4} du = \lim_{R \rightarrow \infty} \frac{4}{3} u^{3/4} \Big|_{\ln 2}^R \\ &= -\frac{4}{3} \left((\ln 2)^{3/4} + \lim_{R \rightarrow \infty} R^{3/4} \right) \end{aligned}$$

The integral diverges, so the original series diverges as well.

$$31. \sum_{n=1}^{\infty} \frac{\ln n}{n^{1.05}}$$

SOLUTION Choose N so that for $n \geq N$ we have $\ln n \leq n^{0.01}$. Then

$$\sum_{n=N}^{\infty} \frac{\ln n}{n^{1.05}} \leq \sum_{n=N}^{\infty} \frac{n^{0.01}}{n^{1.05}} = \sum_{n=N}^{\infty} \frac{1}{n^{1.04}}$$

This is a convergent p -series, so by the Comparison Test, the original series converges as well.

$$32. \sum_{n=2}^{\infty} \frac{1}{(\ln n)^2}$$

SOLUTION Choose N so that for $n \geq N$ we have $\ln n < n^{0.25}$ so that $\ln^2 n < n^{0.5}$. Then

$$\sum_{n=N}^{\infty} \frac{1}{(\ln n)^2} > \sum_{n=N}^{\infty} \frac{1}{n^{0.5}}$$

This is a divergent p -series, so by the Comparison Test, the original series diverges as well.

33. Show that

$$S = \frac{1}{2} - \frac{1}{2} + \frac{1}{3} - \frac{1}{3} + \frac{1}{4} - \frac{1}{4} + \cdots$$

converges by computing the partial sums. Does it converge absolutely?

SOLUTION The sequence of partial sums is

$$S_1 = \frac{1}{2}$$

$$S_2 = S_1 - \frac{1}{2} = 0$$

$$S_3 = S_2 + \frac{1}{3} = \frac{1}{3}$$

$$S_4 = S_3 - \frac{1}{3} = 0$$

and, in general,

$$S_N = \begin{cases} \frac{1}{N}, & \text{for odd } N \\ 0, & \text{for even } N \end{cases}$$

Thus, $\lim_{N \rightarrow \infty} S_N = 0$, and the series converges to 0. The positive series is

$$\frac{1}{2} + \frac{1}{2} + \frac{1}{3} + \frac{1}{3} + \frac{1}{4} + \frac{1}{4} + \cdots = 2 \sum_{n=2}^{\infty} \frac{1}{n};$$

which diverges. Therefore, the original series converges conditionally, not absolutely.

34. The Leibniz Test cannot be applied to

$$\frac{1}{2} - \frac{1}{3} + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{2^3} - \frac{1}{3^3} + \cdots$$

Why not? Show that it converges by another method.

SOLUTION The sequence of terms $\{a_n\}$ for this alternating series is

$$\frac{1}{2}, \frac{1}{3}, \frac{1}{2^2}, \frac{1}{3^2}, \frac{1}{2^3}, \frac{1}{3^3}, \dots, \frac{1}{2^n}, \frac{1}{3^n}, \frac{1}{2^{n+1}}, \frac{1}{3^{n+1}}, \dots$$

Now,

$$\frac{1}{3^2} = \frac{1}{9} < \frac{1}{8} = \frac{1}{2^3}.$$

Moreover, if we assume that

$$\frac{1}{3^k} < \frac{1}{2^{k+1}}$$

for some k , then

$$\frac{1}{3^{k+1}} = \frac{1}{3} \cdot \frac{1}{3^k} < \frac{1}{3} \cdot \frac{1}{2^{k+1}} < \frac{1}{2} \cdot \frac{1}{2^{k+1}} = \frac{1}{2^{k+2}}.$$

Thus, by mathematical induction,

$$\frac{1}{3^n} < \frac{1}{2^{n+1}}$$

for all $n \geq 2$. The sequence $\{a_n\}$ is therefore not decreasing, and the Leibniz Test does not apply.

We may express the given series as


$$\sum_{n=1}^{\infty} \left(\frac{1}{2^n} - \frac{1}{3^n} \right).$$

Because

$$\sum_{n=1}^{\infty} \frac{1}{2^n} = \sum_{n=1}^{\infty} \left(\frac{1}{2} \right)^n \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{3^n} = \sum_{n=1}^{\infty} \left(\frac{1}{3} \right)^n$$

are both convergent geometric series, it follows that this series converges, and

$$\sum_{n=1}^{\infty} \left(\frac{1}{2^n} - \frac{1}{3^n} \right) = \sum_{n=1}^{\infty} \left(\frac{1}{2} \right)^n - \sum_{n=1}^{\infty} \left(\frac{1}{3} \right)^n = \frac{\frac{1}{2}}{1 - \frac{1}{2}} - \frac{\frac{1}{3}}{1 - \frac{1}{3}} = 1 - \frac{1}{2} = \frac{1}{2}.$$

35.  **Assumptions Matter** Show by counterexample that the Leibniz Test does not remain true if the sequence a_n tends to zero but is not assumed nonincreasing. *Hint:* Consider

$$R = \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{8} + \frac{1}{4} - \frac{1}{16} + \cdots + \left(\frac{1}{n} - \frac{1}{2^n} \right) + \cdots$$

SOLUTION Let

$$R = \frac{1}{2} - \frac{1}{4} + \frac{1}{3} - \frac{1}{8} + \frac{1}{4} - \frac{1}{16} + \cdots + \left(\frac{1}{n+1} - \frac{1}{2^{n+1}} \right) + \cdots$$

This is an alternating series with

$$a_n = \begin{cases} \frac{1}{k+1}, & n = 2k-1 \\ \frac{1}{2^{k+1}}, & n = 2k \end{cases}$$

Note that $a_n \rightarrow 0$ as $n \rightarrow \infty$, but the sequence $\{a_n\}$ is not decreasing. We will now establish that R diverges.

For sake of contradiction, suppose that R converges. The geometric series

$$\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$$

converges, so the sum of R and this geometric series must also converge; however,

$$R + \sum_{n=1}^{\infty} \frac{1}{2^{n+1}} = \sum_{n=2}^{\infty} \frac{1}{n},$$

which diverges because the harmonic series diverges. Thus, the series R must diverge.

36. Determine whether the following series converges conditionally:

$$1 - \frac{1}{3} + \frac{1}{2} - \frac{1}{5} + \frac{1}{3} - \frac{1}{7} + \frac{1}{4} - \frac{1}{9} + \frac{1}{5} - \frac{1}{11} + \cdots$$

SOLUTION Although the signs alternate, the terms a_n are not decreasing, so we cannot apply the Leibniz Test. However, we may express the series as

$$\sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{2n+1} \right) = \sum_{n=1}^{\infty} \frac{n+1}{n(2n+1)}.$$

Using the Limit Comparison Test and comparing with the harmonic series, we find

$$L = \lim_{n \rightarrow \infty} \frac{\frac{n+1}{n(2n+1)}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n+1}{2n+1} = \frac{1}{2}.$$

Because $L > 0$, we conclude that the series

$$1 - \frac{1}{3} + \frac{1}{2} - \frac{1}{5} + \frac{1}{3} - \frac{1}{7} + \frac{1}{4} - \frac{1}{9} + \frac{1}{5} - \frac{1}{11} + \cdots$$

diverges.

37. Prove that if $\sum a_n$ converges absolutely, then $\sum a_n^2$ also converges. Then give an example where $\sum a_n$ is only conditionally convergent and $\sum a_n^2$ diverges.

SOLUTION Suppose the series $\sum a_n$ converges absolutely. Because $\sum |a_n|$ converges, we know that

$$\lim_{n \rightarrow \infty} |a_n| = 0.$$

Therefore, there exists a positive integer N such that $|a_n| < 1$ for all $n \geq N$. It then follows that for $n \geq N$,

$$0 \leq a_n^2 = |a_n|^2 = |a_n| \cdot |a_n| < |a_n| \cdot 1 = |a_n|.$$

By the Comparison Test we can then conclude that $\sum a_n^2$ also converges.

Consider the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$. This series converges by the Leibniz Test, but the corresponding positive series is a divergent p -series; that is, $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$ is conditionally convergent. Now, $\sum_{n=1}^{\infty} a_n^2$ is the divergent harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$. Thus, $\sum a_n^2$ need not converge if $\sum a_n$ is only conditionally convergent.

Further Insights and Challenges

38. Prove the following variant of the Leibniz Test: If $\{a_n\}$ is a positive, decreasing sequence with $\lim_{n \rightarrow \infty} a_n = 0$, then the series

$$a_1 + a_2 - 2a_3 + a_4 + a_5 - 2a_6 + \cdots$$

converges. *Hint:* Show that S_{3N} is increasing and bounded by $a_1 + a_2$, and continue as in the proof of the Leibniz Test.

SOLUTION Following the hint, we first examine the sequence $\{S_{3N}\}$. Now,

$$S_{3N+3} = S_{3(N+1)} = S_{3N} + a_{3N+1} + a_{3N+2} - 2a_{3N+3} = S_{3N} + (a_{3N+1} - a_{3N+3}) + (a_{3N+2} - a_{3N+3}) \geq S_{3N}$$

because $\{a_n\}$ is a decreasing sequence. Moreover,

$$\begin{aligned} S_{3N} &= a_1 + a_2 - \sum_{k=1}^{N-1} (2a_{3k} - a_{3k+1} - a_{3k+2}) - 2a_{3N} \\ &= a_1 + a_2 - \sum_{k=1}^{N-1} [(a_{3k} - a_{3k+1}) + (a_{3k} - a_{3k+2}) - 2a_{3N}] \leq a_1 + a_2 \end{aligned}$$

again because $\{a_n\}$ is a decreasing sequence. Thus, $\{S_{3N}\}$ is an increasing sequence with an upper bound; hence, $\{S_{3N}\}$ converges. Next,

$$S_{3N+1} = S_{3N} + a_{3N+1} \quad \text{and} \quad S_{3N+2} = S_{3N} + a_{3N+1} + a_{3N+2}.$$

Given that $\lim_{n \rightarrow \infty} a_n = 0$, it follows that

$$\lim_{N \rightarrow \infty} S_{3N+1} = \lim_{N \rightarrow \infty} S_{3N+2} = \lim_{N \rightarrow \infty} S_{3N}.$$

Having just established that $\lim_{N \rightarrow \infty} S_{3N}$ exists, it follows that the sequences $\{S_{3N+1}\}$ and $\{S_{3N+2}\}$ converge to the same limit. Finally, we can conclude that the sequence of partial sums $\{S_N\}$ converges, so the given series converges.

39. Use Exercise 38 to show that the following series converges:

$$S = \frac{1}{\ln 2} + \frac{1}{\ln 3} - \frac{2}{\ln 4} + \frac{1}{\ln 5} + \frac{1}{\ln 6} - \frac{2}{\ln 7} + \cdots$$

SOLUTION The given series has the structure of the generic series from Exercise 38 with $a_n = \frac{1}{\ln(n+1)}$. Because a_n is a positive, decreasing sequence with $\lim_{n \rightarrow \infty} a_n = 0$, we can conclude from Exercise 38 that the given series converges.

40. Prove the conditional convergence of

$$R = 1 + \frac{1}{2} + \frac{1}{3} - \frac{3}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} - \frac{3}{8} + \cdots$$

SOLUTION Using Exercise 38 as a template, we first examine the sequence $\{R_{4N}\}$. Now,

$$\begin{aligned} R_{4N+4} &= R_{4(N+1)} = R_{4N} + \frac{1}{4N+1} + \frac{1}{4N+2} + \frac{1}{4N+3} - \frac{3}{4N+4} \\ &= R_{4N} + \left(\frac{1}{4N+1} - \frac{1}{4N+4} \right) + \left(\frac{1}{4N+2} - \frac{1}{4N+4} \right) + \left(\frac{1}{4N+3} - \frac{1}{4N+4} \right) \geq R_{4N}. \end{aligned}$$

Moreover,

$$R_{4N} = 1 + \frac{1}{2} + \frac{1}{3} - \sum_{k=1}^{N-1} \left(\frac{3}{4k} - \frac{1}{4k+1} - \frac{1}{4k+2} - \frac{1}{4k+3} \right) - \frac{3}{4N} \leq 1 + \frac{1}{2} + \frac{1}{3}.$$

Thus, $\{R_{4N}\}$ is an increasing sequence with an upper bound; hence, $\{R_{4N}\}$ converges. Next,

$$\begin{aligned}R_{4N+1} &= R_{4N} + \frac{1}{4N+1}; \\R_{4N+2} &= R_{4N} + \frac{1}{4N+1} + \frac{1}{4N+2}; \text{ and} \\R_{4N+3} &= R_{4N} + \frac{1}{4N+1} + \frac{1}{4N+2} + \frac{1}{4N+3},\end{aligned}$$

so

$$\lim_{n \rightarrow \infty} R_{4N+1} = \lim_{N \rightarrow \infty} R_{4N+2} = \lim_{N \rightarrow \infty} R_{4N+3} = \lim_{N \rightarrow \infty} R_{4N}.$$

Having just established that $\lim_{N \rightarrow \infty} R_{4N}$ exists, it follows that the sequences $\{R_{4N+1}\}$, $\{R_{4N+2}\}$ and $\{R_{4N+3}\}$ converge to the same limit. Finally, we can conclude that the sequence of partial sums $\{R_N\}$ converges, so the series R converges.

Now, consider the positive series

$$R^+ = 1 + \frac{1}{2} + \frac{1}{3} + \frac{3}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{3}{8} + \cdots$$

Because the terms in this series are greater than or equal to the corresponding terms in the divergent harmonic series, it follows from the Comparison Test that R^+ diverges. Thus, by definition, R converges conditionally.

41. Show that the following series diverges:

$$S = 1 + \frac{1}{2} + \frac{1}{3} - \frac{2}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} - \frac{2}{8} + \cdots$$

Hint: Use the result of Exercise 40 to write S as the sum of a convergent series and a divergent series.

SOLUTION Let

$$R = 1 + \frac{1}{2} + \frac{1}{3} - \frac{3}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} - \frac{3}{8} + \cdots$$

and

$$S = 1 + \frac{1}{2} + \frac{1}{3} - \frac{2}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} - \frac{2}{8} + \cdots$$

For sake of contradiction, suppose the series S converges. From Exercise 40, we know that the series R converges. Thus, the series $S - R$ must converge; however,

$$S - R = \frac{1}{4} + \frac{1}{8} + \frac{1}{12} + \cdots = \frac{1}{4} \sum_{k=1}^{\infty} \frac{1}{k},$$

which diverges because the harmonic series diverges. Thus, the series S must diverge.

42. Prove that

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(\ln n)^a}{n}$$

converges for all exponents a . *Hint:* Show that $f(x) = (\ln x)^a/x$ is decreasing for x sufficiently large.

SOLUTION This is an alternating series with $a_n = \frac{(\ln n)^a}{n}$. Following the hint, consider the function $f(x) = \frac{(\ln x)^a}{x}$. Now,

$$f'(x) = \frac{a(\ln x)^{a-1} - (\ln x)^a}{x^2} = \frac{(\ln x)^{a-1}}{x^2} (a - \ln x),$$

so $f'(x) < 0$ and f is decreasing for $x > e^a$. If $a \leq 0$, then it is clear that

$$\lim_{x \rightarrow \infty} \frac{(\ln x)^a}{x} = 0;$$

if $a > 0$, then repeated use of L'Hôpital's Rule leads to the same conclusion. Let N be any integer greater than e^a ; then,

$\{a_n\}$ is a decreasing sequence for $n \geq N$ which converges to zero and the series $\sum_{n=N}^{\infty} (-1)^{n+1} \frac{(\ln n)^a}{n}$ converges by the

Leibniz Test. Finally, the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(\ln n)^a}{n}$ also converges.

43. We say that $\{b_n\}$ is a rearrangement of $\{a_n\}$ if $\{b_n\}$ has the same terms as $\{a_n\}$ but occurring in a different order. Show that if $\{b_n\}$ is a rearrangement of $\{a_n\}$ and $S = \sum_{n=1}^{\infty} a_n$ converges absolutely, then $T = \sum_{n=1}^{\infty} b_n$ also converges absolutely.

(This result does not hold if S is only conditionally convergent.) *Hint:* Prove that the partial sums $\sum_{n=1}^N |b_n|$ are bounded.

It can be shown further that $S = T$.

SOLUTION Suppose the series $S = \sum_{n=1}^{\infty} a_n$ converges absolutely and denote the corresponding positive series by

$$S^+ = \sum_{n=1}^{\infty} |a_n|.$$

Further, let $T_N = \sum_{n=1}^N |b_n|$ denote the N th partial sum of the series $\sum_{n=1}^{\infty} |b_n|$. Because $\{b_n\}$ is a rearrangement of $\{a_n\}$, we know that

$$0 \leq T_N \leq \sum_{n=1}^{\infty} |a_n| = S^+;$$

that is, the sequence $\{T_N\}$ is bounded. Moreover,

$$T_{N+1} = \sum_{n=1}^{N+1} |b_n| = T_N + |b_{N+1}| \geq T_N;$$

that is, $\{T_N\}$ is increasing. It follows that $\{T_N\}$ converges, so the series $\sum_{n=1}^{\infty} |b_n|$ converges, which means the series $\sum_{n=1}^{\infty} b_n$ converges absolutely.

44. Assumptions Matter In 1829, Lejeune Dirichlet pointed out that the great French mathematician Augustin Louis Cauchy made a mistake in a published paper by improperly assuming the Limit Comparison Test to be valid for nonpositive series. Here are Dirichlet's two series:

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}, \quad \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}} \left(1 + \frac{(-1)^n}{\sqrt{n}}\right)$$

Explain how they provide a counterexample to the Limit Comparison Test when the series are not assumed to be positive.

SOLUTION Let

$$R = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}} \quad \text{and} \quad S = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}} \left(1 + \frac{(-1)^n}{\sqrt{n}}\right)$$

R is an alternating series that converges by the Leibniz Test; however, we cannot apply the Leibniz Test to S because the absolute value of the terms in S is not decreasing. Because

$$L = \lim_{n \rightarrow \infty} \frac{\frac{(-1)^n}{\sqrt{n}} \left(1 + \frac{(-1)^n}{\sqrt{n}}\right)}{\frac{(-1)^n}{\sqrt{n}}} = \lim_{n \rightarrow \infty} \left(1 + \frac{(-1)^n}{\sqrt{n}}\right) = 1,$$

if the Limit Comparison Test were valid for nonpositive series, we would conclude that S converges. However, if we assume that S converges, then the series $S - R$ would also converge. But

$$S - R = \sum_{n=1}^{\infty} \left(\frac{(-1)^n}{\sqrt{n}} + \frac{1}{n} - \frac{(-1)^n}{\sqrt{n}}\right) = \sum_{n=1}^{\infty} \frac{1}{n},$$

which is the divergent harmonic series. Thus, S diverges, and the Limit Comparison Test is not valid for nonpositive series.

10.5 The Ratio and Root Tests

Preliminary Questions

1. In the Ratio Test, is ρ equal to $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ or $\lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$?

SOLUTION In the Ratio Test ρ is the limit $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$.

2. Is the Ratio Test conclusive for $\sum_{n=1}^{\infty} \frac{1}{2^n}$? Is it conclusive for $\sum_{n=1}^{\infty} \frac{1}{n}$?

SOLUTION The general term of $\sum_{n=1}^{\infty} \frac{1}{2^n}$ is $a_n = \frac{1}{2^n}$; thus,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2^{n+1}} \cdot \frac{2^n}{1} = \frac{1}{2},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2} < 1.$$

Consequently, the Ratio Test guarantees that the series $\sum_{n=1}^{\infty} \frac{1}{2^n}$ converges.

The general term of $\sum_{n=1}^{\infty} \frac{1}{n}$ is $a_n = \frac{1}{n}$; thus,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{n+1} \cdot \frac{n}{1} = \frac{n}{n+1},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} = 1.$$

The Ratio Test is therefore inconclusive for the series $\sum_{n=1}^{\infty} \frac{1}{n}$.

3. Can the Ratio Test be used to show convergence if the series is only conditionally convergent?

SOLUTION No. The Ratio Test can only establish absolute convergence and divergence, not conditional convergence.

Exercises

In Exercises 1–20, apply the Ratio Test to determine convergence or divergence, or state that the Ratio Test is inconclusive.

1. $\sum_{n=1}^{\infty} \frac{1}{5^n}$

SOLUTION With $a_n = \frac{1}{5^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{5^{n+1}} \cdot \frac{5^n}{1} = \frac{1}{5} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{5} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{1}{5^n}$ converges by the Ratio Test.

2. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1} n}{5^n}$

SOLUTION With $a_n = \frac{(-1)^{n-1} n}{5^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{n+1}{5^{n+1}} \cdot \frac{5^n}{n} = \frac{n+1}{5n} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{5} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1} n}{5^n}$ converges by the Ratio Test.

$$3. \sum_{n=1}^{\infty} \frac{1}{n^n}$$

SOLUTION With $a_n = \frac{1}{n^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{(n+1)^{n+1}} \cdot \frac{n^n}{1} = \frac{1}{n+1} \left(\frac{n}{n+1} \right)^n = \frac{1}{n+1} \left(1 + \frac{1}{n} \right)^{-n},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 \cdot \frac{1}{e} = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{1}{n^n}$ converges by the Ratio Test.

$$4. \sum_{n=0}^{\infty} \frac{3n+2}{5n^3+1}$$

SOLUTION With $a_n = \frac{3n+2}{5n^3+1}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{3(n+1)+2}{5(n+1)^3+1} \cdot \frac{5n^3+1}{3n+2} = \frac{3n+5}{3n+2} \cdot \frac{5n^3+1}{5(n+1)^3+1},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot 1 = 1.$$

Therefore, for the series $\sum_{n=0}^{\infty} \frac{3n+2}{5n^3+1}$, the Ratio Test is inconclusive.

We can show that this series converges by using the Limit Comparison Test and comparing with the convergent p -series

$$\sum_{n=1}^{\infty} \frac{1}{n^2}.$$

$$5. \sum_{n=1}^{\infty} \frac{n}{n^2+1}$$

SOLUTION With $a_n = \frac{n}{n^2+1}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{n+1}{(n+1)^2+1} \cdot \frac{n^2+1}{n} = \frac{n+1}{n} \cdot \frac{n^2+1}{n^2+2n+2},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot 1 = 1.$$

Therefore, for the series $\sum_{n=1}^{\infty} \frac{n}{n^2+1}$, the Ratio Test is inconclusive.

We can show that this series diverges by using the Limit Comparison Test and comparing with the divergent harmonic series.

$$6. \sum_{n=1}^{\infty} \frac{2^n}{n}$$

SOLUTION With $a_n = \frac{2^n}{n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{n+1}}{n+1} \cdot \frac{n}{2^n} = \frac{2n}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 2 > 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{2^n}{n}$ diverges by the Ratio Test.

$$7. \sum_{n=1}^{\infty} \frac{2^n}{n^{100}}$$

SOLUTION With $a_n = \frac{2^n}{n^{100}}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{n+1}}{(n+1)^{100}} \cdot \frac{n^{100}}{2^n} = 2 \left(\frac{n}{n+1} \right)^{100} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 2 \cdot 1^{100} = 2 > 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{2^n}{n^{100}}$ diverges by the Ratio Test.

$$8. \sum_{n=1}^{\infty} \frac{n^3}{3n^2}$$

SOLUTION With $a_n = \frac{n^3}{3n^2}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^3}{3(n+1)^2} \cdot \frac{3n^2}{n^3} = \left(\frac{n+1}{n} \right)^3 \cdot \frac{1}{3} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1^3 \cdot 0 = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n^3}{3n^2}$ converges by the Ratio Test.

$$9. \sum_{n=1}^{\infty} \frac{10^n}{2n^2}$$

SOLUTION With $a_n = \frac{10^n}{2n^2}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{10^{n+1}}{2(n+1)^2} \cdot \frac{2n^2}{10^n} = 10 \cdot \frac{1}{2} \cdot \frac{n^2}{(n+1)^2} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 10 \cdot 0 = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{10^n}{2n^2}$ converges by the Ratio Test.

$$10. \sum_{n=1}^{\infty} \frac{e^n}{n!}$$

SOLUTION With $a_n = \frac{e^n}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{e^{n+1}}{(n+1)!} \cdot \frac{n!}{e^n} = \frac{e}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{e^n}{n!}$ converges by the Ratio Test.

$$11. \sum_{n=1}^{\infty} \frac{e^n}{n^n}$$

SOLUTION With $a_n = \frac{e^n}{n^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{e^{n+1}}{(n+1)^{n+1}} \cdot \frac{n^n}{e^n} = \frac{e}{n+1} \left(\frac{n}{n+1} \right)^n = \frac{e}{n+1} \left(1 + \frac{1}{n} \right)^{-n},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 \cdot \frac{1}{e} = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{e^n}{n^n}$ converges by the Ratio Test.

$$12. \sum_{n=1}^{\infty} \frac{n^{40}}{n!}$$

SOLUTION With $a_n = \frac{n^{40}}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^{40}}{(n+1)!} \cdot \frac{n!}{n^{40}} = \frac{1}{n+1} \left(\frac{n+1}{n} \right)^{40} = \frac{1}{n+1} \left(1 + \frac{1}{n} \right)^{40},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 \cdot 1 = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n^{40}}{n!}$ converges by the Ratio Test.

$$13. \sum_{n=0}^{\infty} \frac{n!}{6^n}$$

SOLUTION With $a_n = \frac{n!}{6^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)!}{6^{n+1}} \cdot \frac{6^n}{n!} = \frac{n+1}{6} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty > 1.$$

Therefore, the series $\sum_{n=0}^{\infty} \frac{n!}{6^n}$ diverges by the Ratio Test.

$$14. \sum_{n=1}^{\infty} \frac{n!}{n^9}$$

SOLUTION With $a_n = \frac{n!}{n^9}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)!}{(n+1)^9} \cdot \frac{n^9}{n!} = \frac{n^9}{(n+1)^8} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty > 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n!}{n^9}$ diverges by the Ratio Test.

$$15. \sum_{n=2}^{\infty} \frac{1}{n \ln n}$$

SOLUTION With $a_n = \frac{1}{n \ln n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{(n+1) \ln(n+1)} \cdot \frac{n \ln n}{1} = \frac{n}{n+1} \frac{\ln n}{\ln(n+1)},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot \lim_{n \rightarrow \infty} \frac{\ln n}{\ln(n+1)}.$$

Now,

$$\lim_{n \rightarrow \infty} \frac{\ln n}{\ln(n+1)} = \lim_{x \rightarrow \infty} \frac{\ln x}{\ln(x+1)} = \lim_{x \rightarrow \infty} \frac{1/(x+1)}{1/x} = \lim_{x \rightarrow \infty} \frac{x}{x+1} = 1.$$

Thus, $\rho = 1$, and the Ratio Test is inconclusive for the series $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$.

Using the Integral Test, we can show that the series $\sum_{n=2}^{\infty} \frac{1}{n \ln n}$ diverges.

$$16. \sum_{n=1}^{\infty} \frac{1}{(2n)!}$$

SOLUTION With $a_n = \frac{1}{(2n)!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{(2n+2)!} \cdot \frac{(2n)!}{1} = \frac{1}{(2n+2)(2n+1)} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{1}{(2n)!}$ converges by the Ratio Test.

$$17. \sum_{n=1}^{\infty} \frac{n^2}{(2n+1)!}$$

SOLUTION With $a_n = \frac{n^2}{(2n+1)!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^2}{(2n+3)!} \cdot \frac{(2n+1)!}{n^2} = \left(\frac{n+1}{n} \right)^2 \frac{1}{(2n+3)(2n+2)},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1^2 \cdot 0 = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n^2}{(2n+1)!}$ converges by the Ratio Test.

$$18. \sum_{n=1}^{\infty} \frac{(n!)^3}{(3n)!}$$

SOLUTION With $a_n = \frac{(n!)^3}{(3n)!}$,

$$\begin{aligned} \left| \frac{a_{n+1}}{a_n} \right| &= \frac{((n+1)!)^3}{(3(n+1))!} \cdot \frac{(3n)!}{(n!)^3} = \frac{(n+1)^3}{(3n+3)(3n+2)(3n+1)} = \frac{n^3 + 3n^2 + 3n + 1}{27n^3 + 54n^2 + 33n + 6} \\ &= \frac{1 + 3n^{-1} + 3n^{-2} + 1n^{-3}}{27 + 54n^{-1} + 33n^{-2} + 6n^{-3}} \end{aligned}$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{27} < 1$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{(n!)^3}{(3n)!}$ converges by the Ratio Test.

$$19. \sum_{n=2}^{\infty} \frac{1}{2^n + 1}$$

SOLUTION With $a_n = \frac{1}{2^n + 1}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2^{n+1} + 1} \cdot \frac{2^n + 1}{1} = \frac{1 + 2^{-n}}{2 + 2^{-n}}$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2} < 1$$

Therefore, the series $\sum_{n=2}^{\infty} \frac{1}{2^n + 1}$ converges by the Ratio Test.

$$20. \sum_{n=2}^{\infty} \frac{1}{\ln n}$$

SOLUTION With $a_n = \frac{1}{\ln n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{\ln n} \cdot \frac{\ln(n+1)}{1} = \frac{\ln(n+1)}{\ln n}$$

and (using L'Hôpital's rule)

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} \ln(x+1)}{\frac{d}{dx} \ln x} = \lim_{x \rightarrow \infty} \frac{x}{x+1} = 1$$

Therefore, the Ratio Test is inconclusive for $\sum_{n=2}^{\infty} \frac{1}{\ln n}$. This series can be shown to diverge using the Comparison Test with the harmonic series since $\ln n < n$ for $n \geq 2$.

21. Show that $\sum_{n=1}^{\infty} n^k 3^{-n}$ converges for all exponents k .

SOLUTION With $a_n = n^k 3^{-n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^k 3^{-(n+1)}}{n^k 3^{-n}} = \frac{1}{3} \left(1 + \frac{1}{n} \right)^k,$$

and, for all k ,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{3} \cdot 1 = \frac{1}{3} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} n^k 3^{-n}$ converges for all exponents k by the Ratio Test.

22. Show that $\sum_{n=1}^{\infty} n^2 x^n$ converges if $|x| < 1$.

SOLUTION With $a_n = n^2 x^n$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^2 |x|^{n+1}}{n^2 |x|^n} = \left(1 + \frac{1}{n} \right)^2 |x| \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot |x| = |x|.$$

Therefore, by the Ratio Test, the series $\sum_{n=1}^{\infty} n^2 x^n$ converges provided $|x| < 1$.

23. Show that $\sum_{n=1}^{\infty} 2^n x^n$ converges if $|x| < \frac{1}{2}$.

SOLUTION With $a_n = 2^n x^n$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{n+1} |x|^{n+1}}{2^n |x|^n} = 2|x| \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 2|x|.$$

Therefore, $\rho < 1$ and the series $\sum_{n=1}^{\infty} 2^n x^n$ converges by the Ratio Test provided $|x| < \frac{1}{2}$.

24. Show that $\sum_{n=1}^{\infty} \frac{r^n}{n!}$ converges for all r .

SOLUTION With $a_n = \frac{r^n}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|r|^{n+1}}{(n+1)!} \cdot \frac{n!}{|r|^n} = \frac{|r|}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 \cdot |r| = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{r^n}{n!}$ converges by the Ratio Test for all r .

25. Show that $\sum_{n=1}^{\infty} \frac{r^n}{n}$ converges if $|r| < 1$.

SOLUTION With $a_n = \frac{r^n}{n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|r|^{n+1}}{n+1} \cdot \frac{n}{|r|^n} = |r| \frac{n}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot |r| = |r|.$$

Therefore, by the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{r^n}{n}$ converges provided $|r| < 1$.

26. Is there any value of k such that $\sum_{n=1}^{\infty} \frac{2^n}{n^k}$ converges?

SOLUTION With $a_n = \frac{2^n}{n^k}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{n+1}}{(n+1)^k} \cdot \frac{n^k}{2^n} = 2 \left(\frac{n}{n+1} \right)^k,$$

and, for all k ,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 2 \cdot 1^k = 2 > 1.$$

Therefore, by the Ratio Test, there is no value for k such that the series $\sum_{n=1}^{\infty} \frac{2^n}{n^k}$ converges.

27. Show that $\sum_{n=1}^{\infty} \frac{n!}{n^n}$ converges. *Hint:* Use $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$.

SOLUTION With $a_n = \frac{n!}{n^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)!}{(n+1)^{n+1}} \cdot \frac{n^n}{n!} = \left(\frac{n}{n+1} \right)^n = \left(1 + \frac{1}{n} \right)^{-n},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{e} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n!}{n^n}$ converges by the Ratio Test.

In Exercises 28–33, assume that $|a_{n+1}/a_n|$ converges to $\rho = \frac{1}{3}$. What can you say about the convergence of the given series?

28. $\sum_{n=1}^{\infty} na_n$

SOLUTION Let $b_n = na_n$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \frac{n+1}{n} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot \frac{1}{3} = \frac{1}{3} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} na_n$ converges by the Ratio Test.

29. $\sum_{n=1}^{\infty} n^3 a_n$

SOLUTION Let $b_n = n^3 a_n$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \left(\frac{n+1}{n} \right)^3 \left| \frac{a_{n+1}}{a_n} \right| = 1^3 \cdot \frac{1}{3} = \frac{1}{3} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} n^3 a_n$ converges by the Ratio Test.

$$30. \sum_{n=1}^{\infty} 2^n a_n$$

SOLUTION Let $b_n = 2^n a_n$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \frac{2^{n+1}}{2^n} \left| \frac{a_{n+1}}{a_n} \right| = 2 \cdot \frac{1}{3} = \frac{2}{3} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} 2^n a_n$ converges by the Ratio Test.

$$31. \sum_{n=1}^{\infty} 3^n a_n$$

SOLUTION Let $b_n = 3^n a_n$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \frac{3^{n+1}}{3^n} \left| \frac{a_{n+1}}{a_n} \right| = 3 \cdot \frac{1}{3} = 1.$$

Therefore, the Ratio Test is inconclusive for the series $\sum_{n=1}^{\infty} 3^n a_n$.

$$32. \sum_{n=1}^{\infty} 4^n a_n$$

SOLUTION Let $b_n = 4^n a_n$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \frac{4^{n+1}}{4^n} \left| \frac{a_{n+1}}{a_n} \right| = 4 \cdot \frac{1}{3} = \frac{4}{3} > 1.$$

Therefore, the series $\sum_{n=1}^{\infty} 4^n a_n$ diverges by the Ratio Test.

$$33. \sum_{n=1}^{\infty} a_n^2$$

SOLUTION Let $b_n = a_n^2$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|^2 = \left(\frac{1}{3} \right)^2 = \frac{1}{9} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} a_n^2$ converges by the Ratio Test.

34. Assume that $|a_{n+1}/a_n|$ converges to $\rho = 4$. Does $\sum_{n=1}^{\infty} a_n^{-1}$ converge (assume that $a_n \neq 0$ for all n)?

SOLUTION Let $b_n = a_n^{-1}$. Then

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{b_{n+1}}{b_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \frac{1}{\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|} = \frac{1}{4} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} a_n^{-1}$ converges by the Ratio Test.

35. Is the Ratio Test conclusive for the p -series $\sum_{n=1}^{\infty} \frac{1}{n^p}$?

SOLUTION With $a_n = \frac{1}{n^p}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{(n+1)^p} \cdot \frac{n^p}{1} = \left(\frac{n}{n+1} \right)^p \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1^p = 1.$$

Therefore, the Ratio Test is inconclusive for the p -series $\sum_{n=1}^{\infty} \frac{1}{n^p}$.

In Exercises 36–41, use the Root Test to determine convergence or divergence (or state that the test is inconclusive).

$$36. \sum_{n=0}^{\infty} \frac{1}{10^n}$$

SOLUTION With $a_n = \frac{1}{10^n}$,

$$\sqrt[n]{a_n} = \sqrt[n]{\frac{1}{10^n}} = \frac{1}{10} \quad \text{and} \quad \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \frac{1}{10} < 1.$$

Therefore, the series $\sum_{n=0}^{\infty} \frac{1}{10^n}$ converges by the Root Test.

$$37. \sum_{n=1}^{\infty} \frac{1}{n^n}$$

SOLUTION With $a_n = \frac{1}{n^n}$,

$$\sqrt[n]{a_n} = \sqrt[n]{\frac{1}{n^n}} = \frac{1}{n} \quad \text{and} \quad \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{1}{n^n}$ converges by the Root Test.

$$38. \sum_{k=0}^{\infty} \left(\frac{k}{k+10} \right)^k$$

SOLUTION With $a_k = \left(\frac{k}{k+10} \right)^k$,

$$\sqrt[k]{a_k} = \sqrt[k]{\left(\frac{k}{k+10} \right)^k} = \frac{k}{k+10} \quad \text{and} \quad \lim_{k \rightarrow \infty} \sqrt[k]{a_k} = 1.$$

Therefore, the Root Test is inconclusive for the series $\sum_{k=0}^{\infty} \left(\frac{k}{k+10} \right)^k$. Because

$$\lim_{k \rightarrow \infty} a_k = \lim_{k \rightarrow \infty} \left(1 + \frac{10}{k} \right)^{-k} = \lim_{k \rightarrow \infty} \left[\left(1 + \frac{10}{k} \right)^{k/10} \right]^{-10} = e^{-10} \neq 0,$$

this series diverges by the Divergence Test.

$$39. \sum_{k=0}^{\infty} \left(\frac{k}{3k+1} \right)^k$$

SOLUTION With $a_k = \left(\frac{k}{3k+1} \right)^k$,

$$\sqrt[k]{a_k} = \sqrt[k]{\left(\frac{k}{3k+1} \right)^k} = \frac{k}{3k+1} \quad \text{and} \quad \lim_{k \rightarrow \infty} \sqrt[k]{a_k} = \frac{1}{3} < 1.$$

Therefore, the series $\sum_{k=0}^{\infty} \left(\frac{k}{3k+1} \right)^k$ converges by the Root Test.

$$40. \sum_{n=1}^{\infty} \left(1 + \frac{1}{n} \right)^{-n}$$

SOLUTION With $a_n = \left(1 + \frac{1}{n} \right)^{-n}$,

$$\sqrt[n]{a_n} = \sqrt[n]{\left(1 + \frac{1}{n} \right)^{-n}} = \left(1 + \frac{1}{n} \right)^{-1} \quad \text{and} \quad \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 1^{-1} = 1.$$

Therefore, the Root Test is inconclusive for the series $\sum_{n=1}^{\infty} \left(1 + \frac{1}{n} \right)^{-n}$.

Because

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^{-n} = \lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{n}\right)^n\right]^{-1} = e^{-1} \neq 0,$$

this series diverges by the Divergence Test.

$$41. \sum_{n=4}^{\infty} \left(1 + \frac{1}{n}\right)^{-n^2}$$

SOLUTION With $a_k = \left(1 + \frac{1}{n}\right)^{-n^2}$,

$$\sqrt[n]{a_n} = \sqrt[n]{\left(1 + \frac{1}{n}\right)^{-n^2}} = \left(1 + \frac{1}{n}\right)^{-n} \quad \text{and} \quad \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = e^{-1} < 1.$$

Therefore, the series $\sum_{n=4}^{\infty} \left(1 + \frac{1}{n}\right)^{-n^2}$ converges by the Root Test.

42. Prove that $\sum_{n=1}^{\infty} \frac{2^{n^2}}{n!}$ diverges. *Hint:* Use $2^{n^2} = (2^n)^n$ and $n! \leq n^n$.

SOLUTION Because $n! \leq n^n$,

$$\frac{2^{n^2}}{n!} \geq \frac{2^{n^2}}{n^n}.$$

Now, let $a_n = \frac{2^{n^2}}{n^n}$. Then

$$\sqrt[n]{a_n} = \sqrt[n]{\frac{2^{n^2}}{n^n}} = \frac{2^n}{n},$$

and

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \frac{2^n}{n} = \lim_{x \rightarrow \infty} \frac{2^x}{x} = \lim_{x \rightarrow \infty} \frac{2^x \ln 2}{1} = \infty > 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{2^{n^2}}{n^n}$ diverges by the Root Test. By the Comparison Test, we can then conclude that the series

$$\sum_{n=1}^{\infty} \frac{2^{n^2}}{n!} \text{ also diverges.}$$

In Exercises 43–56, determine convergence or divergence using any method covered in the text so far.

$$43. \sum_{n=1}^{\infty} \frac{2^n + 4^n}{7^n}$$

SOLUTION Because the series

$$\sum_{n=1}^{\infty} \frac{2^n}{7^n} = \sum_{n=1}^{\infty} \left(\frac{2}{7}\right)^n \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{4^n}{7^n} = \sum_{n=1}^{\infty} \left(\frac{4}{7}\right)^n$$

are both convergent geometric series, it follows that

$$\sum_{n=1}^{\infty} \frac{2^n + 4^n}{7^n} = \sum_{n=1}^{\infty} \left(\frac{2}{7}\right)^n + \sum_{n=1}^{\infty} \left(\frac{4}{7}\right)^n$$

also converges.

$$44. \sum_{n=1}^{\infty} \frac{n^3}{n!}$$

SOLUTION The presence of the factorial suggests applying the Ratio Test. With $a_n = \frac{n^3}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^3}{(n+1)!} \cdot \frac{n!}{n^3} = \frac{(n+1)^2}{n^3} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0 < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n^3}{n!}$ converges by the Ratio Test.

$$45. \sum_{n=1}^{\infty} \frac{n^3}{5^n}$$

SOLUTION The presence of the exponential term suggests applying the Ratio Test. With $a_n = \frac{n^3}{5^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^3}{5^{n+1}} \cdot \frac{5^n}{n^3} = \frac{1}{5} \left(1 + \frac{1}{n} \right)^3 \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{5} \cdot 1^3 = \frac{1}{5} < 1.$$

Therefore, the series $\sum_{n=1}^{\infty} \frac{n^3}{5^n}$ converges by the Ratio Test.

$$46. \sum_{n=2}^{\infty} \frac{1}{n(\ln n)^3}$$

SOLUTION The general term in this series suggests applying the Integral Test. Let $f(x) = \frac{1}{x(\ln x)^3}$. This function is continuous, positive and decreasing for $x \geq 2$, so the Integral Test does apply. Now

$$\int_2^{\infty} \frac{dx}{x(\ln x)^3} = \lim_{R \rightarrow \infty} \int_2^R \frac{dx}{x(\ln x)^3} = \lim_{R \rightarrow \infty} \int_{\ln 2}^{\ln R} \frac{du}{u^3} = -\frac{1}{2} \lim_{R \rightarrow \infty} \left(\frac{1}{(\ln R)^2} - \frac{1}{(\ln 2)^2} \right) = \frac{1}{2(\ln 2)^2}.$$

The integral converges; hence, the series $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^3}$ also converges.

$$47. \sum_{n=2}^{\infty} \frac{1}{\sqrt{n^3 - n^2}}$$

SOLUTION This series is similar to a p -series; because

$$\frac{1}{\sqrt{n^3 - n^2}} \approx \frac{1}{\sqrt{n^3}} = \frac{1}{n^{3/2}}$$

for large n , we will apply the Limit Comparison Test comparing with the p -series with $p = \frac{3}{2}$. Now,

$$L = \lim_{n \rightarrow \infty} \frac{\frac{1}{\sqrt{n^3 - n^2}}}{\frac{1}{n^{3/2}}} = \lim_{n \rightarrow \infty} \sqrt{\frac{n^3}{n^3 - n^2}} = 1.$$

The p -series with $p = \frac{3}{2}$ converges and L exists; therefore, the series $\sum_{n=2}^{\infty} \frac{1}{\sqrt{n^3 - n^2}}$ also converges.

$$48. \sum_{n=1}^{\infty} \frac{n^2 + 4n}{3n^4 + 9}$$

SOLUTION This series is similar to a p -series; because

$$\frac{n^2 + 4n}{3n^4 + 9} \approx \frac{n^2}{\sqrt{3n^4}} = \frac{1}{3n^2}$$

for large n , we will apply the Limit Comparison Test comparing with the p -series with $p = 2$. Now,

$$L = \lim_{n \rightarrow \infty} \frac{\frac{n^2 + 4n}{3n^4 + 9}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{n^4 + 4n^3}{3n^4 + 9} = \frac{1}{3}.$$

The p -series with $p = 2$ converges and L exists; therefore, the series $\sum_{n=1}^{\infty} \frac{n^2 + 4n}{3n^4 + 9}$ also converges.

$$49. \sum_{n=1}^{\infty} n^{-0.8}$$

SOLUTION

$$\sum_{n=1}^{\infty} n^{-0.8} = \sum_{n=1}^{\infty} \frac{1}{n^{0.8}}$$

so that this is a divergent p -series.

$$50. \sum_{n=1}^{\infty} (0.8)^{-n} n^{-0.8}$$

SOLUTION

$$\sum_{n=1}^{\infty} (0.8)^{-n} n^{-0.8} = \sum_{n=1}^{\infty} (0.8^{-1})^n n^{-0.8} = \sum_{n=1}^{\infty} \frac{1.25^n}{n^{0.8}}$$

With $a_n = \frac{1.25^n}{n^{0.8}}$ we have

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{1.25^{n+1}}{(n+1)^{0.8}} \cdot \frac{n^{0.8}}{1.25^n} = 1.25 \left(\frac{n}{n+1} \right)^{0.8}$$

so that

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1.25 > 1$$

Thus the original series diverges, by the Ratio Test.

$$51. \sum_{n=1}^{\infty} 4^{-2n+1}$$

SOLUTION Observe

$$\sum_{n=1}^{\infty} 4^{-2n+1} = \sum_{n=1}^{\infty} 4 \cdot (4^{-2})^n = \sum_{n=1}^{\infty} 4 \left(\frac{1}{16} \right)^n$$

is a geometric series with $r = \frac{1}{16}$; therefore, this series converges.

$$52. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$$

SOLUTION This is an alternating series with $a_n = \frac{1}{\sqrt{n}}$. Because a_n forms a decreasing sequence which converges to

zero, the series $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$ converges by the Leibniz Test.

$$53. \sum_{n=1}^{\infty} \sin \frac{1}{n^2}$$

SOLUTION Here, we will apply the Limit Comparison Test, comparing with the p -series with $p = 2$. Now,

$$L = \lim_{n \rightarrow \infty} \frac{\sin \frac{1}{n^2}}{\frac{1}{n^2}} = \lim_{u \rightarrow 0} \frac{\sin u}{u} = 1,$$

where $u = \frac{1}{n^2}$. The p -series with $p = 2$ converges and L exists; therefore, the series $\sum_{n=1}^{\infty} \sin \frac{1}{n^2}$ also converges.

$$54. \sum_{n=1}^{\infty} (-1)^n \cos \frac{1}{n}$$

SOLUTION Because

$$\lim_{n \rightarrow \infty} \cos \frac{1}{n} = \cos 0 = 1 \neq 0,$$

the general term in the series $\sum_{n=1}^{\infty} (-1)^n \cos \frac{1}{n}$ does not tend toward zero; therefore, the series diverges by the Divergence Test.

$$55. \sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n}}$$

SOLUTION Because

$$\lim_{n \rightarrow \infty} \frac{2^n}{\sqrt{n}} = \lim_{x \rightarrow \infty} \frac{2^x}{\sqrt{x}} = \lim_{x \rightarrow \infty} \frac{2^x \ln 2}{\frac{1}{2\sqrt{x}}} = \lim_{x \rightarrow \infty} 2^{x+1} \sqrt{x} \ln 2 = \infty \neq 0,$$

the general term in the series $\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n}}$ does not tend toward zero; therefore, the series diverges by the Divergence Test.


$$56. \sum_{n=1}^{\infty} \left(\frac{n}{n+12} \right)^n$$

SOLUTION Because the general term has the form of a function of n raised to the n th power, we might be tempted to use the Root Test; however, the Root Test is inconclusive for this series. Instead, note

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \left(1 + \frac{12}{n} \right)^{-n} = \lim_{n \rightarrow \infty} \left[\left(1 + \frac{12}{n} \right)^{n/12} \right]^{-12} = e^{-12} \neq 0.$$

Therefore, the series diverges by the Divergence Test.

Further Insights and Challenges

57.  **Proof of the Root Test** Let $S = \sum_{n=0}^{\infty} a_n$ be a positive series, and assume that $L = \lim_{n \rightarrow \infty} \sqrt[n]{a_n}$ exists.

(a) Show that S converges if $L < 1$. *Hint:* Choose R with $L < R < 1$ and show that $a_n \leq R^n$ for n sufficiently large. Then compare with the geometric series $\sum R^n$.

(b) Show that S diverges if $L > 1$.

SOLUTION Suppose $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = L$ exists.

(a) If $L < 1$, let $\epsilon = \frac{1-L}{2}$. By the definition of a limit, there is a positive integer N such that

$$-\epsilon \leq \sqrt[n]{a_n} - L \leq \epsilon$$

for $n \geq N$. From this, we conclude that

$$0 \leq \sqrt[n]{a_n} \leq L + \epsilon$$

for $n \geq N$. Now, let $R = L + \epsilon$. Then

$$R = L + \frac{1-L}{2} = \frac{L+1}{2} < \frac{1+1}{2} = 1,$$

and

$$0 \leq \sqrt[n]{a_n} \leq R \quad \text{or} \quad 0 \leq a_n \leq R^n$$

for $n \geq N$. Because $0 \leq R < 1$, the series $\sum_{n=N}^{\infty} R^n$ is a convergent geometric series, so the series $\sum_{n=N}^{\infty} a_n$ converges by

the Comparison Test. Therefore, the series $\sum_{n=0}^{\infty} a_n$ also converges.

(b) If $L > 1$, let $\epsilon = \frac{L-1}{2}$. By the definition of a limit, there is a positive integer N such that

$$-\epsilon \leq \sqrt[n]{a_n} - L \leq \epsilon$$

for $n \geq N$. From this, we conclude that

$$L - \epsilon \leq \sqrt[n]{a_n}$$

for $n \geq N$. Now, let $R = L - \epsilon$. Then

$$R = L - \frac{L-1}{2} = \frac{L+1}{2} > \frac{1+1}{2} = 1,$$

and

$$R \leq \sqrt[n]{a_n} \quad \text{or} \quad R^n \leq a_n$$

for $n \geq N$. Because $R > 1$, the series $\sum_{n=N}^{\infty} R^n$ is a divergent geometric series, so the series $\sum_{n=N}^{\infty} a_n$ diverges by the

Comparison Test. Therefore, the series $\sum_{n=0}^{\infty} a_n$ also diverges.

58. Show that the Ratio Test does not apply, but verify convergence using the Comparison Test for the series

$$\frac{1}{2} + \frac{1}{3^2} + \frac{1}{2^3} + \frac{1}{3^4} + \frac{1}{2^5} + \cdots$$

SOLUTION The general term of the series is:

$$a_n = \begin{cases} \frac{1}{2^n} & n \text{ odd} \\ \frac{1}{3^n} & n \text{ even} \end{cases}$$

First use the Ratio Test. If n is even,

$$\frac{a_{n+1}}{a_n} = \frac{\frac{1}{2^{n+1}}}{\frac{1}{3^n}} = \frac{3^n}{2^{n+1}} = \frac{1}{2} \cdot \left(\frac{3}{2}\right)^n$$

whereas, if n is odd,

$$\frac{a_{n+1}}{a_n} = \frac{\frac{1}{3^{n+1}}}{\frac{1}{2^n}} = \frac{2^n}{3^{n+1}} = \frac{1}{3} \cdot \left(\frac{2}{3}\right)^n$$

Since $\lim_{n \rightarrow \infty} \frac{1}{3} \cdot \left(\frac{2}{3}\right)^n = 0$ and $\lim_{n \rightarrow \infty} \frac{1}{2} \cdot \left(\frac{3}{2}\right)^n = \infty$, the sequence $\frac{a_{n+1}}{a_n}$ does not converge, and the Ratio Test is inconclusive.

However, we have $0 \leq a_n \leq \frac{1}{2^n}$ for all n , so the series converges by comparison with the convergent geometric series

$$\sum_{n=1}^{\infty} \frac{1}{2^n}$$

59. Let $S = \sum_{n=1}^{\infty} \frac{c^n n!}{n^n}$, where c is a constant.

(a) Prove that S converges absolutely if $|c| < e$ and diverges if $|c| > e$.

(b) It is known that $\lim_{n \rightarrow \infty} \frac{e^n n!}{n^{n+1/2}} = \sqrt{2\pi}$. Verify this numerically.

(c) Use the Limit Comparison Test to prove that S diverges for $c = e$.

SOLUTION

(a) With $a_n = \frac{c^n n!}{n^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|c|^{n+1} (n+1)!}{(n+1)^{n+1}} \cdot \frac{n^n}{|c|^n n!} = |c| \left(\frac{n}{n+1} \right)^n = |c| \left(1 + \frac{1}{n} \right)^{-n},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |c|e^{-1}.$$

Thus, by the Ratio Test, the series $\sum_{n=1}^{\infty} \frac{c^n n!}{n^n}$ converges when $|c|e^{-1} < 1$, or when $|c| < e$. The series diverges when $|c| > e$.

(b) The table below lists the value of $\frac{e^n n!}{n^{n+1/2}}$ for several increasing values of n . Since $\sqrt{2\pi} = 2.506628275$, the numerical evidence verifies that

$$\lim_{n \rightarrow \infty} \frac{e^n n!}{n^{n+1/2}} = \sqrt{2\pi}.$$

n	100	1000	10000	100000
$\frac{e^n n!}{n^{n+1/2}}$	2.508717995	2.506837169	2.506649163	2.506630363

(c) With $c = e$, the series S becomes $\sum_{n=1}^{\infty} \frac{e^n n!}{n^n}$. Using the result from part (b),

$$L = \lim_{n \rightarrow \infty} \frac{\frac{e^n n!}{n^n}}{\sqrt{n}} = \lim_{n \rightarrow \infty} \frac{e^n n!}{n^{n+1/2}} = \sqrt{2\pi}.$$

Because the series $\sum_{n=1}^{\infty} \sqrt{n}$ diverges by the Divergence Test and $L > 0$, we conclude that $\sum_{n=1}^{\infty} \frac{e^n n!}{n^n}$ diverges by the Limit Comparison Test.

10.6 Power Series

Preliminary Questions

1. Suppose that $\sum a_n x^n$ converges for $x = 5$. Must it also converge for $x = 4$? What about $x = -3$?

SOLUTION The power series $\sum a_n x^n$ is centered at $x = 0$. Because the series converges for $x = 5$, the radius of convergence must be at least 5 and the series converges absolutely at least for the interval $|x| < 5$. Both $x = 4$ and $x = -3$ are inside this interval, so the series converges for $x = 4$ and for $x = -3$.

2. Suppose that $\sum a_n (x - 6)^n$ converges for $x = 10$. At which of the points (a)–(d) must it also converge?

(a) $x = 8$ (b) $x = 11$ (c) $x = 3$ (d) $x = 0$

SOLUTION The given power series is centered at $x = 6$. Because the series converges for $x = 10$, the radius of convergence must be at least $|10 - 6| = 4$ and the series converges absolutely at least for the interval $|x - 6| < 4$, or $2 < x < 10$.

- (a) $x = 8$ is inside the interval $2 < x < 10$, so the series converges for $x = 8$.
 (b) $x = 11$ is not inside the interval $2 < x < 10$, so the series may or may not converge for $x = 11$.
 (c) $x = 3$ is inside the interval $2 < x < 10$, so the series converges for $x = 3$.
 (d) $x = 0$ is not inside the interval $2 < x < 10$, so the series may or may not converge for $x = 0$.

3. What is the radius of convergence of $F(3x)$ if $F(x)$ is a power series with radius of convergence $R = 12$?

SOLUTION If the power series $F(x)$ has radius of convergence $R = 12$, then the power series $F(3x)$ has radius of convergence $R = \frac{12}{3} = 4$.

4. The power series $F(x) = \sum_{n=1}^{\infty} n x^n$ has radius of convergence $R = 1$. What is the power series expansion of $F'(x)$ and what is its radius of convergence?

SOLUTION We obtain the power series expansion for $F'(x)$ by differentiating the power series expansion for $F(x)$ term-by-term. Thus,

$$F'(x) = \sum_{n=1}^{\infty} n^2 x^{n-1}.$$

The radius of convergence for this series is $R = 1$, the same as the radius of convergence for the series expansion for $F(x)$.

Exercises

1. Use the Ratio Test to determine the radius of convergence R of $\sum_{n=0}^{\infty} \frac{x^n}{2^n}$. Does it converge at the endpoints $x = \pm R$?

SOLUTION With $a_n = \frac{x^n}{2^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|^{n+1}}{2^{n+1}} \cdot \frac{2^n}{|x|^n} = \frac{|x|}{2} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|}{2}.$$

By the Ratio Test, the series converges when $\rho = \frac{|x|}{2} < 1$, or $|x| < 2$, and diverges when $\rho = \frac{|x|}{2} > 1$, or $|x| > 2$. The radius of convergence is therefore $R = 2$. For $x = -2$, the left endpoint, the series becomes $\sum_{n=0}^{\infty} (-1)^n$, which is divergent. For $x = 2$, the right endpoint, the series becomes $\sum_{n=0}^{\infty} 1$, which is also divergent. Thus the series diverges at both endpoints.

2. Use the Ratio Test to show that $\sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}2^n}$ has radius of convergence $R = 2$. Then determine whether it converges at the endpoints $R = \pm 2$.

SOLUTION With $a_n = \frac{x^n}{\sqrt{n}2^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|^{n+1}}{\sqrt{n+1} \cdot 2^{n+1}} \cdot \frac{\sqrt{n} \cdot 2^n}{|x|^n} = \frac{|x|}{2} \cdot \sqrt{\frac{n}{n+1}} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|}{2} \cdot 1 = \frac{|x|}{2}.$$

By the Ratio Test, the series converges when $\rho = \frac{|x|}{2} < 1$, or $|x| < 2$, and diverges when $\rho = \frac{|x|}{2} > 1$, or $|x| > 2$. The radius of convergence is therefore $R = 2$.

For the endpoint $x = 2$, the series becomes

$$\sum_{n=1}^{\infty} \frac{2^n}{\sqrt{n} \cdot 2^n} = \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}},$$

which is a divergent p -series. For the endpoint $x = -2$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt{n} \cdot 2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}.$$

This alternating series converges by the Leibniz Test, but its associated positive series is a divergent p -series. Thus, the series for $x = -2$ is conditionally convergent.

3. Show that the power series (a)–(c) have the same radius of convergence. Then show that (a) diverges at both endpoints, (b) converges at one endpoint but diverges at the other, and (c) converges at both endpoints.

$$\text{(a)} \sum_{n=1}^{\infty} \frac{x^n}{3^n} \qquad \text{(b)} \sum_{n=1}^{\infty} \frac{x^n}{n3^n} \qquad \text{(c)} \sum_{n=1}^{\infty} \frac{x^n}{n^2 3^n}$$

SOLUTION

(a) With $a_n = \frac{x^n}{3^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{3^{n+1}} \cdot \frac{3^n}{x^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x}{3} \right| = \left| \frac{x}{3} \right|$$

Then $\rho < 1$ if $|x| < 3$, so that the radius of convergence is $R = 3$. For the endpoint $x = 3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{3^n}{3^n} = \sum_{n=1}^{\infty} 1,$$

which diverges by the Divergence Test. For the endpoint $x = -3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-3)^n}{3^n} = \sum_{n=1}^{\infty} (-1)^n,$$

which also diverges by the Divergence Test.

(b) With $a_n = \frac{x^n}{n3^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)3^{n+1}} \cdot \frac{n3^n}{x^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x}{3} \left(\frac{n}{n+1} \right) \right| = \left| \frac{x}{3} \right|.$$

Then $\rho < 1$ when $|x| < 3$, so that the radius of convergence is $R = 3$. For the endpoint $x = 3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{3^n}{n3^n} = \sum_{n=1}^{\infty} \frac{1}{n},$$

which is the divergent harmonic series. For the endpoint $x = -3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-3)^n}{n3^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n},$$

which converges by the Leibniz Test.

(c) With $a_n = \frac{x^n}{n^2 3^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)^2 3^{n+1}} \cdot \frac{n^2 3^n}{x^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x}{3} \left(\frac{n}{n+1} \right)^2 \right| = \left| \frac{x}{3} \right|$$

Then $\rho < 1$ when $|x| < 3$, so that the radius of convergence is $R = 3$. For the endpoint $x = 3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{3^n}{n^2 3^n} = \sum_{n=1}^{\infty} \frac{1}{n^2},$$

which is a convergent p -series. For the endpoint $x = -3$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-3)^n}{n^2 3^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2},$$

which converges by the Leibniz Test.

4. Repeat Exercise 3 for the following series:

(a) $\sum_{n=1}^{\infty} \frac{(x-5)^n}{9^n}$

(b) $\sum_{n=1}^{\infty} \frac{(x-5)^n}{n 9^n}$

(c) $\sum_{n=1}^{\infty} \frac{(x-5)^n}{n^2 9^n}$

SOLUTION

(a) With $a_n = \frac{(x-5)^n}{9^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x-5)^{n+1}}{9^{n+1}} \cdot \frac{9^n}{(x-5)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x-5}{9} \right| = \left| \frac{x-5}{9} \right|$$

Then $\rho < 1$ when $|x-5| < 9$, so that the radius of convergence is $R = 9$. Because the series is centered at $x = 5$, the series converges absolutely on the interval $|x-5| < 9$, or $-4 < x < 14$. For the endpoint $x = 14$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(14-5)^n}{9^n} = \sum_{n=1}^{\infty} 1,$$

which diverges by the Divergence Test. For the endpoint $x = -4$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-4-5)^n}{9^n} = \sum_{n=1}^{\infty} (-1)^n,$$

which also diverges by the Divergence Test.

(b) With $a_n = \frac{(x-5)^n}{n 9^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x-5)^{n+1}}{(n+1) 9^{n+1}} \cdot \frac{n 9^n}{(x-5)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x-5}{9} \cdot \frac{n}{n+1} \right| = \left| \frac{x-5}{9} \right|.$$

Then $\rho < 1$ when $|x-5| < 9$, so that the radius of convergence is $R = 9$. Because the series is centered at $x = 5$, the series converges absolutely on the interval $|x-5| < 9$, or $-4 < x < 14$. For the endpoint $x = 14$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(14-5)^n}{n 9^n} = \sum_{n=1}^{\infty} \frac{1}{n},$$

which is the divergent harmonic series. For the endpoint $x = -4$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-4-5)^n}{n 9^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n},$$

which converges by the Leibniz Test.

(c) With $a_n = \frac{(x-5)^n}{n^2 9^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x-5)^{n+1}}{(n+1)^2 9^{n+1}} \cdot \frac{n^2 9^n}{(x-5)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x-5}{9} \left(\frac{n}{n+1} \right)^2 \right| = \left| \frac{x-5}{9} \right|.$$

Then $\rho < 1$ when $|x - 5| < 9$, so that the radius of convergence is $R = 9$. Because the series is centered at $x = 5$, the series converges absolutely on the interval $|x - 5| < 9$, or $-4 < x < 14$. For the endpoint $x = 14$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(14-5)^n}{n^2 9^n} = \sum_{n=1}^{\infty} \frac{1}{n^2},$$

which is a convergent p -series. For the endpoint $x = -4$, the series becomes

$$\sum_{n=1}^{\infty} \frac{(-4-5)^n}{n^2 9^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2},$$

which converges by the Leibniz Test.

5. Show that $\sum_{n=0}^{\infty} n^n x^n$ diverges for all $x \neq 0$.

SOLUTION With $a_n = n^n x^n$, and assuming $x \neq 0$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)^{n+1} x^{n+1}}{n^n x^n} \right| = \lim_{n \rightarrow \infty} \left| x \left(1 + \frac{1}{n}\right)^n (n+1) \right| = \infty$$

$\rho < 1$ only if $x = 0$, so that the radius of convergence is therefore $R = 0$. In other words, the power series converges only for $x = 0$.

6. For which values of x does $\sum_{n=0}^{\infty} n! x^n$ converge?

SOLUTION With $a_n = n! x^n$, and assuming $x \neq 0$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)! x^{n+1}}{n! x^n} \right| = \lim_{n \rightarrow \infty} |(n+1)x| = \infty$$

$\rho < 1$ only if $x = 0$, so that the radius of convergence is $R = 0$. In other words, the power series converges only for $x = 0$.

7. Use the Ratio Test to show that $\sum_{n=0}^{\infty} \frac{x^{2n}}{3^n}$ has radius of convergence $R = \sqrt{3}$.

SOLUTION With $a_n = \frac{x^{2n}}{3^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{2(n+1)}}{3^{n+1}} \cdot \frac{3^n}{x^{2n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^2}{3} \right| = \left| \frac{x^2}{3} \right|$$

Then $\rho < 1$ when $|x^2| < 3$, or $x = \sqrt{3}$, so the radius of convergence is $R = \sqrt{3}$.

8. Show that $\sum_{n=0}^{\infty} \frac{x^{3n+1}}{64^n}$ has radius of convergence $R = 4$.

SOLUTION With $a_n = \frac{x^{3n+1}}{64^n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{3(n+1)+1}}{64^{n+1}} \cdot \frac{64^n}{x^{3n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^3}{64} \right| = \left| \frac{x^3}{64} \right|$$

Then $\rho < 1$ when $|x|^3 < 64$ or $|x| = 4$, so the radius of convergence is $R = 4$.

In Exercises 9–34, find the interval of convergence.

9. $\sum_{n=0}^{\infty} n x^n$

SOLUTION With $a_n = n x^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)x^{n+1}}{n x^n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{n+1}{n} \right| = |x|$$

Then $\rho < 1$ when $|x| < 1$, so that the radius of convergence is $R = 1$, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=0}^{\infty} n$, which diverges by the Divergence Test.

For the endpoint $x = -1$, the series becomes $\sum_{n=1}^{\infty} (-1)^n n$, which also diverges by the Divergence Test. Thus, the series $\sum_{n=0}^{\infty} nx^n$ converges for $-1 < x < 1$ and diverges elsewhere.

$$10. \sum_{n=1}^{\infty} \frac{2^n}{n} x^n$$

SOLUTION With $a_n = \frac{2^n}{n} x^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{2^{n+1} x^{n+1}}{n+1} \cdot \frac{n}{2^n x^n} \right| = \lim_{n \rightarrow \infty} \left| 2x \frac{n}{n+1} \right| = |2x|$$

$\rho < 1$ when $|x| < \frac{1}{2}$, so the radius of convergence is $R = \frac{1}{2}$, and the series converges absolutely on the interval $|x| < \frac{1}{2}$, or $-\frac{1}{2} < x < \frac{1}{2}$. For the endpoint $x = \frac{1}{2}$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{n}$, which is the divergent harmonic series. For the endpoint $x = -\frac{1}{2}$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=1}^{\infty} \frac{x^n}{n} x^n$ converges for $-\frac{1}{2} \leq x < \frac{1}{2}$ and diverges elsewhere.

$$11. \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n+1}}{2^n n}$$

SOLUTION With $a_n = (-1)^n \frac{x^{2n+1}}{2^n n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{x^{2(n+1)+1}}{2^{n+1}(n+1)} \cdot \frac{2^n n}{x^{2n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^2}{2} \cdot \frac{n}{n+1} \right| = \left| \frac{x^2}{2} \right|$$

Then $\rho < 1$ when $|x| < \sqrt{2}$, so the radius of convergence is $R = \sqrt{2}$, and the series converges absolutely on the interval $-\sqrt{2} < x < \sqrt{2}$. For the endpoint $x = -\sqrt{2}$, the series becomes $\sum_{n=1}^{\infty} (-1)^n \frac{-\sqrt{2}}{n} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\sqrt{2}}{n}$, which converges by the Leibniz test. For the endpoint $x = \sqrt{2}$, the series becomes $\sum_{n=1}^{\infty} (-1)^n \frac{\sqrt{2}}{n}$ which also converges by the Leibniz test.

Thus the series $\sum_{n=1}^{\infty} (-1)^n \frac{x^{2n+1}}{2^n n}$ converges for $-\sqrt{2} \leq x \leq \sqrt{2}$ and diverges elsewhere.

$$12. \sum_{n=0}^{\infty} (-1)^n \frac{n}{4^n} x^{2n}$$

SOLUTION With $a_n = (-1)^n \frac{n}{4^n} x^{2n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)x^{2(n+1)}}{4^{n+1}} \cdot \frac{4^n}{nx^{2n}} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^2}{4} \cdot \frac{n+1}{n} \right| = \left| \frac{x^2}{4} \right|$$

Then $\rho < 1$ when $|x^2| < 4$, or $|x| < 2$, so the radius of convergence is $R = 2$, and the series converges absolutely for $-2 < x < 2$. At both endpoints $x = \pm 2$, the series becomes $\sum_{n=0}^{\infty} (-1)^n n$, which diverges by the Divergence Test. Thus,

the series $\sum_{n=0}^{\infty} (-1)^n \frac{n}{4^n} x^{2n}$ converges for $-2 < x < 2$ and diverges elsewhere.

$$13. \sum_{n=4}^{\infty} \frac{x^n}{n^5}$$

SOLUTION With $a_n = \frac{x^n}{n^5}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)^5} \cdot \frac{n^5}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \left(\frac{n}{n+1} \right)^5 \right| = |x|$$

Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is $R = 1$, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{n^5}$, which is a convergent p -series. For the

endpoint $x = -1$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^5}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=4}^{\infty} \frac{x^n}{n^5}$ converges for $-1 \leq x \leq 1$ and diverges elsewhere.

$$14. \sum_{n=8}^{\infty} n^7 x^n$$

SOLUTION With $a_n = n^7 x^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)^7 x^{n+1}}{n^7 x^n} \right| = \lim_{n \rightarrow \infty} \left| x \left(\frac{n+1}{n} \right)^7 \right| = |x|$$

Then $\rho < 1$ when $|x| < 1$, so that the radius of convergence is $R = 1$, and the series converges absolutely on the interval $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=8}^{\infty} n^7$, which diverges by the Divergence test; for the endpoints

$x = -1$, the series becomes $\sum_{n=8}^{\infty} (-1)^n n^7$, which also diverges by the Divergence test. Thus the series $\sum_{n=8}^{\infty} n^7 x^n$ converges for $-1 < x < 1$ and diverges elsewhere.

$$15. \sum_{n=0}^{\infty} \frac{x^n}{(n!)^2}$$

SOLUTION With $a_n = \frac{x^n}{(n!)^2}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{((n+1)!)^2} \cdot \frac{(n!)^2}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \left(\frac{1}{n+1} \right)^2 \right| = 0$$

$\rho < 1$ for all x , so the radius of convergence is $R = \infty$, and the series converges absolutely for all x .

$$16. \sum_{n=0}^{\infty} \frac{8^n}{n!} x^n$$

SOLUTION With $a_n = \frac{8^n x^n}{n!}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{8^{n+1} x^{n+1}}{(n+1)!} \cdot \frac{n!}{8^n x^n} \right| = \lim_{n \rightarrow \infty} \left| 8x \cdot \frac{1}{n+1} \right| = 0$$

$\rho < 1$ for all x , so the radius of convergence is $R = \infty$, and the series converges absolutely for all x .

$$17. \sum_{n=0}^{\infty} \frac{(2n)!}{(n!)^3} x^n$$

SOLUTION With $a_n = \frac{(2n)! x^n}{(n!)^3}$, and assuming $x \neq 0$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(2(n+1))! x^{n+1}}{((n+1)!)^3} \cdot \frac{(n!)^3}{(2n)! x^n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{(2n+2)(2n+1)}{(n+1)^3} \right| \\ &= \lim_{n \rightarrow \infty} \left| x \frac{4n^2 + 6n + 2}{n^3 + 3n^2 + 3n + 1} \right| = \lim_{n \rightarrow \infty} \left| x \frac{4n^{-1} + 6n^{-2} + 2n^{-3}}{1 + 3n^{-1} + 3n^{-2} + n^{-3}} \right| = 0 \end{aligned}$$

Then $\rho < 1$ for all x , so the radius of convergence is $R = \infty$, and the series converges absolutely for all x .

$$18. \sum_{n=0}^{\infty} \frac{4^n}{(2n+1)!} x^{2n-1}$$

SOLUTION With $a_n = \frac{4^n x^{2n-1}}{(2n+1)!}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{4^{n+1} x^{2n+1}}{(2n+3)!} \cdot \frac{(2n+1)!}{4^n x^{2n-1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{4x^2}{(2n+3)(2n+2)} \right| = 0$$

Then ρ is always less than 1, and the series converges absolutely for all x .

$$19. \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{\sqrt{n^2+1}}$$

SOLUTION With $a_n = \frac{(-1)^n x^n}{\sqrt{n^2+1}}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} x^{n+1}}{\sqrt{n^2+2n+2}} \cdot \frac{\sqrt{n^2+1}}{(-1)^n x^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| x \frac{\sqrt{n^2+1}}{\sqrt{n^2+2n+2}} \right| = \lim_{n \rightarrow \infty} \left| x \sqrt{\frac{n^2+1}{n^2+2n+2}} \right| = \lim_{n \rightarrow \infty} \left| x \sqrt{\frac{1+1/n^2}{1+2/n+2/n^2}} \right| \\ &= |x| \end{aligned}$$

Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is $R = 1$, and the series converges absolutely on the interval $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^2+1}}$, which converges by the Leibniz Test. For the

endpoint $x = -1$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n^2+1}}$, which diverges by the Limit Comparison Test comparing with the divergent harmonic series. Thus, the series $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{\sqrt{n^2+1}}$ converges for $-1 < x \leq 1$ and diverges elsewhere.

$$20. \sum_{n=0}^{\infty} \frac{x^n}{n^4+2}$$

SOLUTION With $a_n = \frac{x^n}{n^4+2}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)^4+2} \cdot \frac{n^4+2}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{n^4+2}{n^4+4n^3+6n^2+4n+3} \right| = |x|$$

$\rho < 1$ when $|x| < 1$, so the radius of convergence is $R = 1$, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{n^4+2}$. Because $\frac{1}{n^4+2} < \frac{1}{n^4}$ and the series $\sum_{n=0}^{\infty} \frac{1}{n^4}$ is a convergent p -series, the endpoint series converges by the Comparison Test. For the endpoint $x = -1$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^4+2}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=0}^{\infty} \frac{x^n}{n^4+2}$ converges for $-1 \leq x \leq 1$ and diverges elsewhere.

$$21. \sum_{n=15}^{\infty} \frac{x^{2n+1}}{3n+1}$$

SOLUTION With $a_n = \frac{x^{2n+1}}{3n+1}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{2n+3}}{3n+4} \cdot \frac{3n+1}{x^{2n+1}} \right| = \lim_{n \rightarrow \infty} \left| x^2 \frac{3n+1}{3n+4} \right| = |x^2|$$

Then $\rho < 1$ when $|x^2| < 1$, so the radius of convergence is $R = 1$, and the series converges absolutely for $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=15}^{\infty} \frac{1}{3n+1}$, which diverges by the Limit Comparison Test comparing

with the divergent harmonic series. For the endpoint $x = -1$, the series becomes $\sum_{n=15}^{\infty} \frac{-1}{3n+1}$, which also diverges by the Limit Comparison Test comparing with the divergent harmonic series. Thus, the series $\sum_{n=15}^{\infty} \frac{x^{2n+1}}{3n+1}$ converges for $-1 < x < 1$ and diverges elsewhere.

$$22. \sum_{n=1}^{\infty} \frac{x^n}{n-4 \ln n}$$

SOLUTION With $a_n = \frac{x^n}{n-4 \ln n}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)-4 \ln(n+1)} \cdot \frac{n-4 \ln n}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{n-4 \ln n}{(n+1)-4 \ln(n+1)} \right| \\ &= \lim_{n \rightarrow \infty} \left| x \frac{1-4(\ln n)/n}{1+n^{-1}-4(\ln(n+1))/n} \right| = |x| \end{aligned}$$

Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{n-4 \ln n}$. Because $\frac{1}{n-4 \ln n} > \frac{1}{n}$ and $\sum_{n=1}^{\infty} \frac{1}{n}$ is the divergent harmonic series, the endpoint series diverges by the Comparison Test. For the endpoint $x = -1$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n-4 \ln n}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=1}^{\infty} \frac{x^n}{n-4 \ln n}$ converges for $-1 \leq x < 1$ and diverges elsewhere.

$$23. \sum_{n=2}^{\infty} \frac{x^n}{\ln n}$$

SOLUTION With $a_n = \frac{x^n}{\ln n}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{\ln(n+1)} \cdot \frac{\ln n}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{\ln(n+1)}{\ln n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{1/(n+1)}{1/n} \right| = \lim_{n \rightarrow \infty} \left| x \frac{n}{n+1} \right| = |x|$$

using L'Hôpital's rule. Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=2}^{\infty} \frac{1}{\ln n}$. Because $\frac{1}{\ln n} > \frac{1}{n}$ and

$\sum_{n=2}^{\infty} \frac{1}{n}$ is the divergent harmonic series, the endpoint series diverges by the Comparison Test. For the endpoint $x = -1$,

the series becomes $\sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=2}^{\infty} \frac{x^n}{\ln n}$ converges for $-1 \leq x < 1$ and diverges elsewhere.

$$24. \sum_{n=2}^{\infty} \frac{x^{3n+2}}{\ln n}$$

SOLUTION With $a_n = \frac{x^{3n+2}}{\ln n}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{3n+5}}{\ln(n+1)} \cdot \frac{\ln n}{x^{3n+2}} \right| = \lim_{n \rightarrow \infty} \left| x^3 \cdot \frac{\ln(n+1)}{\ln n} \right| = \lim_{n \rightarrow \infty} \left| x^3 \cdot \frac{1/(n+1)}{1/n} \right| \\ &= \lim_{n \rightarrow \infty} \left| x^3 \cdot \frac{n}{n+1} \right| = |x^3| \end{aligned}$$

using L'Hôpital's rule. Thus $\rho < 1$ when $|x^3| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=2}^{\infty} \frac{1}{\ln n}$. Because $\frac{1}{\ln n} > \frac{1}{n}$ and

$\sum_{n=2}^{\infty} \frac{1}{n}$ is the divergent harmonic series, the endpoint series diverges by the Comparison Test. For the endpoint $x = -1$,

the series becomes $\sum_{n=2}^{\infty} \frac{(-1)^{3n+2}}{\ln n} = \sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=2}^{\infty} \frac{x^{3n+2}}{\ln n}$ converges for $-1 \leq x < 1$ and diverges elsewhere.

$$25. \sum_{n=1}^{\infty} n(x-3)^n$$

SOLUTION With $a_n = n(x-3)^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)(x-3)^{n+1}}{n(x-3)^n} \right| = \lim_{n \rightarrow \infty} \left| (x-3) \cdot \frac{n+1}{n} \right| = |x-3|$$

Then $\rho < 1$ when $|x-3| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x-3| < 1$, or $2 < x < 4$. For the endpoint $x = 4$, the series becomes $\sum_{n=1}^{\infty} n$, which diverges by the Divergence Test.

For the endpoint $x = 2$, the series becomes $\sum_{n=1}^{\infty} (-1)^n n$, which also diverges by the Divergence Test. Thus, the series

$\sum_{n=1}^{\infty} n(x-3)^n$ converges for $2 < x < 4$ and diverges elsewhere.

$$26. \sum_{n=1}^{\infty} \frac{(-5)^n (x-3)^n}{n^2}$$

SOLUTION With $a_n = \frac{(-5)^n (x-3)^n}{n^2}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-5)^{n+1} (x-3)^{n+1}}{(n+1)^2} \cdot \frac{n^2}{(-5)^n (x-3)^n} \right| = \lim_{n \rightarrow \infty} \left| 5(x-3) \cdot \frac{n^2}{n^2 + 2n + 1} \right| \\ &= \lim_{n \rightarrow \infty} \left| 5(x-3) \cdot \frac{1}{1 + 2n^{-1} + n^{-2}} \right| = |5(x-3)| \end{aligned}$$

Then $\rho < 1$ when $|5(x-3)| < 1$, or $|x-3| < \frac{1}{5}$. Thus the radius of convergence is $\frac{1}{5}$, and the series converges absolutely on the interval $|x-3| < \frac{1}{5}$, or $\frac{14}{5} < x < \frac{16}{5}$. For the endpoint $x = \frac{16}{5}$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$, which converges

by the Leibniz Test. For the endpoint $x = \frac{14}{5}$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{n^2}$, which is a convergent p -series. Thus, the series

$\sum_{n=1}^{\infty} \frac{(-5)^n (x-3)^n}{n^2}$ converges for $\frac{14}{5} \leq x \leq \frac{16}{5}$ and diverges elsewhere.

$$27. \sum_{n=1}^{\infty} (-1)^n n^5 (x-7)^n$$

SOLUTION With $a_n = (-1)^n n^5 (x-7)^n$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} (n+1)^5 (x-7)^{n+1}}{(-1)^n n^5 (x-7)^n} \right| = \lim_{n \rightarrow \infty} \left| (x-7) \cdot \frac{(n+1)^5}{n^5} \right| \\ &= \lim_{n \rightarrow \infty} \left| (x-7) \cdot \frac{n^5 + \dots}{n^5} \right| = |x-7| \end{aligned}$$

Then $\rho < 1$ when $|x-7| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x-7| < 1$, or $6 < x < 8$. For the endpoint $x = 6$, the series becomes $\sum_{n=1}^{\infty} (-1)^{2n} n^5 = \sum_{n=1}^{\infty} n^5$, which diverges by the

Divergence Test. For the endpoint $x = 8$, the series becomes $\sum_{n=1}^{\infty} (-1)^n n^5$, which also diverges by the Divergence Test.

Thus, the series $\sum_{n=1}^{\infty} (-1)^n n^5 (x-7)^n$ converges for $6 < x < 8$ and diverges elsewhere.

$$28. \sum_{n=0}^{\infty} 27^n (x-1)^{3n+2}$$

SOLUTION With $a_n = 27^n (x-1)^{3n+2}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{27^{n+1} (x-1)^{3n+5}}{27^n (x-1)^{3n+2}} \right| = \lim_{n \rightarrow \infty} |27(x-1)^3| = |27(x-1)^3|$$

Then $\rho < 1$ when $|27(x-1)^3| < 1$, or when $|(x-1)^3| < \frac{1}{27}$, so when $|x-1| < \frac{1}{3}$. Thus the radius of convergence is $\frac{1}{3}$, and the series converges absolutely when $\frac{2}{3} < x < \frac{4}{3}$. For the endpoint $x = \frac{2}{3}$, the series becomes $\sum_{n=0}^{\infty} 27^n \left(\frac{-1}{3}\right)^{3n+2} = \frac{1}{9} \sum_{n=0}^{\infty} (-1)^n$ which diverges by the Divergence test. For the endpoint $x = \frac{4}{3}$, the series becomes $\sum_{n=0}^{\infty} 27^n \left(\frac{1}{3}\right)^{3n+2} = \frac{1}{9} \sum_{n=0}^{\infty} 1$, which also diverges by the Divergence Test. Thus the series $\sum_{n=0}^{\infty} 27^n (x-1)^{3n+2}$ converges for $\frac{2}{3} < x < \frac{4}{3}$ and diverges elsewhere.

$$29. \sum_{n=1}^{\infty} \frac{2^n}{3n} (x+3)^n$$

SOLUTION With $a_n = \frac{2^n (x+3)^n}{3n}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{2^{n+1} (x+3)^{n+1}}{3(n+1)} \cdot \frac{3n}{2^n (x+3)^n} \right| = \lim_{n \rightarrow \infty} \left| 2(x+3) \cdot \frac{3n}{3n+3} \right| \\ &= \lim_{n \rightarrow \infty} \left| 2(x+3) \cdot \frac{1}{1+1/n} \right| = |2(x+3)| \end{aligned}$$

Then $\rho < 1$ when $|2(x+3)| < 1$, so when $|x+3| < \frac{1}{2}$. Thus the radius of convergence is $\frac{1}{2}$, and the series converges absolutely on the interval $|x+3| < \frac{1}{2}$, or $-\frac{7}{2} < x < -\frac{5}{2}$. For the endpoint $x = -\frac{5}{2}$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{3n}$, which diverges because it is a multiple of the divergent harmonic series. For the endpoint $x = -\frac{7}{2}$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{3n}$, which converges by the Leibniz Test. Thus, the series $\sum_{n=1}^{\infty} \frac{2^n}{3n} (x+3)^n$ converges for $-\frac{7}{2} \leq x < -\frac{5}{2}$ and diverges elsewhere.

$$30. \sum_{n=0}^{\infty} \frac{(x-4)^n}{n!}$$

SOLUTION With $a_n = \frac{(x-4)^n}{n!}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x-4)^{n+1}}{(n+1)!} \cdot \frac{n!}{(x-4)^n} \right| = \lim_{n \rightarrow \infty} \left| (x-4) \frac{1}{n} \right| = 0$$

Thus $\rho < 1$ for all x , so the radius of convergence is infinite, and $\sum_{n=0}^{\infty} \frac{(x-4)^n}{n!}$ converges for all x .

$$31. \sum_{n=0}^{\infty} \frac{(-5)^n}{n!} (x+10)^n$$

SOLUTION With $a_n = \frac{(-5)^n}{n!} (x+10)^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(-5)^{n+1} (x+10)^{n+1}}{(n+1)!} \cdot \frac{n!}{(-5)^n (x+10)^n} \right| = \lim_{n \rightarrow \infty} \left| 5(x+10) \frac{1}{n} \right| = 0$$

Thus $\rho < 1$ for all x , so the radius of convergence is infinite, and $\sum_{n=0}^{\infty} \frac{(-5)^n}{n!} (x+10)^n$ converges for all x .

$$32. \sum_{n=10}^{\infty} n!(x+5)^n$$

SOLUTION With $a_n = n!(x+5)^n$, and assuming $x+5 \neq 0$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)!(x+5)^{n+1}}{n!(x+5)^n} \right| = \lim_{n \rightarrow \infty} |(n+1)(x+5)| = \infty$$

Thus $\rho < 1$ only if $x+5 = 0$, so the radius of convergence is zero, and $\sum_{n=10}^{\infty} n!(x+5)^n$ converges only for $x = -5$.

$$33. \sum_{n=12}^{\infty} e^n(x-2)^n$$

SOLUTION With $a_n = e^n(x-2)^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{e^{n+1}(x-2)^{n+1}}{e^n(x-2)^n} \right| = \lim_{n \rightarrow \infty} |e(x-2)| = |e(x-2)|$$

Thus $\rho < 1$ when $|e(x-2)| < 1$, so when $|x-2| < e^{-1}$. Thus the radius of convergence is e^{-1} , and the series converges absolutely on the interval $|x-2| < e^{-1}$, or $2 - e^{-1} < x < 2 + e^{-1}$. For the endpoint $x = 2 + e^{-1}$, the series becomes $\sum_{n=1}^{\infty} 1$, which diverges by the Divergence Test. For the endpoint $x = 2 - e^{-1}$, the series becomes $\sum_{n=1}^{\infty} (-1)^n$, which also

diverges by the Divergence Test. Thus, the series $\sum_{n=12}^{\infty} e^n(x-2)^n$ converges for $2 - e^{-1} < x < 2 + e^{-1}$ and diverges elsewhere.

$$34. \sum_{n=2}^{\infty} \frac{(x+4)^n}{(n \ln n)^2}$$

SOLUTION With $a_n = \frac{(x+4)^n}{(n \ln n)^2}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(x+4)^{n+1}}{((n+1) \ln(n+1))^2} \cdot \frac{(n \ln n)^2}{(x+4)^n} \right| = \lim_{n \rightarrow \infty} \left| (x+4) \cdot \left(\frac{n}{n+1} \cdot \frac{\ln n}{\ln(n+1)} \right)^2 \right| = |x+4|$$

applying L'Hôpital's rule to evaluate the second term in the product. Thus $\rho < 1$ when $|x+4| < 1$, so the radius of convergence is 1, and the series converges absolutely on the interval $|x+4| < 1$, or $-5 < x < -3$. For the endpoint $x = -3$, the series becomes $\sum_{n=1}^{\infty} \frac{1}{(n \ln n)^2}$, which converges by the Limit Comparison Test comparing with the convergent

p -series $\sum_{n=2}^{\infty} \frac{1}{n^2}$. For the endpoint $x = -5$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{(n \ln n)^2}$, which converges by the Leibniz Test. Thus,

the series $\sum_{n=2}^{\infty} \frac{(x+4)^n}{(n \ln n)^2}$ converges for $-5 \leq x \leq -3$ and diverges elsewhere.

In Exercises 35–40, use Eq. (2) to expand the function in a power series with center $c = 0$ and determine the interval of convergence.

$$35. f(x) = \frac{1}{1-3x}$$

SOLUTION Substituting $3x$ for x in Eq. (2), we obtain

$$\frac{1}{1-3x} = \sum_{n=0}^{\infty} (3x)^n = \sum_{n=0}^{\infty} 3^n x^n.$$

This series is valid for $|3x| < 1$, or $|x| < \frac{1}{3}$.

$$36. f(x) = \frac{1}{1+3x}$$

SOLUTION Substituting $-3x$ for x in Eq. (2), we obtain

$$\frac{1}{1+3x} = \sum_{n=0}^{\infty} (-3x)^n = \sum_{n=0}^{\infty} (-3)^n x^n.$$

This series is valid for $|-3x| < 1$, or $|x| < \frac{1}{3}$.

$$37. f(x) = \frac{1}{3-x}$$

SOLUTION First write

$$\frac{1}{3-x} = \frac{1}{3} \cdot \frac{1}{1-\frac{x}{3}}.$$

Substituting $\frac{x}{3}$ for x in Eq. (2), we obtain

$$\frac{1}{1-\frac{x}{3}} = \sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n = \sum_{n=0}^{\infty} \frac{x^n}{3^n};$$

Thus,

$$\frac{1}{3-x} = \frac{1}{3} \sum_{n=0}^{\infty} \frac{x^n}{3^n} = \sum_{n=0}^{\infty} \frac{x^n}{3^{n+1}}.$$

This series is valid for $|x/3| < 1$, or $|x| < 3$.

$$38. f(x) = \frac{1}{4+3x}$$

SOLUTION First write

$$\frac{1}{4+3x} = \frac{1}{4} \cdot \frac{1}{1+\frac{3x}{4}}.$$

Substituting $-\frac{3x}{4}$ for x in Eq. (2), we obtain

$$\frac{1}{1+\frac{3x}{4}} = \sum_{n=0}^{\infty} \left(-\frac{3x}{4}\right)^n = \sum_{n=0}^{\infty} (-1)^n \frac{3^n x^n}{4^n};$$

Thus,

$$\frac{1}{4+3x} = \frac{1}{4} \sum_{n=0}^{\infty} (-1)^n \frac{3^n x^n}{4^n} = \sum_{n=0}^{\infty} (-1)^n \frac{3^n x^n}{4^{n+1}}.$$

This series is valid for $|-3x/4| < 1$, or $|x| < \frac{4}{3}$.

$$39. f(x) = \frac{1}{1+x^2}$$

SOLUTION Substituting $-x^2$ for x in Eq. (2), we obtain

$$\frac{1}{1+x^2} = \sum_{n=0}^{\infty} (-x^2)^n = \sum_{n=0}^{\infty} (-1)^n x^{2n}$$

This series is valid for $|x| < 1$.

$$40. f(x) = \frac{1}{16+2x^3}$$

SOLUTION First rewrite

$$\frac{1}{16+2x^3} = \frac{1}{16} \cdot \frac{1}{1+\frac{x^3}{8}}$$

Now substitute $-\frac{x^3}{8}$ for x in Eq. (2) to obtain

$$\frac{1}{1+\frac{x^3}{8}} = \sum_{n=0}^{\infty} \left(-\frac{x^3}{8}\right)^n = \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n}}{8^n}$$

Thus,

$$\frac{1}{16 + 2x^3} = \frac{1}{16} \cdot \frac{1}{1 + \frac{x^3}{8}} = \frac{1}{16} \cdot \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n}}{8}$$

This series is valid for $|x^3| < 8$, or $|x| < 2$.

41. Use the equalities

$$\frac{1}{1-x} = \frac{1}{-3-(x-4)} = \frac{-\frac{1}{3}}{1 + \left(\frac{x-4}{3}\right)}$$

to show that for $|x-4| < 3$,

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{(x-4)^n}{3^{n+1}}$$

SOLUTION Substituting $-\frac{x-4}{3}$ for x in Eq. (2), we obtain

$$\frac{1}{1 + \left(\frac{x-4}{3}\right)} = \sum_{n=0}^{\infty} \left(-\frac{x-4}{3}\right)^n = \sum_{n=0}^{\infty} (-1)^n \frac{(x-4)^n}{3^n}.$$

Thus,

$$\frac{1}{1-x} = -\frac{1}{3} \sum_{n=0}^{\infty} (-1)^n \frac{(x-4)^n}{3^n} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{(x-4)^n}{3^{n+1}}.$$

This series is valid for $|\frac{x-4}{3}| < 1$, or $|x-4| < 3$.

42. Use the method of Exercise 41 to expand $1/(1-x)$ in power series with centers $c = 2$ and $c = -2$. Determine the interval of convergence.

SOLUTION For $c = 2$, write

$$\frac{1}{1-x} = \frac{1}{-1-(x-2)} = -\frac{1}{1+(x-2)}.$$

Substituting $-(x-2)$ for x in Eq. (2), we obtain

$$\frac{1}{1+(x-2)} = \sum_{n=0}^{\infty} (-(x-2))^n = \sum_{n=0}^{\infty} (-1)^n (x-2)^n.$$

Thus,

$$\frac{1}{1-x} = -\sum_{n=0}^{\infty} (-1)^n (x-2)^n = \sum_{n=0}^{\infty} (-1)^{n+1} (x-2)^n.$$

This series is valid for $|(x-2)| < 1$, or $|x-2| < 1$.

For $c = -2$, write

$$\frac{1}{1-x} = \frac{1}{3-(x+2)} = \frac{1}{3} \cdot \frac{1}{1 - \frac{x+2}{3}}.$$

Substituting $\frac{x+2}{3}$ for x in Eq. (2), we obtain

$$\frac{1}{1 - \frac{x+2}{3}} = \sum_{n=0}^{\infty} \left(\frac{x+2}{3}\right)^n = \sum_{n=0}^{\infty} \frac{(x+2)^n}{3^n}.$$

Thus,

$$\frac{1}{1-x} = \frac{1}{3} \sum_{n=0}^{\infty} \frac{(x+2)^n}{3^n} = \sum_{n=0}^{\infty} \frac{(x+2)^n}{3^{n+1}}.$$

This series is valid for $|\frac{x+2}{3}| < 1$, or $|x+2| < 3$.

43. Use the method of Exercise 41 to expand $1/(4-x)$ in a power series with center $c = 5$. Determine the interval of convergence.

SOLUTION First write

$$\frac{1}{4-x} = \frac{1}{-1-(x-5)} = -\frac{1}{1+(x-5)}.$$

Substituting $-(x-5)$ for x in Eq. (2), we obtain

$$\frac{1}{1+(x-5)} = \sum_{n=0}^{\infty} (-(x-5))^n = \sum_{n=0}^{\infty} (-1)^n (x-5)^n.$$

Thus,

$$\frac{1}{4-x} = -\sum_{n=0}^{\infty} (-1)^n (x-5)^n = \sum_{n=0}^{\infty} (-1)^{n+1} (x-5)^n.$$

This series is valid for $|-(x-5)| < 1$, or $|x-5| < 1$.

44. Find a power series that converges only for x in $[2, 6)$.

SOLUTION The power series must be centered at $c = \frac{6+2}{2} = 4$, with radius of convergence $R = 2$. Consider the following series:

$$\sum_{n=1}^{\infty} \frac{(x-4)^n}{n2^n}.$$

With $a_n = \frac{1}{n2^n}$,

$$r = \lim_{n \rightarrow \infty} \frac{n2^n}{(n+1)2^{n+1}} = \frac{1}{2} \lim_{n \rightarrow \infty} \frac{n}{n+1} = \frac{1}{2}.$$

The radius of convergence is therefore $R = r^{-1} = 2$, and the series converges absolutely for $|x-4| < 2$, or $2 < x < 6$.

For the endpoint $x = 6$, the series becomes $\sum_{n=1}^{\infty} \frac{(6-4)^n}{n \cdot 2^n} = \sum_{n=1}^{\infty} \frac{1}{n}$, which is the divergent harmonic series. For the

endpoint $x = 2$, the series becomes $\sum_{n=1}^{\infty} \frac{(2-4)^n}{n \cdot 2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$, which converges by the Leibniz Test. Therefore, the series converges for $2 \leq x < 6$, as desired.

45. Apply integration to the expansion

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n = 1 - x + x^2 - x^3 + \dots$$

to prove that for $-1 < x < 1$,

$$\ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

SOLUTION To obtain the first expansion, substitute $-x$ for x in Eq. (2):

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-x)^n = \sum_{n=0}^{\infty} (-1)^n x^n.$$

This expansion is valid for $| -x | < 1$, or $-1 < x < 1$.

Upon integrating both sides of the above equation, we find

$$\ln(1+x) = \int \frac{dx}{1+x} = \int \left(\sum_{n=0}^{\infty} (-1)^n x^n \right) dx.$$

Integrating the series term-by-term then yields

$$\ln(1+x) = C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1}.$$

To determine the constant C , set $x = 0$. Then $0 = \ln(1+0) = C$. Finally,

$$\ln(1+x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}.$$

46. Use the result of Exercise 45 to prove that

$$\ln \frac{3}{2} = \frac{1}{2} - \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} - \frac{1}{4 \cdot 2^4} + \cdots$$

Use your knowledge of alternating series to find an N such that the partial sum S_N approximates $\ln \frac{3}{2}$ to within an error of at most 10^{-3} . Confirm using a calculator to compute both S_N and $\ln \frac{3}{2}$.

SOLUTION In the previous exercise we found that

$$\ln(1+x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{n+1}}{n+1}.$$

Setting $x = \frac{1}{2}$ yields:

$$\ln \frac{3}{2} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{\left(\frac{1}{2}\right)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n2^n} = \frac{1}{2} - \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} - \frac{1}{4 \cdot 2^4} + \cdots$$

Note that the series for $\ln \frac{3}{2}$ is an alternating series with $a_n = \frac{1}{n2^n}$. The error in approximating $\ln \frac{3}{2}$ by the partial sum S_N is therefore bounded by

$$\left| \ln \frac{3}{2} - S_N \right| < a_{N+1} = \frac{1}{(N+1)2^{N+1}}.$$

To obtain an error of at most 10^{-3} , we must find an N such that

$$\frac{1}{(N+1)2^{N+1}} < 10^{-3} \quad \text{or} \quad (N+1)2^{N+1} > 1000.$$

For $N = 6$, $(N+1)2^{N+1} = 7 \cdot 2^7 = 896 < 1000$, but for $N = 7$, $(N+1)2^{N+1} = 8 \cdot 2^8 = 2048 > 1000$; hence, the smallest value for N is $N = 7$. The corresponding approximation is

$$S_7 = \frac{1}{2} - \frac{1}{2 \cdot 2^2} + \frac{1}{3 \cdot 2^3} - \frac{1}{4 \cdot 2^4} + \frac{1}{5 \cdot 2^5} - \frac{1}{6 \cdot 2^6} + \frac{1}{7 \cdot 2^7} = 0.405803571.$$

Now, $\ln \frac{3}{2} = 0.405465108$, so

$$\left| \ln \frac{3}{2} - S_7 \right| = 3.385 \times 10^{-4} < 10^{-3}.$$

47. Let $F(x) = (x+1)\ln(1+x) - x$.

(a) Apply integration to the result of Exercise 45 to prove that for $-1 < x < 1$,

$$F(x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{n+1}}{n(n+1)}$$

(b) Evaluate at $x = \frac{1}{2}$ to prove

$$\frac{3}{2} \ln \frac{3}{2} - \frac{1}{2} = \frac{1}{1 \cdot 2 \cdot 2^2} - \frac{1}{2 \cdot 3 \cdot 2^3} + \frac{1}{3 \cdot 4 \cdot 2^4} - \frac{1}{4 \cdot 5 \cdot 2^5} + \cdots$$

(c) Use a calculator to verify that the partial sum S_4 approximates the left-hand side with an error no greater than the term a_5 of the series.

SOLUTION

(a) Note that

$$\int \ln(x+1) dx = (x+1)\ln(x+1) - x + C$$

Then integrating both sides of the result of Exercise 45 gives

$$(x+1)\ln(x+1) - x = \int \ln(x+1) dx = \int \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} dx$$

For $-1 < x < 1$, which is the interval of convergence of the series in Exercise 45, therefore, we can integrate term by term to get

$$(x+1)\ln(x+1) - x = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \int x^n dx = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \cdot \frac{x^{n+1}}{n+1} + C = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{n+1}}{n(n+1)} + C$$

(noting that $(-1)^{n-1} = (-1)^{n+1}$). To determine C , evaluate both sides at $x = 0$ to get

$$0 = \ln 1 - 0 = 0 + C$$

so that $C = 0$ and we get finally

$$(x+1)\ln(x+1) - x = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{n+1}}{n(n+1)}$$

(b) Evaluating the result of part(a) at $x = \frac{1}{2}$ gives

$$\begin{aligned} \frac{3}{2} \ln \frac{3}{2} - \frac{1}{2} &= \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n(n+1)2^{n+1}} \\ &= \frac{1}{1 \cdot 2 \cdot 2^2} - \frac{1}{2 \cdot 3 \cdot 2^3} + \frac{1}{3 \cdot 4 \cdot 2^4} - \frac{1}{4 \cdot 5 \cdot 2^5} + \dots \end{aligned}$$

(c)

$$\begin{aligned} S_4 &= \frac{1}{1 \cdot 2 \cdot 2^2} - \frac{1}{2 \cdot 3 \cdot 2^3} + \frac{1}{3 \cdot 4 \cdot 2^4} - \frac{1}{4 \cdot 5 \cdot 2^5} = 0.1078125 \\ a_5 &= \frac{1}{5 \cdot 6 \cdot 2^6} \approx 0.0005208 \\ \frac{3}{2} \ln \frac{3}{2} - \frac{1}{2} &\approx 0.10819766 \end{aligned}$$

and

$$\left| S_4 - \frac{3}{2} \ln \frac{3}{2} - \frac{1}{2} \right| \approx 0.0003852 < a_5$$

48. Prove that for $|x| < 1$,

$$\int \frac{dx}{x^4+1} = x - \frac{x^5}{5} + \frac{x^9}{9} - \dots$$

Use the first two terms to approximate $\int_0^{1/2} dx/(x^4+1)$ numerically. Use the fact that you have an alternating series to show that the error in this approximation is at most 0.00022.**SOLUTION** Substitute $-x^4$ for x in Eq. (2) to get

$$\frac{1}{1+x^4} = \sum_{n=0}^{\infty} (-x^4)^n = \sum_{n=0}^{\infty} (-1)^n x^{4n}$$

This is valid for $|x| < 1$, so for x in that range we can integrate the right-hand side term by term to get

$$\begin{aligned} \int \frac{1}{1+x^4} dx &= \sum_{n=0}^{\infty} \int (-1)^n x^{4n} dx = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+1}}{4n+1} + C \\ &= x - \frac{x^5}{5} + \frac{x^9}{9} - \frac{x^{13}}{13} + \dots + C \end{aligned}$$

Using the first two terms, we have

$$\int_0^{1/2} \frac{1}{1+x^4} dx \approx \frac{1}{2} - \frac{1}{2^5 \cdot 5} = \frac{79}{160} = 0.49375$$

Since this is an alternating series, the error in the approximation is bounded by the first unused term, so by

$$\frac{1}{2^9 \cdot 9} = \frac{1}{4608} \approx 0.000217 < 0.00022$$

49. Use the result of Example 7 to show that

$$F(x) = \frac{x^2}{1 \cdot 2} - \frac{x^4}{3 \cdot 4} + \frac{x^6}{5 \cdot 6} - \frac{x^8}{7 \cdot 8} + \cdots$$

is an antiderivative of $f(x) = \tan^{-1} x$ satisfying $F(0) = 0$. What is the radius of convergence of this power series?

SOLUTION For $-1 < x < 1$, which is the interval of convergence for the power series for arctangent, we can integrate term-by-term, so integrate that power series to get

$$\begin{aligned} F(x) &= \int \tan^{-1} x \, dx = \sum_{n=0}^{\infty} \int \frac{(-1)^n x^{2n+1}}{2n+1} \, dx = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+2}}{(2n+1)(2n+2)} \\ &= \frac{x^2}{1 \cdot 2} - \frac{x^4}{3 \cdot 4} + \frac{x^6}{5 \cdot 6} - \frac{x^8}{7 \cdot 8} + \cdots + C \end{aligned}$$

If we assume $F(0) = 0$, then we have $C = 0$. The radius of convergence of this power series is the same as that of the original power series, which is 1.

50. Verify that function $F(x) = x \tan^{-1} x - \frac{1}{2} \log(x^2 + 1)$ is an antiderivative of $f(x) = \tan^{-1} x$ satisfying $F(0) = 0$. Then use the result of Exercise 49 with $x = \frac{\pi}{6}$ to show that

$$\frac{\pi}{6\sqrt{3}} - \frac{1}{2} \ln \frac{4}{3} = \frac{1}{1 \cdot 2(3)} - \frac{1}{3 \cdot 4(3^2)} + \frac{1}{5 \cdot 6(3^3)} - \frac{1}{7 \cdot 8(3^4)} + \cdots$$

Use a calculator to compare the value of the left-hand side with the partial sum S_4 of the series on the right.

SOLUTION We have

$$F'(x) = \tan^{-1} x + \frac{x}{1+x^2} - \frac{1}{2} \cdot \frac{1}{x^2+1} \cdot 2x = \tan^{-1} x + \frac{x}{1+x^2} - \frac{x}{1+x^2} = \tan^{-1} x$$

so that $F(x)$ is an antiderivative of $\tan^{-1} x$, and clearly $F(0) = 0$. So applying Exercise 49 for this F , and setting $x = \frac{1}{\sqrt{3}}$, gives

$$\begin{aligned} \frac{1}{\sqrt{3}} \tan^{-1} \frac{1}{\sqrt{3}} - \frac{1}{2} \ln \left(\frac{1}{3} + 1 \right) &= \frac{\pi}{6\sqrt{3}} - \frac{1}{2} \ln \frac{4}{3} \\ &= \frac{(1/\sqrt{3})^2}{1 \cdot 2} - \frac{(1/\sqrt{3})^4}{3 \cdot 4} + \frac{(1/\sqrt{3})^6}{5 \cdot 6} - \frac{(1/\sqrt{3})^8}{7 \cdot 8} + \cdots \\ &= \frac{1}{1 \cdot 2(3)} - \frac{1}{3 \cdot 4(3^2)} + \frac{1}{5 \cdot 6(3^3)} - \frac{1}{7 \cdot 8(3^4)} + \cdots \end{aligned}$$

Now, we have

$$\begin{aligned} S_4 &= \frac{1}{1 \cdot 2(3)} - \frac{1}{3 \cdot 4(3^2)} + \frac{1}{5 \cdot 6(3^3)} - \frac{1}{7 \cdot 8(3^4)} = \frac{3593}{22680} \approx 0.1548215 \\ \frac{\pi}{6\sqrt{3}} - \frac{1}{2} \ln \frac{4}{3} &\approx 0.158459 \end{aligned}$$

so the two differ by less than 0.00004.

51. Evaluate $\sum_{n=1}^{\infty} \frac{n}{2^n}$. *Hint:* Use differentiation to show that

$$(1-x)^{-2} = \sum_{n=1}^{\infty} nx^{n-1} \quad (\text{for } |x| < 1)$$

SOLUTION Differentiate both sides of Eq. (2) to obtain

$$\frac{1}{(1-x)^2} = \sum_{n=1}^{\infty} nx^{n-1}.$$

Setting $x = \frac{1}{2}$ then yields

$$\sum_{n=1}^{\infty} \frac{n}{2^{n-1}} = \frac{1}{\left(1 - \frac{1}{2}\right)^2} = 4.$$

Divide this equation by 2 to obtain

$$\sum_{n=1}^{\infty} \frac{n}{2^n} = 2.$$

52. Use the power series for $(1+x^2)^{-1}$ and differentiation to prove that for $|x| < 1$,

$$\frac{2x}{(x^2+1)^2} = \sum_{n=1}^{\infty} (-1)^{n-1} (2n)x^{2n-1}$$

SOLUTION From Exercise 39, we know that for $-1 < x < 1$,

$$\frac{1}{1+x^2} = \sum_{n=0}^{\infty} (-1)^n x^{2n}$$

Thus for x in this range, we can differentiate both sides, and differentiate the right-hand side term by term, to get

$$\frac{d}{dx} \frac{1}{1+x^2} = \frac{-2x}{(x^2+1)^2} = \sum_{n=1}^{\infty} (-1)^n 2nx^{2n-1}$$

(Note the change in the lower limit of summation, since the $n = 0$ term is a constant, whose derivative is zero). Cancelling the minus sign on the left gives

$$\frac{2x}{(x^2+1)^2} = \sum_{n=1}^{\infty} (-1)^{n-1} (2n)x^{2n-1}$$

53. Show that the following series converges absolutely for $|x| < 1$ and compute its sum:

$$F(x) = 1 - x - x^2 + x^3 - x^4 - x^5 + x^6 - x^7 - x^8 + \dots$$

Hint: Write $F(x)$ as a sum of three geometric series with common ratio x^3 .

SOLUTION Because the coefficients in the power series are all ± 1 , we find

$$r = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1.$$

The radius of convergence is therefore $R = r^{-1} = 1$, and the series converges absolutely for $|x| < 1$.

By Exercise 43 of Section 11.4, any rearrangement of the terms of an absolutely convergent series yields another absolutely convergent series with the same sum as the original series. Following the hint, we now rearrange the terms of $F(x)$ as the sum of three geometric series:

$$\begin{aligned} F(x) &= (1 + x^3 + x^6 + \dots) - (x + x^4 + x^7 + \dots) - (x^2 + x^5 + x^8 + \dots) \\ &= \sum_{n=0}^{\infty} (x^3)^n - \sum_{n=0}^{\infty} x(x^3)^n - \sum_{n=0}^{\infty} x^2(x^3)^n = \frac{1}{1-x^3} - \frac{x}{1-x^3} - \frac{x^2}{1-x^3} = \frac{1-x-x^2}{1-x^3}. \end{aligned}$$

54. Show that for $|x| < 1$,

$$\frac{1+2x}{1+x+x^2} = 1+x-2x^2+x^3+x^4-2x^5+x^6+x^7-2x^8+\dots$$

Hint: Use the hint from Exercise 53.

SOLUTION The terms in the series on the right-hand side are either of the form x^n or $-2x^n$ for some n . Because

$$\lim_{n \rightarrow \infty} \sqrt[n]{2} = \lim_{n \rightarrow \infty} \sqrt[n]{1} = 1,$$

it follows that

$$\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = |x|.$$

Hence, by the Root Test, the series converges absolutely for $|x| < 1$.

By Exercise 43 of Section 11.4, any rearrangement of the terms of an absolutely convergent series yields another absolutely convergent series with the same sum as the original series. If we let S denote the sum of the series, then

$$\begin{aligned} S &= (1+x^3+x^6+\dots) + (x+x^4+x^7+\dots) - 2(x^2+x^5+x^8+\dots) \\ &= \frac{1}{1-x^3} + \frac{x}{1-x^3} - \frac{2x^2}{1-x^3} = \frac{1+x-2x^2}{1-x^3} = \frac{(1-x)(2x+1)}{(1-x)(1+x+x^2)} = \frac{2x+1}{1+x+x^2}. \end{aligned}$$

55. Find all values of x such that $\sum_{n=1}^{\infty} \frac{x^{n^2}}{n!}$ converges.

SOLUTION With $a_n = \frac{x^{n^2}}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|^{(n+1)^2}}{(n+1)!} \cdot \frac{n!}{|x|^{n^2}} = \frac{|x|^{2n+1}}{n+1}.$$

if $|x| \leq 1$, then

$$\lim_{n \rightarrow \infty} \frac{|x|^{2n+1}}{n+1} = 0,$$

and the series converges absolutely. On the other hand, if $|x| > 1$, then

$$\lim_{n \rightarrow \infty} \frac{|x|^{2n+1}}{n+1} = \infty,$$

and the series diverges. Thus, $\sum_{n=1}^{\infty} \frac{x^{n^2}}{n!}$ converges for $-1 \leq x \leq 1$ and diverges elsewhere.

56. Find all values of x such that the following series converges:

$$F(x) = 1 + 3x + x^2 + 27x^3 + x^4 + 243x^5 + \dots$$

SOLUTION Observe that $F(x)$ can be written as the sum of two geometric series:

$$F(x) = (1 + x^2 + x^4 + \dots) + (3x + 27x^3 + 243x^5 + \dots) = \sum_{n=0}^{\infty} (x^2)^n + \sum_{n=0}^{\infty} 3x(9x^2)^n$$

The first geometric series converges for $|x^2| < 1$, or $|x| < 1$; the second geometric series converges for $|9x^2| < 1$, or $|x| < \frac{1}{3}$. Since both geometric series must converge for $F(x)$ to converge, we find that $F(x)$ converges for $|x| < \frac{1}{3}$, the intersection of the intervals of convergence for the two geometric series.

57. Find a power series $P(x) = \sum_{n=0}^{\infty} a_n x^n$ satisfying the differential equation $y' = -y$ with initial condition $y(0) = 1$.

Then use Theorem 1 of Section 5.8 to conclude that $P(x) = e^{-x}$.

SOLUTION Let $P(x) = \sum_{n=0}^{\infty} a_n x^n$ and note that $P(0) = a_0$; thus, to satisfy the initial condition $P(0) = 1$, we must take $a_0 = 1$. Now,

$$P'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1},$$

so

$$P'(x) + P(x) = \sum_{n=1}^{\infty} na_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} [(n+1)a_{n+1} + a_n] x^n.$$

In order for this series to be equal to zero, the coefficient of x^n must be equal to zero for each n ; thus

$$(n+1)a_{n+1} + a_n = 0 \quad \text{or} \quad a_{n+1} = -\frac{a_n}{n+1}.$$

Starting from $a_0 = 1$, we then calculate

$$a_1 = -\frac{a_0}{1} = -1;$$

$$a_2 = -\frac{a_1}{2} = \frac{1}{2};$$

$$a_3 = -\frac{a_2}{3} = -\frac{1}{6} = -\frac{1}{3!};$$

and, in general,

$$a_n = (-1)^n \frac{1}{n!}.$$

Hence,

$$P(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n!}.$$

The solution to the initial value problem $y' = -y$, $y(0) = 1$ is $y = e^{-x}$. Because this solution is unique, it follows that

$$P(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{n!} = e^{-x}.$$

58. Let $C(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$.

(a) Show that $C(x)$ has an infinite radius of convergence.

(b) Prove that $C(x)$ and $f(x) = \cos x$ are both solutions of $y'' = -y$ with initial conditions $y(0) = 1$, $y'(0) = 0$. This initial value problem has a unique solution, so we have $C(x) = \cos x$ for all x .

SOLUTION

(a) Consider the series

$$C(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}.$$

With $a_n = (-1)^n \frac{x^{2n}}{(2n)!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{|x|^{2n+2}}{(2n+2)!} \cdot \frac{(2n)!}{|x|^{2n}} = \frac{|x|^2}{(2n+2)(2n+1)},$$

and

$$r = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0.$$

The radius of convergence for $C(x)$ is therefore $R = r^{-1} = \infty$.

(b) Differentiating the series defining $C(x)$ term-by-term, we find

$$C'(x) = \sum_{n=1}^{\infty} (-1)^n (2n) \frac{x^{2n-1}}{(2n)!} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-1}}{(2n-1)!}$$

and

$$\begin{aligned} C''(x) &= \sum_{n=1}^{\infty} (-1)^n (2n-1) \frac{x^{2n-2}}{(2n-1)!} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-2}}{(2n-2)!} \\ &= \sum_{n=0}^{\infty} (-1)^{n+1} \frac{x^{2n}}{(2n)!} = -\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = -C(x). \end{aligned}$$

Moreover, $C(0) = 1$ and $C'(0) = 0$.

59. Use the power series for $y = e^x$ to show that

$$\frac{1}{e} = \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \cdots$$

Use your knowledge of alternating series to find an N such that the partial sum S_N approximates e^{-1} to within an error of at most 10^{-3} . Confirm this using a calculator to compute both S_N and e^{-1} .

SOLUTION Recall that the series for e^x is

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Setting $x = -1$ yields

$$e^{-1} = 1 - 1 + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \cdots = \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \cdots$$

This is an alternating series with $a_n = \frac{1}{(n+1)!}$. The error in approximating e^{-1} with the partial sum S_N is therefore bounded by

$$|S_N - e^{-1}| \leq a_{N+1} = \frac{1}{(N+2)!}$$

To make the error at most 10^{-3} , we must choose N such that

$$\frac{1}{(N+2)!} \leq 10^{-3} \quad \text{or} \quad (N+2)! \geq 1000.$$

For $N = 4$, $(N+2)! = 6! = 720 < 1000$, but for $N = 5$, $(N+2)! = 7! = 5040$; hence, $N = 5$ is the smallest value that satisfies the error bound. The corresponding approximation is

$$S_5 = \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \frac{1}{5!} + \frac{1}{6!} = 0.368055555$$

Now, $e^{-1} = 0.367879441$, so

$$|S_5 - e^{-1}| = 1.761 \times 10^{-4} < 10^{-3}.$$

60. Let $P(x) = \sum_{n=0}^{\infty} a_n x^n$ be a power series solution to $y' = 2xy$ with initial condition $y(0) = 1$.

(a) Show that the odd coefficients a_{2k+1} are all zero.

(b) Prove that $a_{2k} = a_{2k-2}/k$ and use this result to determine the coefficients a_{2k} .

SOLUTION Let $P(x) = \sum_{n=0}^{\infty} a_n x^n$ and note that $P(0) = a_0$; thus, to satisfy the initial condition $P(0) = 1$, we must take $a_0 = 1$. Now,

$$P'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1},$$

so

$$\begin{aligned} P'(x) - 2xP(x) &= \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} 2a_n x^{n+1} = \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=2}^{\infty} 2a_{n-2} x^{n-1} \\ &= a_1 + \sum_{n=2}^{\infty} [n a_n - 2a_{n-2}] x^{n-1}. \end{aligned}$$

In order for this series to be equal to zero, the coefficient of x^n must be equal to zero for each n ; thus, $a_1 = 0$ and

$$n a_n - 2a_{n-2} = 0 \quad \text{or} \quad a_n = \frac{2a_{n-2}}{n}.$$

(a) We know that $a_1 = 0$ and

$$a_n = \frac{2a_{n-2}}{n}.$$

Thus,

$$a_3 = \frac{2a_1}{3} = 0;$$

$$a_5 = \frac{2a_3}{5} = 0;$$

$$a_7 = \frac{2a_5}{7} = 0;$$

and, in general, $a_{2k+1} = 0$ for all k .

(b) Replace n by $2k$ in the equation

$$a_n = \frac{2a_{n-2}}{n} \quad \text{to obtain} \quad a_{2k} = \frac{2a_{2k-2}}{2k} = \frac{a_{2k-2}}{k}.$$

Starting from $a_0 = 1$, we then calculate

$$a_2 = \frac{a_0}{1} = 1 = \frac{1}{1!};$$

$$a_4 = \frac{a_2}{2} = \frac{1}{2} = \frac{1}{2!};$$

$$a_6 = \frac{a_4}{3} = \frac{1}{6} = \frac{1}{3!};$$

and, in general, $a_{2k} = \frac{1}{k!}$.

61. Find a power series $P(x)$ satisfying the differential equation

$$y'' - xy' + y = 0$$

9

with initial condition $y(0) = 1$, $y'(0) = 0$. What is the radius of convergence of the power series?

SOLUTION Let $P(x) = \sum_{n=0}^{\infty} a_n x^n$. Then

$$P'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} \quad \text{and} \quad P''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}.$$

Note that $P(0) = a_0$ and $P'(0) = a_1$; in order to satisfy the initial conditions $P(0) = 1$, $P'(0) = 0$, we must have $a_0 = 1$ and $a_1 = 0$. Now,

$$\begin{aligned} P''(x) - xP'(x) + P(x) &= \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n \\ &= \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n \\ &= 2a_2 + a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1)a_{n+2} - n a_n + a_n] x^n. \end{aligned}$$

In order for this series to be equal to zero, the coefficient of x^n must be equal to zero for each n ; thus, $2a_2 + a_0 = 0$ and $(n+2)(n+1)a_{n+2} - (n-1)a_n = 0$, or

$$a_2 = -\frac{1}{2}a_0 \quad \text{and} \quad a_{n+2} = \frac{n-1}{(n+2)(n+1)}a_n.$$

Starting from $a_1 = 0$, we calculate

$$a_3 = \frac{1-1}{(3)(2)}a_1 = 0;$$

$$a_5 = \frac{2}{(5)(4)}a_3 = 0;$$

$$a_7 = \frac{4}{(7)(6)}a_5 = 0;$$

and, in general, all of the odd coefficients are zero. As for the even coefficients, we have $a_0 = 1$, $a_2 = -\frac{1}{2}$,

$$a_4 = \frac{1}{(4)(3)}a_2 = -\frac{1}{4!};$$

$$a_6 = \frac{3}{(6)(5)}a_4 = -\frac{3}{6!};$$

$$a_8 = \frac{5}{(8)(7)}a_6 = -\frac{15}{8!}$$

and so on. Thus,

$$P(x) = 1 - \frac{1}{2}x^2 - \frac{1}{4!}x^4 - \frac{3}{6!}x^6 - \frac{15}{8!}x^8 - \dots$$

To determine the radius of convergence, treat this as a series in the variable x^2 , and observe that

$$r = \lim_{k \rightarrow \infty} \left| \frac{a_{2k+2}}{a_{2k}} \right| = \lim_{k \rightarrow \infty} \frac{2k-1}{(2k+2)(2k+1)} = 0.$$

Thus, the radius of convergence is $R = r^{-1} = \infty$.

62. Find a power series satisfying Eq. (9) with initial condition $y(0) = 0$, $y'(0) = 1$.

SOLUTION Let $P(x) = \sum_{n=0}^{\infty} a_n x^n$ be a solution to Eq. (9). From the previous exercise, we know that

$$a_2 = -\frac{1}{2}a_0 \quad \text{and} \quad a_{n+2} = \frac{n-1}{(n+2)(n+1)}a_n.$$

To satisfy the initial condition $P(0) = 0$, $P'(0) = 1$, we must have $a_0 = 0$ and $a_1 = 1$. Then

$$a_2 = -\frac{1}{2}a_0 = 0;$$

$$a_4 = \frac{1}{(4)(3)}a_2 = 0;$$

$$a_6 = \frac{3}{(6)(5)}a_4 = 0;$$

and, in general, all of the even coefficients are zero. As in the previous exercise, all of the odd coefficients past a_1 are also equal to zero. Thus,

$$P(x) = x.$$

63. Prove that

$$J_2(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k+2} k! (k+3)!} x^{2k+2}$$

is a solution of the Bessel differential equation of order 2:

$$x^2 y'' + x y' + (x^2 - 4)y = 0$$

SOLUTION Let $J_2(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k+2} k! (k+2)!} x^{2k+2}$. Then

$$J_2'(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (k+1)}{2^{2k+1} k! (k+2)!} x^{2k+1}$$


$$J_2''(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (k+1)(2k+1)}{2^{2k+1} k! (k+2)!} x^{2k}$$

and

$$x^2 J_2''(x) + x J_2'(x) + (x^2 - 4)J_2(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (k+1)(2k+1)}{2^{2k+1} k! (k+2)!} x^{2k+2} + \sum_{k=0}^{\infty} \frac{(-1)^k (k+1)}{2^{2k+1} k! (k+2)!} x^{2k+2}$$

$$- \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k+2} k! (k+2)!} x^{2k+4} - \sum_{k=0}^{\infty} \frac{(-1)^k}{2^{2k} k! (k+2)!} x^{2k+2}$$

$$\begin{aligned}
&= \sum_{k=0}^{\infty} \frac{(-1)^k k(k+2)}{2^{2k} k!(k+2)!} x^{2k+2} + \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2^{2k} (k-1)!(k+1)!} x^{2k+2} \\
&= \sum_{k=1}^{\infty} \frac{(-1)^k}{2^{2k} (k-1)!(k+1)!} x^{2k+2} - \sum_{k=1}^{\infty} \frac{(-1)^k}{2^{2k} (k-1)!(k+1)!} x^{2k+2} = 0.
\end{aligned}$$

64.  Why is it impossible to expand $f(x) = |x|$ as a power series that converges in an interval around $x = 0$? Explain using Theorem 2.

SOLUTION Suppose that there exists a $c > 0$ such that f can be represented by a power series on the interval $(-c, c)$; that is,

$$|x| = \sum_{n=0}^{\infty} a_n x^n$$

for $|x| < c$. Then it follows by Theorem 2 that $|x|$ is differentiable on $(-c, c)$. This contradicts the well known property that $f(x) = |x|$ is not differentiable at the point $x = 0$.

Further Insights and Challenges

65. Suppose that the coefficients of $F(x) = \sum_{n=0}^{\infty} a_n x^n$ are *periodic*; that is, for some whole number $M > 0$, we have $a_{M+n} = a_n$. Prove that $F(x)$ converges absolutely for $|x| < 1$ and that

$$F(x) = \frac{a_0 + a_1 x + \cdots + a_{M-1} x^{M-1}}{1 - x^M}$$

Hint: Use the hint for Exercise 53.

SOLUTION Suppose the coefficients of $F(x)$ are periodic, with $a_{M+n} = a_n$ for some whole number M and all n . The $F(x)$ can be written as the sum of M geometric series:

$$\begin{aligned}
F(x) &= a_0 (1 + x^M + x^{2M} + \cdots) + a_1 (x + x^{M+1} + x^{2M+1} + \cdots) + \\
&= a_2 (x^2 + x^{M+2} + x^{2M+2} + \cdots) + \cdots + a_{M-1} (x^{M-1} + x^{2M-1} + x^{3M-1} + \cdots) \\
&= \frac{a_0}{1 - x^M} + \frac{a_1 x}{1 - x^M} + \frac{a_2 x^2}{1 - x^M} + \cdots + \frac{a_{M-1} x^{M-1}}{1 - x^M} = \frac{a_0 + a_1 x + a_2 x^2 + \cdots + a_{M-1} x^{M-1}}{1 - x^M}.
\end{aligned}$$

As each geometric series converges absolutely for $|x| < 1$, it follows that $F(x)$ also converges absolutely for $|x| < 1$.

66. Continuity of Power Series Let $F(x) = \sum_{n=0}^{\infty} a_n x^n$ be a power series with radius of convergence $R > 0$.

(a) Prove the inequality

$$|x^n - y^n| \leq n|x - y|(|x|^{n-1} + |y|^{n-1}) \quad \boxed{10}$$

Hint: $x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \cdots + y^{n-1})$.

(b) Choose R_1 with $0 < R_1 < R$. Show that the infinite series $M = \sum_{n=0}^{\infty} 2n|a_n|R_1^n$ converges. *Hint:* Show that $n|a_n|R_1^n < |a_n|x^n$ for all n sufficiently large if $R_1 < x < R$.

(c) Use Eq. (10) to show that if $|x| < R_1$ and $|y| < R_1$, then $|F(x) - F(y)| \leq M|x - y|$.

(d) Prove that if $|x| < R$, then $F(x)$ is continuous at x . *Hint:* Choose R_1 such that $|x| < R_1 < R$. Show that if $\epsilon > 0$ is given, then $|F(x) - F(y)| \leq \epsilon$ for all y such that $|x - y| < \delta$, where δ is any positive number that is less than ϵ/M and $R_1 - |x|$ (see Figure 6).




FIGURE 6 If $x > 0$, choose $\delta > 0$ less than ϵ/M and $R_1 - x$.

SOLUTION

(a) Take the absolute value of both sides of the identity

$$x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \cdots + y^{n-1}),$$

and then apply the triangle inequality to obtain

$$|x^n - y^n| \leq |x - y| (|x|^{n-1} + |x|^{n-2}|y| + |x|^{n-3}|y|^2 + \cdots + |x||y|^{n-2} + |y|^{n-1}).$$

Now, if $|x| \leq |y|$ then $|x|^{n-k}|y|^{k-1} \leq |y|^{n-k}|y|^{k-1} = |y|^{n-1}$, and if $|y| \leq |x|$ then $|x|^{n-k}|y|^{k-1} \leq |x|^{n-k}|x|^{k-1} = |x|^{n-1}$. In either case, $|x|^{n-k}|y|^{k-1} \leq |x|^{n-1} + |y|^{n-1}$. Thus,

$$\begin{aligned} |x^n - y^n| &\leq |x - y| \left(|x|^{n-1} + (n-2)(|x|^{n-1} + |y|^{n-1}) + |y|^{n-1} \right) \\ &= (n-1)|x - y| \left(|x|^{n-1} + |y|^{n-1} \right) \leq n|x - y| \left(|x|^{n-1} + |y|^{n-1} \right). \end{aligned}$$

(b) Let $0 < R_1 < R$. Then,

$$\rho = \lim_{n \rightarrow \infty} \frac{2(n+1)|a_{n+1}|R_1^{n+1}}{2n|a_n|R_1^n} = R_1 \lim_{n \rightarrow \infty} \frac{n+1}{n} \cdot \left| \frac{a_{n+1}}{a_n} \right| = R_1 \cdot 1 \cdot \frac{1}{R} = \frac{R_1}{R} < 1.$$

Thus, the series $M = \sum_{n=0}^{\infty} 2n|a_n|R_1^n$ converges by the Ratio Test.

(c) Suppose $|x| < R_1$ and $|y| < R_1$. Then

$$\begin{aligned} |F(x) - F(y)| &= \left| \sum_{n=0}^{\infty} a_n x^n - \sum_{n=0}^{\infty} a_n y^n \right| \leq \sum_{n=0}^{\infty} |a_n| |x^n - y^n| \leq \sum_{n=0}^{\infty} n|a_n| |x - y| \left(|x|^{n-1} + |y|^{n-1} \right) \\ &\leq |x - y| \sum_{n=0}^{\infty} n|a_n| \left(R_1^{n-1} + R_1^{n-1} \right) = M|x - y| \end{aligned}$$

(d) Let $|x| < R$, and let R_1 be a number such that $|x| < R_1 < R$. Then by part (b), $M = \sum_{n=0}^{\infty} 2n|a_n|R_1^n$ is finite and by part (c)

$$|F(x) - F(y)| \leq M|x - y|$$

for $|y| < R_1$. Now, let $\epsilon > 0$, and choose $\delta > 0$ so that $\delta < \frac{\epsilon}{M}$ and $\delta < R_1 - |x|$. Then, whenever $|y - x| < \delta$,

$$|y| = |(y - x) + x| \leq |y - x| + |x| < \delta + |x| < R_1,$$

so

$$|F(x) - F(y)| < M|x - y| < M\delta < M \cdot \frac{\epsilon}{M} = \epsilon.$$

Thus, F is continuous at x .

10.7 Taylor Series

Preliminary Questions

1. Determine $f(0)$ and $f'''(0)$ for a function $f(x)$ with Maclaurin series

$$T(x) = 3 + 2x + 12x^2 + 5x^3 + \dots$$

SOLUTION The Maclaurin series for a function f has the form

$$f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

Matching this general expression with the given series, we find $f(0) = 3$ and $\frac{f'''(0)}{3!} = 5$. From this latter equation, it follows that $f'''(0) = 30$.

2. Determine $f(-2)$ and $f^{(4)}(-2)$ for a function with Taylor series

$$T(x) = 3(x+2) + (x+2)^2 - 4(x+2)^3 + 2(x+2)^4 + \dots$$

SOLUTION The Taylor series for a function f centered at $x = -2$ has the form

$$f(-2) + \frac{f'(-2)}{1!}(x+2) + \frac{f''(-2)}{2!}(x+2)^2 + \frac{f'''(-2)}{3!}(x+2)^3 + \frac{f^{(4)}(-2)}{4!}(x+2)^4 + \dots$$

Matching this general expression with the given series, we find $f(-2) = 0$ and $\frac{f^{(4)}(-2)}{4!} = 2$. From this latter equation, it follows that $f^{(4)}(-2) = 48$.

3. What is the easiest way to find the Maclaurin series for the function $f(x) = \sin(x^2)$?

SOLUTION The easiest way to find the Maclaurin series for $\sin(x^2)$ is to substitute x^2 for x in the Maclaurin series for $\sin x$.

4. Find the Taylor series for $f(x)$ centered at $c = 3$ if $f(3) = 4$ and $f'(x)$ has a Taylor expansion

$$f'(x) = \sum_{n=1}^{\infty} \frac{(x-3)^n}{n}$$

SOLUTION Integrating the series for $f'(x)$ term-by-term gives

$$f(x) = C + \sum_{n=1}^{\infty} \frac{(x-3)^{n+1}}{n(n+1)}.$$

Substituting $x = 3$ then yields

$$f(3) = C = 4;$$

so

$$f(x) = 4 + \sum_{n=1}^{\infty} \frac{(x-3)^{n+1}}{n(n+1)}.$$

5. Let $T(x)$ be the Maclaurin series of $f(x)$. Which of the following guarantees that $f(2) = T(2)$?

- (a) $T(x)$ converges for $x = 2$.
- (b) The remainder $R_k(2)$ approaches a limit as $k \rightarrow \infty$.
- (c) The remainder $R_k(2)$ approaches zero as $k \rightarrow \infty$.

SOLUTION The correct response is (c): $f(2) = T(2)$ if and only if the remainder $R_k(2)$ approaches zero as $k \rightarrow \infty$.

Exercises

1. Write out the first four terms of the Maclaurin series of $f(x)$ if

$$f(0) = 2, \quad f'(0) = 3, \quad f''(0) = 4, \quad f'''(0) = 12$$

SOLUTION The first four terms of the Maclaurin series of $f(x)$ are

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 = 2 + 3x + \frac{4}{2}x^2 + \frac{12}{6}x^3 = 2 + 3x + 2x^2 + 2x^3.$$

2. Write out the first four terms of the Taylor series of $f(x)$ centered at $c = 3$ if

$$f(3) = 1, \quad f'(3) = 2, \quad f''(3) = 12, \quad f'''(3) = 3$$

SOLUTION The first four terms of the Taylor series centered at $c = 3$ are:

$$\begin{aligned} f(3) + f'(3)(x-3) + \frac{f''(3)}{2!}(x-3)^2 + \frac{f'''(3)}{3!}(x-3)^3 &= 1 + 2(x-3) + \frac{12}{2}(x-3)^2 + \frac{3}{6}(x-3)^3 \\ &= 1 + 2(x-3) + 6(x-3)^2 + \frac{1}{2}(x-3)^3. \end{aligned}$$

In Exercises 3–18, find the Maclaurin series and find the interval on which the expansion is valid.

3. $f(x) = \frac{1}{1-2x}$

SOLUTION Substituting $2x$ for x in the Maclaurin series for $\frac{1}{1-x}$ gives

$$\frac{1}{1-2x} = \sum_{n=0}^{\infty} (2x)^n = \sum_{n=0}^{\infty} 2^n x^n.$$

This series is valid for $|2x| < 1$, or $|x| < \frac{1}{2}$.

4. $f(x) = \frac{x}{1-x^4}$

SOLUTION Substituting x^4 for x in the Maclaurin series for $\frac{1}{1-x}$ gives

$$\frac{1}{1-x^4} = \sum_{n=0}^{\infty} (x^4)^n = \sum_{n=0}^{\infty} x^{4n}.$$

Therefore

$$\frac{x}{1-x^4} = x \sum_{n=0}^{\infty} x^{4n} = \sum_{n=0}^{\infty} x^{4n+1}.$$

This series is valid for $|x^4| < 1$, or $|x| < 1$.

5. $f(x) = \cos 3x$

SOLUTION Substituting $3x$ for x in the Maclaurin series for $\cos x$ gives

$$\cos 3x = \sum_{n=0}^{\infty} (-1)^n \frac{(3x)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{9^n x^{2n}}{(2n)!}.$$

This series is valid for all x .

6. $f(x) = \sin(2x)$

SOLUTION Substituting $2x$ for x in the Maclaurin series for $\sin x$ gives

$$\sin 2x = \sum_{n=0}^{\infty} (-1)^n \frac{(2x)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{2^{2n+1} x^{2n+1}}{(2n+1)!}.$$

This series is valid for all x .

7. $f(x) = \sin(x^2)$

SOLUTION Substituting x^2 for x in the Maclaurin series for $\sin x$ gives

$$\sin x^2 = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{(2n+1)!}.$$

This series is valid for all x .

8. $f(x) = e^{4x}$

SOLUTION Substituting $4x$ for x in the Maclaurin series for e^x gives

$$e^{4x} = \sum_{n=0}^{\infty} \frac{(4x)^n}{n!} = \sum_{n=0}^{\infty} \frac{4^n x^n}{n!}.$$

This series is valid for all x .

9. $f(x) = \ln(1-x^2)$

SOLUTION Substituting $-x^2$ for x in the Maclaurin series for $\ln(1+x)$ gives

$$\ln(1-x^2) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (-x^2)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{2n-1} x^{2n}}{n} = - \sum_{n=1}^{\infty} \frac{x^{2n}}{n}.$$

This series is valid for $|x| < 1$.

10. $f(x) = (1-x)^{-1/2}$

SOLUTION Substituting $-x$ for x and using $a = -\frac{1}{2}$ in the Binomial series gives

$$(1-x)^{-1/2} = \sum_{n=0}^{\infty} \binom{-\frac{1}{2}}{n} (-x)^n = \sum_{n=0}^{\infty} (-1)^n \binom{-\frac{1}{2}}{n} x^n.$$

This series is valid for $|x| < 1$.

11. $f(x) = \tan^{-1}(x^2)$

SOLUTION Substituting x^2 for x in the Maclaurin series for $\tan^{-1} x$ gives

$$\tan^{-1}(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1}.$$

This series is valid for $|x| \leq 1$.

12. $f(x) = x^2 e^{x^2}$

SOLUTION First substitute x^2 for x in the Maclaurin series for e^x to obtain

$$e^{x^2} = \sum_{n=0}^{\infty} \frac{(x^2)^n}{n!} = \sum_{n=0}^{\infty} \frac{x^{2n}}{n!}.$$

Now, multiply by x^2 to obtain

$$x^2 e^{x^2} = x^2 \sum_{n=0}^{\infty} \frac{x^{2n}}{n!} = \sum_{n=0}^{\infty} \frac{x^{2n+2}}{n!}.$$

This series is valid for all x .

13. $f(x) = e^{x-2}$

SOLUTION $e^{x-2} = e^{-2} e^x$; thus,

$$e^{x-2} = e^{-2} \sum_{n=0}^{\infty} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{x^n}{e^2 n!}.$$

This series is valid for all x .

14. $f(x) = \frac{1 - \cos x}{x}$

SOLUTION $\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$, so

$$\frac{1 - \cos x}{x} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n)!}$$

This series is valid for all x .

15. $f(x) = \ln(1 - 5x)$

SOLUTION Substituting $-5x$ for x in the Maclaurin series for $\ln(1 + x)$ gives

$$\ln(1 - 5x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (-5x)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{2n-1} 5^n x^n}{n} = - \sum_{n=1}^{\infty} \frac{5^n x^n}{n}.$$

This series is valid for $|5x| < 1$, or $|x| < \frac{1}{5}$, and for $x = -\frac{1}{5}$.

16. $f(x) = (x^2 + 2x)e^x$

SOLUTION Using the Maclaurin series for e^x , we find

$$\begin{aligned} (x^2 + 2x)e^x &= x^2 \sum_{n=0}^{\infty} \frac{x^n}{n!} + 2x \sum_{n=0}^{\infty} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{x^{n+2}}{n!} + \sum_{n=0}^{\infty} \frac{2x^{n+1}}{n!} = 2x + \sum_{n=1}^{\infty} \left(\frac{1}{(n-1)!} + \frac{2}{n!} \right) x^{n+1} \\ &= 2x + \sum_{n=1}^{\infty} \frac{n+2}{n!} x^{n+1} = \sum_{n=0}^{\infty} \frac{n+2}{n!} x^{n+1}. \end{aligned}$$

This series is valid for all x .

17. $f(x) = \sinh x$

SOLUTION Recall that

$$\sinh x = \frac{1}{2}(e^x - e^{-x}).$$

Therefore,

$$\sinh x = \frac{1}{2} \left(\sum_{n=0}^{\infty} \frac{x^n}{n!} - \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right) = \sum_{n=0}^{\infty} \frac{x^n}{2(n!)} (1 - (-1)^n).$$

Now,

$$1 - (-1)^n = \begin{cases} 0, & n \text{ even} \\ 2, & n \text{ odd} \end{cases}$$

so

$$\sinh x = \sum_{k=0}^{\infty} 2 \frac{x^{2k+1}}{2(2k+1)!} = \sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!}.$$

This series is valid for all x .

18. $f(x) = \cosh x$

SOLUTION Recall that

$$\cosh x = \frac{1}{2}(e^x + e^{-x}).$$

Therefore,

$$\cosh x = \frac{1}{2} \left(\sum_{n=0}^{\infty} \frac{x^n}{n!} + \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right) = \sum_{n=0}^{\infty} \frac{x^n}{2(n!)} (1 + (-1)^n).$$

Now,

$$1 + (-1)^n = \begin{cases} 0, & n \text{ odd} \\ 2, & n \text{ even} \end{cases}$$

so

$$\cosh x = \sum_{k=0}^{\infty} 2 \frac{x^{2k}}{2(2k)!} = \sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!}.$$

This series is valid for all x .

In Exercises 19–28, find the terms through degree four of the Maclaurin series of $f(x)$. Use multiplication and substitution as necessary.

19. $f(x) = e^x \sin x$

SOLUTION Multiply the fourth-order Taylor Polynomials for e^x and $\sin x$:

$$\begin{aligned} & \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} \right) \left(x - \frac{x^3}{6} \right) \\ &= x + x^2 - \frac{x^3}{6} + \frac{x^3}{2} - \frac{x^4}{6} + \frac{x^4}{6} + \text{higher-order terms} \\ &= x + x^2 + \frac{x^3}{3} + \text{higher-order terms.} \end{aligned}$$

The terms through degree four in the Maclaurin series for $f(x) = e^x \sin x$ are therefore

$$x + x^2 + \frac{x^3}{3}.$$

20. $f(x) = e^x \ln(1-x)$

SOLUTION Multiply the fourth order Taylor Polynomials for e^x and $\ln(1-x)$:

$$\begin{aligned} & \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} \right) \left(-x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} \right) \\ &= -x - \frac{x^2}{2} - x^2 - \frac{x^3}{3} - \frac{x^3}{2} - \frac{x^3}{2} - \frac{x^4}{4} - \frac{x^4}{3} - \frac{x^4}{4} - \frac{x^4}{6} + \text{higher-order terms} \\ &= -x - \frac{3x^2}{2} - \frac{4x^3}{3} - x^4 + \text{higher-order terms.} \end{aligned}$$

The first four terms of the Maclaurin series for $f(x) = e^x \ln(1-x)$ are therefore

$$-x - \frac{3x^2}{2} - \frac{4x^3}{3} - x^4.$$

21. $f(x) = \frac{\sin x}{1-x}$

SOLUTION Multiply the fourth order Taylor Polynomials for $\sin x$ and $\frac{1}{1-x}$:

$$\begin{aligned} & \left(x - \frac{x^3}{6}\right) \left(1 + x + x^2 + x^3 + x^4\right) \\ &= x + x^2 - \frac{x^3}{6} + x^3 + x^4 - \frac{x^4}{6} + \text{higher-order terms} \\ &= x + x^2 + \frac{5x^3}{6} + \frac{5x^4}{6} + \text{higher-order terms.} \end{aligned}$$

The terms through order four of the Maclaurin series for $f(x) = \frac{\sin x}{1-x}$ are therefore

$$x + x^2 + \frac{5x^3}{6} + \frac{5x^4}{6}.$$

22. $f(x) = \frac{1}{1 + \sin x}$

SOLUTION Substituting $\sin x$ for x in the Maclaurin series for $\frac{1}{1-x}$ and then using the Maclaurin series for $\sin x$ gives

$$\begin{aligned} \frac{1}{1 + \sin x} &= 1 - \sin x + \sin^2 x - \sin^3 x + \sin^4 x - \dots \\ &= 1 - \left(x - \frac{x^3}{6} + \dots\right) + \left(x - \frac{x^3}{6} + \dots\right)^2 - \left(x - \frac{x^3}{6} + \dots\right)^3 + \left(x - \frac{x^3}{6} + \dots\right)^4 - \dots \\ &= 1 - x + \frac{x^3}{6} + x^2 - \frac{x^4}{3} - x^3 + x^4 = 1 - x + x^2 - \frac{5x^3}{6} + \frac{2x^4}{3} \end{aligned}$$

Therefore, the terms of the Maclaurin series for $f(x) = \frac{1}{1 + \sin x}$ through order four are

$$1 - x + x^2 - \frac{5x^3}{6} + \frac{2x^4}{3}.$$

23. $f(x) = (1+x)^{1/4}$

SOLUTION The first five generalized binomial coefficients for $a = \frac{1}{4}$ are

$$1, \quad \frac{1}{4}, \quad \frac{\frac{1}{4} \binom{-3}{4}}{2!} = -\frac{3}{32}, \quad \frac{\frac{1}{4} \binom{-3}{4} \binom{-7}{4}}{3!} = \frac{7}{128}, \quad \frac{\frac{1}{4} \binom{-3}{4} \binom{-7}{4} \binom{-11}{4}}{4!} = \frac{-77}{2048}$$

Therefore, the first four terms in the binomial series for $(1+x)^{1/4}$ are

$$1 + \frac{1}{4}x - \frac{3}{32}x^2 + \frac{7}{128}x^3 - \frac{77}{2048}x^4$$

24. $f(x) = (1+x)^{-3/2}$

SOLUTION The first five generalized binomial coefficients for $a = -\frac{3}{2}$ are

$$1, \quad -\frac{3}{2}, \quad \frac{-\frac{3}{2} \binom{-5}{2}}{2!} = \frac{15}{8}, \quad \frac{-\frac{3}{2} \binom{-5}{2} \binom{-7}{2}}{3!} = -\frac{35}{16}, \quad \frac{-\frac{3}{2} \binom{-5}{2} \binom{-7}{2} \binom{-9}{2}}{4!} = \frac{315}{128}.$$

Therefore, the first five terms in the binomial series for $f(x) = (1+x)^{-3/2}$ are

$$1 - \frac{3}{2}x + \frac{15}{8}x^2 - \frac{35}{16}x^3 + \frac{315}{128}x^4.$$

25. $f(x) = e^x \tan^{-1} x$

SOLUTION Using the Maclaurin series for e^x and $\tan^{-1} x$, we find

$$\begin{aligned} e^x \tan^{-1} x &= \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots\right) \left(x - \frac{x^3}{3} + \dots\right) = x + x^2 - \frac{x^3}{3} + \frac{x^3}{2} + \frac{x^4}{6} - \frac{x^4}{3} + \dots \\ &= x + x^2 + \frac{1}{6}x^3 - \frac{1}{6}x^4 + \dots \end{aligned}$$

26. $f(x) = \sin(x^3 - x)$

SOLUTION Substitute $x^3 - x$ into the first two terms of the Maclaurin series for $\sin x$:

$$(x^3 - x) - \frac{(x^3 - x)^3}{3!} = x^3 - x - \frac{x^9 - 3x^7 + 3x^5 - x^3}{3!}$$

so that the terms of the Maclaurin series for $\sin(x^3 - x)$ through degree four are

$$-x + \frac{7}{6}x^3$$

27. $f(x) = e^{\sin x}$

SOLUTION Substituting $\sin x$ for x in the Maclaurin series for e^x and then using the Maclaurin series for $\sin x$, we find

$$\begin{aligned} e^{\sin x} &= 1 + \sin x + \frac{\sin^2 x}{2} + \frac{\sin^3 x}{6} + \frac{\sin^4 x}{24} + \dots \\ &= 1 + \left(x - \frac{x^3}{6} + \dots\right) + \frac{1}{2} \left(x - \frac{x^3}{6} + \dots\right)^2 + \frac{1}{6} (x - \dots)^3 + \frac{1}{24} (x - \dots)^4 \\ &= 1 + x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{6}x^3 - \frac{1}{6}x^4 + \frac{1}{24}x^4 + \dots \\ &= 1 + x + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \dots \end{aligned}$$

28. $f(x) = e^{(e^x)}$

SOLUTION With $f(x) = e^{(e^x)}$, we find

$$\begin{aligned} f'(x) &= e^{(e^x)} \cdot e^x \\ f''(x) &= e^{(e^x)} \cdot e^x + e^{(e^x)} \cdot e^{2x} = e^{(e^x)} (e^{2x} + e^x) \\ f'''(x) &= e^{(e^x)} (2e^{2x} + e^x) + e^{(e^x)} (e^{2x} + e^x) e^x \\ &= e^{(e^x)} (e^{3x} + 3e^{2x} + e^x) \\ f^{(4)}(x) &= e^{(e^x)} (3e^{3x} + 6e^{2x} + e^x) + e^{(e^x)} (e^{3x} + 3e^{2x} + e^x) e^x \\ &= e^{(e^x)} (e^{4x} + 6e^{3x} + 7e^{2x} + e^x) \end{aligned}$$

and

$$f(0) = e, \quad f'(0) = e, \quad f''(0) = 2e, \quad f'''(0) = 5e, \quad f^{(4)}(0) = 15e.$$

Therefore, the first four terms of the Maclaurin for $f(x) = e^{(e^x)}$ are

$$e + ex + ex^2 + \frac{5e}{6}x^3 + \frac{5e}{8}x^4.$$

In Exercises 29–38, find the Taylor series centered at c and find the interval on which the expansion is valid.

29. $f(x) = \frac{1}{x}$, $c = 1$

SOLUTION Write

$$\frac{1}{x} = \frac{1}{1 + (x - 1)},$$

and then substitute $-(x - 1)$ for x in the Maclaurin series for $\frac{1}{1-x}$ to obtain

$$\frac{1}{x} = \sum_{n=0}^{\infty} [-(x - 1)]^n = \sum_{n=0}^{\infty} (-1)^n (x - 1)^n.$$

This series is valid for $|x - 1| < 1$.

30. $f(x) = e^{3x}$, $c = -1$

SOLUTION Write

$$e^{3x} = e^{3(x+1)-3} = e^{-3} e^{3(x+1)}.$$

Now, substitute $3(x + 1)$ for x in the Maclaurin series for e^x to obtain

$$e^{3(x+1)} = \sum_{n=0}^{\infty} \frac{(3(x+1))^n}{n!} = \sum_{n=0}^{\infty} \frac{3^n}{n!} (x+1)^n.$$

Thus,

$$e^{3x} = e^{-3} \sum_{n=0}^{\infty} \frac{3^n}{n!} (x+1)^n = \sum_{n=0}^{\infty} \frac{3^n e^{-3}}{n!} (x+1)^n.$$

This series is valid for all x .

31. $f(x) = \frac{1}{1-x}$, $c = 5$

SOLUTION Write

$$\frac{1}{1-x} = \frac{1}{-4 - (x-5)} = -\frac{1}{4} \cdot \frac{1}{1 + \frac{x-5}{4}}.$$

Substituting $-\frac{x-5}{4}$ for x in the Maclaurin series for $\frac{1}{1-x}$ yields

$$\frac{1}{1 + \frac{x-5}{4}} = \sum_{n=0}^{\infty} \left(-\frac{x-5}{4}\right)^n = \sum_{n=0}^{\infty} (-1)^n \frac{(x-5)^n}{4^n}.$$

Thus,

$$\frac{1}{1-x} = -\frac{1}{4} \sum_{n=0}^{\infty} (-1)^n \frac{(x-5)^n}{4^n} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{(x-5)^n}{4^{n+1}}.$$

This series is valid for $\left|\frac{x-5}{4}\right| < 1$, or $|x - 5| < 4$.

32. $f(x) = \sin x$, $c = \frac{\pi}{2}$

SOLUTION Note that the odd derivatives of $\sin x$ are zero at $\frac{\pi}{2}$, and the even derivatives alternate between $+1$ and -1 . Thus the Taylor series centered at $\frac{\pi}{2}$ is

$$\sum_{n=0}^{\infty} (-1)^n \frac{\left(x - \frac{\pi}{2}\right)^{2n}}{(2n)!}$$

33. $f(x) = x^4 + 3x - 1$, $c = 2$

SOLUTION To determine the Taylor series with center $c = 2$, we compute

$$f'(x) = 4x^3 + 3, \quad f''(x) = 12x^2, \quad f'''(x) = 24x,$$

and $f^{(4)}(x) = 24$. All derivatives of order five and higher are zero. Now,

$$f(2) = 21, \quad f'(2) = 35, \quad f''(2) = 48, \quad f'''(2) = 48,$$

and $f^{(4)}(2) = 24$. Therefore, the Taylor series is

$$21 + 35(x-2) + \frac{48}{2}(x-2)^2 + \frac{48}{6}(x-2)^3 + \frac{24}{24}(x-2)^4,$$

or

$$21 + 35(x-2) + 24(x-2)^2 + 8(x-2)^3 + (x-2)^4.$$

34. $f(x) = x^4 + 3x - 1, \quad c = 0$

SOLUTION The function $x^4 + 3x - 1$ is a polynomial in x , hence it is already in the form of a Maclaurin series.

35. $f(x) = \frac{1}{x^2}, \quad c = 4$

SOLUTION We will first find the Taylor series for $\frac{1}{x}$ and then differentiate to obtain the series for $\frac{1}{x^2}$. Write

$$\frac{1}{x} = \frac{1}{4 + (x-4)} = \frac{1}{4} \cdot \frac{1}{1 + \frac{x-4}{4}}.$$

Now substitute $-\frac{x-4}{4}$ for x in the Maclaurin series for $\frac{1}{1-x}$ to obtain

$$\frac{1}{x} = \frac{1}{4} \sum_{n=0}^{\infty} \left(-\frac{x-4}{4}\right)^n = \sum_{n=0}^{\infty} (-1)^n \frac{(x-4)^n}{4^{n+1}}.$$

Differentiating term-by-term yields

$$-\frac{1}{x^2} = \sum_{n=1}^{\infty} (-1)^n n \frac{(x-4)^{n-1}}{4^{n+1}},$$

so that

$$\frac{1}{x^2} = \sum_{n=1}^{\infty} (-1)^{n-1} n \frac{(x-4)^{n-1}}{4^{n+1}} = \sum_{n=0}^{\infty} (-1)^n (n+1) \frac{(x-4)^n}{4^{n+2}}.$$

This series is valid for $\left|\frac{x-4}{4}\right| < 1$, or $|x-4| < 4$.

36. $f(x) = \sqrt{x}, \quad c = 4$

SOLUTION Write

$$\sqrt{x} = \sqrt{4 + (x-4)} = 2\sqrt{1 + \frac{x-4}{4}}.$$

Substituting $\frac{x-4}{4}$ for x in the binomial series with $a = \frac{1}{2}$ yields

$$\sqrt{x} = 2 \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} \left(\frac{x-4}{4}\right)^n = \sum_{n=0}^{\infty} \frac{1}{2^{2n-1}} \binom{\frac{1}{2}}{n} (x-4)^n.$$

This series is valid for $\left|\frac{x-4}{4}\right| < 1$, or $|x-4| < 4$.

37. $f(x) = \frac{1}{1-x^2}, \quad c = 3$

SOLUTION By partial fraction decomposition

$$\frac{1}{1-x^2} = \frac{\frac{1}{2}}{1-x} + \frac{\frac{1}{2}}{1+x},$$

so

$$\frac{1}{1-x^2} = \frac{\frac{1}{2}}{-2 - (x-3)} + \frac{\frac{1}{2}}{4 + (x-3)} = -\frac{1}{4} \cdot \frac{1}{1 + \frac{x-3}{2}} + \frac{1}{8} \cdot \frac{1}{1 + \frac{x-3}{4}}.$$

Substituting $-\frac{x-3}{2}$ for x in the Maclaurin series for $\frac{1}{1-x}$ gives

$$\frac{1}{1 + \frac{x-3}{2}} = \sum_{n=0}^{\infty} \left(-\frac{x-3}{2}\right)^n = \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} (x-3)^n,$$

while substituting $-\frac{x-3}{4}$ for x in the same series gives

$$\frac{1}{1 + \frac{x-3}{4}} = \sum_{n=0}^{\infty} \left(-\frac{x-3}{4}\right)^n = \sum_{n=0}^{\infty} \frac{(-1)^n}{4^n} (x-3)^n.$$

Thus,

$$\begin{aligned} \frac{1}{1-x^2} &= -\frac{1}{4} \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} (x-3)^n + \frac{1}{8} \sum_{n=0}^{\infty} \frac{(-1)^n}{4^n} (x-3)^n = \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{2^{n+2}} (x-3)^n + \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{2n+3}} (x-3)^n \\ &= \sum_{n=0}^{\infty} \left(\frac{(-1)^{n+1}}{2^{n+2}} + \frac{(-1)^n}{2^{2n+3}} \right) (x-3)^n = \sum_{n=0}^{\infty} \frac{(-1)^{n+1} (2^{n+1} - 1)}{2^{2n+3}} (x-3)^n. \end{aligned}$$

This series is valid for $|x-3| < 2$.

38. $f(x) = \frac{1}{3x-2}$, $c = -1$

SOLUTION Write

$$\frac{1}{3x-2} = \frac{1}{-5+3(x+1)} = -\frac{1}{5} \frac{1}{1 - \frac{3(x+1)}{5}},$$

and then substitute $\frac{3(x+1)}{5}$ for x in the Maclaurin series for $\frac{1}{1-x}$ to obtain

$$\frac{1}{1 - \frac{3(x+1)}{5}} = \sum_{n=0}^{\infty} \left(\frac{3(x+1)}{5}\right)^n = \sum_{n=0}^{\infty} \frac{3^n}{5^n} (x+1)^n.$$

Thus,

$$\frac{1}{3x-2} = -\sum_{n=0}^{\infty} \frac{3^n}{5^{n+1}} (x+1)^n.$$

This series is valid for $\left|\frac{3(x+1)}{5}\right| < 1$, or $|x+1| < \frac{5}{3}$.

39. Use the identity $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$ to find the Maclaurin series for $\cos^2 x$.

SOLUTION The Maclaurin series for $\cos 2x$ is

$$\sum_{n=0}^{\infty} (-1)^n \frac{(2x)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{2^{2n} x^{2n}}{(2n)!}$$

so the Maclaurin series for $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$ is

$$\frac{1 + \left(1 + \sum_{n=1}^{\infty} (-1)^n \frac{2^{2n} x^{2n}}{(2n)!}\right)}{2} = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{2^{2n-1} x^{2n}}{(2n)!}$$

40. Show that for $|x| < 1$,

$$\tanh^{-1} x = x + \frac{x^3}{3} + \frac{x^5}{5} + \cdots$$

Hint: Recall that $\frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2}$.

SOLUTION Because

$$\frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2} = \sum_{n=0}^{\infty} (x^2)^n = \sum_{n=0}^{\infty} x^{2n},$$

we have

$$\tanh^{-1} x = C + \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1} = C + x + \frac{x^3}{3} + \frac{x^5}{5} + \cdots.$$

Now, $\tanh^{-1} 0 = 0$, so it follows that $C = 0$, and

$$\tanh^{-1} x = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1} = x + \frac{x^3}{3} + \frac{x^5}{5} + \cdots.$$

41. Use the Maclaurin series for $\ln(1+x)$ and $\ln(1-x)$ to show that

$$\frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) = x + \frac{x^3}{3} + \frac{x^5}{5} + \cdots$$

for $|x| < 1$. What can you conclude by comparing this result with that of Exercise 40?

SOLUTION Using the Maclaurin series for $\ln(1+x)$ and $\ln(1-x)$, we have for $|x| < 1$

$$\begin{aligned} \ln(1+x) - \ln(1-x) &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n - \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} (-x)^n \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n + \sum_{n=1}^{\infty} \frac{x^n}{n} = \sum_{n=1}^{\infty} \frac{1 + (-1)^{n-1}}{n} x^n. \end{aligned}$$

Since $1 + (-1)^{n-1} = 0$ for even n and $1 + (-1)^{n-1} = 2$ for odd n ,

$$\ln(1+x) - \ln(1-x) = \sum_{k=0}^{\infty} \frac{2}{2k+1} x^{2k+1}.$$

Thus,

$$\frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) = \frac{1}{2} (\ln(1+x) - \ln(1-x)) = \frac{1}{2} \sum_{k=0}^{\infty} \frac{2}{2k+1} x^{2k+1} = \sum_{k=0}^{\infty} \frac{x^{2k+1}}{2k+1}.$$

Observe that this is the same series we found in Exercise 40; therefore,

$$\frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) = \tanh^{-1} x.$$

42. Differentiate the Maclaurin series for $\frac{1}{1-x}$ twice to find the Maclaurin series of $\frac{1}{(1-x)^3}$.

SOLUTION Differentiating the Maclaurin series for $\frac{1}{1-x}$ term-by-term, we obtain

$$\frac{1}{(1-x)^2} = \sum_{n=1}^{\infty} n x^{n-1}.$$

Differentiating again then yields

$$\frac{2}{(1-x)^3} = \sum_{n=2}^{\infty} n(n-1) x^{n-2},$$

so that

$$\frac{1}{(1-x)^3} = \sum_{n=2}^{\infty} \frac{n(n-1)}{2} x^{n-2} = \sum_{n=0}^{\infty} \frac{(n+2)(n+1)}{2} x^n.$$

43. Show, by integrating the Maclaurin series for $f(x) = \frac{1}{\sqrt{1-x^2}}$, that for $|x| < 1$,

$$\sin^{-1} x = x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \frac{x^{2n+1}}{2n+1}$$

SOLUTION From Example 10, we know that for $|x| < 1$

$$\frac{1}{\sqrt{1-x^2}} = \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} x^{2n} = 1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} x^{2n},$$

so, for $|x| < 1$,

$$\sin^{-1} x = \int \frac{dx}{\sqrt{1-x^2}} = C + x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \frac{x^{2n+1}}{2n+1}.$$

Since $\sin^{-1} 0 = 0$, we find that $C = 0$. Thus,

$$\sin^{-1} x = x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \frac{x^{2n+1}}{2n+1}.$$

44. Use the first five terms of the Maclaurin series in Exercise 43 to approximate $\sin^{-1} \frac{1}{2}$. Compare the result with the calculator value.

SOLUTION From Exercise 43 we know that for $|x| < 1$,

$$\sin^{-1} x = x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \frac{x^{2n+1}}{2n+1}.$$

The first five terms of the series are:

$$x + \frac{1}{2} \frac{x^3}{3} + \frac{1 \cdot 3}{2 \cdot 4} \frac{x^5}{5} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \frac{x^7}{7} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{2 \cdot 4 \cdot 6 \cdot 8} \frac{x^9}{9} = x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + \frac{35x^9}{1152}$$

Setting $x = \frac{1}{2}$, we obtain the following approximation:

$$\sin^{-1} \frac{1}{2} \approx \frac{1}{2} + \frac{\left(\frac{1}{2}\right)^3}{6} + \frac{3 \cdot \left(\frac{1}{2}\right)^5}{40} + \frac{5 \cdot \left(\frac{1}{2}\right)^7}{112} + \frac{35 \cdot \left(\frac{1}{2}\right)^9}{1152} \approx 0.52358519539.$$

The calculator value is $\sin^{-1} \frac{1}{2} \approx 0.523598775$.

45. How many terms of the Maclaurin series of $f(x) = \ln(1+x)$ are needed to compute $\ln 1.2$ to within an error of at most 0.0001? Make the computation and compare the result with the calculator value.

SOLUTION Substitute $x = 0.2$ into the Maclaurin series for $\ln(1+x)$ to obtain:

$$\ln 1.2 = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(0.2)^n}{n} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{5^n n}.$$

This is an alternating series with $a_n = \frac{1}{n \cdot 5^n}$. Using the error bound for alternating series

$$|\ln 1.2 - S_N| \leq a_{N+1} = \frac{1}{(N+1)5^{N+1}},$$

so we must choose N so that

$$\frac{1}{(N+1)5^{N+1}} < 0.0001 \quad \text{or} \quad (N+1)5^{N+1} > 10,000.$$

For $N = 3$, $(N+1)5^{N+1} = 4 \cdot 5^4 = 2500 < 10,000$, and for $N = 4$, $(N+1)5^{N+1} = 5 \cdot 5^5 = 15,625 > 10,000$; thus, the smallest acceptable value for N is $N = 4$. The corresponding approximation is:

$$S_4 = \sum_{n=1}^4 \frac{(-1)^{n-1}}{5^n \cdot n} = \frac{1}{5} - \frac{1}{5^2 \cdot 2} + \frac{1}{5^3 \cdot 3} - \frac{1}{5^4 \cdot 4} = 0.1822666666.$$

Now, $\ln 1.2 = 0.182321556$, so

$$|\ln 1.2 - S_4| = 5.489 \times 10^{-5} < 0.0001.$$

46. Show that

$$\pi - \frac{\pi^3}{3!} + \frac{\pi^5}{5!} - \frac{\pi^7}{7!} + \cdots$$

converges to zero. How many terms must be computed to get within 0.01 of zero?

SOLUTION Set $x = \pi$ in the Maclaurin series for $\sin x$ to obtain:

$$0 = \sin \pi = \pi - \frac{\pi^3}{3!} + \frac{\pi^5}{5!} - \frac{\pi^7}{7!} + \cdots$$

Using the error bound for an alternating series, we have

$$\left| 0 - \sum_{n=0}^N \frac{(-1)^n \pi^{2n+1}}{(2n+1)!} \right| \leq \frac{\pi^{2N+3}}{(2N+3)!}$$

$N = 4$ is the smallest value for which the error bound is less than 0.01, so five terms are needed.

47. Use the Maclaurin expansion for e^{-t^2} to express the function $F(x) = \int_0^x e^{-t^2} dt$ as an alternating power series in x (Figure 4).

(a) How many terms of the Maclaurin series are needed to approximate the integral for $x = 1$ to within an error of at most 0.001?

(b)  Carry out the computation and check your answer using a computer algebra system.




FIGURE 4 The Maclaurin polynomial $T_{15}(x)$ for $F(t) = \int_0^x e^{-t^2} dt$.

SOLUTION Substituting $-t^2$ for t in the Maclaurin series for e^t yields

$$e^{-t^2} = \sum_{n=0}^{\infty} \frac{(-t^2)^n}{n!} = \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n}}{n!};$$

thus,

$$\int_0^x e^{-t^2} dt = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{n!(2n+1)}.$$

(a) For $x = 1$,

$$\int_0^1 e^{-t^2} dt = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n!(2n+1)}.$$

This is an alternating series with $a_n = \frac{1}{n!(2n+1)}$; therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 e^{-t^2} dt - S_N \right| \leq a_{N+1} = \frac{1}{(N+1)!(2N+3)}.$$

To guarantee the error is at most 0.001, we must choose N so that

$$\frac{1}{(N+1)!(2N+3)} < 0.001 \quad \text{or} \quad (N+1)!(2N+3) > 1000.$$

For $N = 3$, $(N+1)!(2N+3) = 4! \cdot 9 = 216 < 1000$ and for $N = 4$, $(N+1)!(2N+3) = 5! \cdot 11 = 1320 > 1000$; thus, the smallest acceptable value for N is $N = 4$. The corresponding approximation is

$$S_4 = \sum_{n=0}^4 \frac{(-1)^n}{n!(2n+1)} = 1 - \frac{1}{3} + \frac{1}{2! \cdot 5} - \frac{1}{3! \cdot 7} + \frac{1}{4! \cdot 9} = 0.747486772.$$

(b) Using a computer algebra system, we find

$$\int_0^1 e^{-t^2} dt = 0.746824133;$$

therefore

$$\left| \int_0^1 e^{-t^2} dt - S_4 \right| = 6.626 \times 10^{-4} < 10^{-3}.$$

48. Let $F(x) = \int_0^x \frac{\sin t}{t} dt$. Show that

$$F(x) = x - \frac{x^3}{3 \cdot 3!} + \frac{x^5}{5 \cdot 5!} - \frac{x^7}{7 \cdot 7!} + \cdots$$

Evaluate $F(1)$ to three decimal places.

SOLUTION Divide the Maclaurin series for $\sin t$ by t to obtain

$$\frac{\sin t}{t} = \frac{1}{t} \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n}}{(2n+1)!}.$$

Integrating both sides of this equation and using term-by-term integration, we find

$$F(x) = \int_0^x \frac{\sin t}{t} dt = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!(2n+1)} = x - \frac{x^3}{3 \cdot 3!} + \frac{x^5}{5 \cdot 5!} - \frac{x^7}{7 \cdot 7!} + \cdots$$

For $x = 1$,

$$F(1) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)!(2n+1)}.$$

This is an alternating series with $a_n = \frac{1}{(2n+1)!(2n+1)}$; therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 \frac{\sin t}{t} dt - S_N \right| \leq a_{N+1} = \frac{1}{(2N+3)!(2N+3)}.$$

To guarantee the error is at most 0.0005, we must choose N so that

$$\frac{1}{(2N+3)!(2N+3)} < 0.0005 \quad \text{or} \quad (2N+3)!(2N+3) > 2000.$$

For $N = 1$, $(2N+3)!(2N+3) = 5! \cdot 5 = 600 < 2000$ and for $N = 2$, $(2N+3)!(2N+3) = 7! \cdot 7 = 35,280 > 2000$; thus, the smallest acceptable value for N is $N = 2$. The corresponding approximation is

$$S_2 = \sum_{n=0}^2 \frac{(-1)^n}{(2n+1)!(2n+1)} = 1 - \frac{1}{3 \cdot 3!} + \frac{1}{5 \cdot 5!} = 0.946111111.$$

In Exercises 49–52, express the definite integral as an infinite series and find its value to within an error of at most 10^{-4} .

49. $\int_0^1 \cos(x^2) dx$

SOLUTION Substituting x^2 for x in the Maclaurin series for $\cos x$ yields

$$\cos(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n}}{(2n)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n}}{(2n)!};$$

therefore,

$$\int_0^1 \cos(x^2) dx = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+1}}{(2n)!(4n+1)} \Big|_0^1 = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!(4n+1)}.$$

This is an alternating series with $a_n = \frac{1}{(2n)!(4n+1)}$; therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 \cos(x^2) dx - S_N \right| \leq a_{N+1} = \frac{1}{(2N+2)!(4N+5)}.$$

To guarantee the error is at most 0.0001, we must choose N so that

$$\frac{1}{(2N+2)!(4N+5)} < 0.0001 \quad \text{or} \quad (2N+2)!(4N+5) > 10,000.$$

For $N = 2$, $(2N+2)!(4N+5) = 6! \cdot 13 = 9360 < 10,000$ and for $N = 3$, $(2N+2)!(4N+5) = 8! \cdot 17 = 685,440 > 10,000$; thus, the smallest acceptable value for N is $N = 3$. The corresponding approximation is

$$S_3 = \sum_{n=0}^3 \frac{(-1)^n}{(2n)!(4n+1)} = 1 - \frac{1}{5 \cdot 2!} + \frac{1}{9 \cdot 4!} - \frac{1}{13 \cdot 6!} = 0.904522792.$$

50. $\int_0^1 \tan^{-1}(x^2) dx$

SOLUTION Substituting x^2 for x in the Maclaurin series for $\tan^{-1} x$ yields

$$\tan^{-1}(x^2) = \sum_{n=0}^{\infty} (-1)^n \frac{(x^2)^{2n+1}}{2n+1} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1};$$

therefore,

$$\int_0^1 \tan^{-1}(x^2) dx = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+3}}{(2n+1)(4n+3)} \Big|_0^1 = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)(4n+3)}.$$

This is an alternating series with $a_n = \frac{1}{(2n+1)(4n+3)}$; therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 \tan^{-1}(x^2) dx - S_N \right| \leq a_{N+1} = \frac{1}{(2N+3)(4N+7)}.$$

To guarantee the error is at most 0.0001, we must choose N so that

$$\frac{1}{(2N+3)(4N+7)} < 0.0001 \quad \text{or} \quad (2N+3)(4N+7) > 10,000.$$

For $N = 33$, $(2N+3)(4N+7) = (69)(139) = 9591 < 10,000$ and for $N = 34$, $(2N+3)(4N+7) = (71)(143) = 10,153 > 10,000$; thus, the smallest acceptable value for N is $N = 34$. The corresponding approximation is

$$S_{34} = \sum_{n=0}^{34} \frac{(-1)^n}{(2n)!(4n+1)} = 0.297953297.$$

51. $\int_0^1 e^{-x^3} dx$

SOLUTION Substituting $-x^3$ for x in the Maclaurin series for e^x yields

$$e^{-x^3} = \sum_{n=0}^{\infty} \frac{(-x^3)^n}{n!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n}}{n!};$$

therefore,

$$\int_0^1 e^{-x^3} dx = \sum_{n=0}^{\infty} (-1)^n \frac{x^{3n+1}}{n!(3n+1)} \Big|_0^1 = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!(3n+1)}.$$

This is an alternating series with $a_n = \frac{1}{n!(3n+1)}$; therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 e^{-x^3} dx - S_N \right| \leq a_{N+1} = \frac{1}{(N+1)!(3N+4)}.$$

To guarantee the error is at most 0.0001, we must choose N so that

$$\frac{1}{(N+1)!(3N+4)} < 0.0001 \quad \text{or} \quad (N+1)!(3N+4) > 10,000.$$

For $N = 4$, $(N+1)!(3N+4) = 5! \cdot 16 = 1920 < 10,000$ and for $N = 5$, $(N+1)!(3N+4) = 6! \cdot 19 = 13,680 > 10,000$; thus, the smallest acceptable value for N is $N = 5$. The corresponding approximation is

$$S_5 = \sum_{n=0}^5 \frac{(-1)^n}{n!(3n+1)} = 0.807446200.$$

52. $\int_0^1 \frac{dx}{\sqrt{x^4+1}}$

SOLUTION From Example 10, we know that for $|x| < 1$

$$\frac{1}{\sqrt{1-x^2}} = \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} x^{2n} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} x^{2n};$$

therefore,

$$\frac{1}{\sqrt{x^4+1}} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} (-x^2)^{2n} = \sum_{n=0}^{\infty} (-1)^n \frac{(2n)!}{2^{2n}(n!)^2} x^{4n},$$

and

$$\int_0^1 \frac{dx}{\sqrt{x^4+1}} = \sum_{n=0}^{\infty} (-1)^n \frac{(2n)!}{2^{2n}(n!)^2} \frac{x^{4n+1}}{4n+1} \Big|_0^1 = \sum_{n=0}^{\infty} (-1)^n \frac{(2n)!}{2^{2n}(4n+1)(n!)^2}.$$

This is an alternating series with

$$a_n = \frac{(2n)!}{2^{2n}(4n+1)(n!)^2};$$

therefore, the error incurred by using S_N to approximate the value of the definite integral is bounded by

$$\left| \int_0^1 \frac{dx}{\sqrt{x^4+1}} - S_N \right| \leq a_{N+1} = \frac{(2N+2)!}{2^{2N+2}(4N+5)((N+1)!)^2}.$$

To guarantee the error is at most 0.0001, we must choose N so that

$$\frac{(2N+2)!}{2^{2N+2}(4N+5)((N+1)!)^2} < 0.0001.$$

For $N = 124$,

$$\frac{(2N+2)!}{2^{2N+2}(4N+5)((N+1)!)^2} = 0.0001006 > 0.0001,$$

and for $N = 125$,

$$\frac{(2N+2)!}{2^{2N+2}(4N+5)((N+1)!)^2} = 0.00009943 < 0.0001,$$

thus, the smallest acceptable value for N is $N = 125$. The corresponding approximation is

$$S_{125} = \sum_{n=0}^{125} (-1)^n \frac{(2n)!}{2^{2n}(4n+1)(n!)^2} = 0.926987328.$$

In Exercises 53–56, express the integral as an infinite series.

53. $\int_0^x \frac{1 - \cos(t)}{t} dt$, for all x

SOLUTION The Maclaurin series for $\cos t$ is

$$\cos t = \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n}}{(2n)!} = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n}}{(2n)!},$$

so

$$1 - \cos t = - \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n}}{(2n)!} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n}}{(2n)!},$$

and

$$\frac{1 - \cos t}{t} = \frac{1}{t} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n}}{(2n)!} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n-1}}{(2n)!}.$$

Thus,

$$\int_0^x \frac{1 - \cos(t)}{t} dt = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n}}{(2n)!2n} \Big|_0^x = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n}}{(2n)!2n}.$$

54. $\int_0^x \frac{t - \sin t}{t} dt$, for all x

SOLUTION The Maclaurin series for $\sin t$ is

$$\sin t = \sum_{n=0}^{\infty} (-1)^n \frac{t^{2n+1}}{(2n+1)!} = t + \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n+1}}{(2n+1)!},$$

so

$$t - \sin t = - \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n+1}}{(2n+1)!} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n+1}}{(2n+1)!},$$

and

$$\frac{t - \sin t}{t} = \frac{1}{t} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n+1}}{(2n+1)!} = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n}}{(2n+1)!}.$$

Thus,

$$\int_0^x \frac{t - \sin(t)}{t} dt = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{t^{2n+1}}{(2n+1)!(2n+1)} \Big|_0^x = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n+1}}{(2n+1)!(2n+1)}.$$

55. $\int_0^x \ln(1+t^2) dt$, for $|x| < 1$

SOLUTION Substituting t^2 for t in the Maclaurin series for $\ln(1+t)$ yields

$$\ln(1+t^2) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{(t^2)^n}{n} = \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n}}{n}.$$

Thus,

$$\int_0^x \ln(1+t^2) dt = \sum_{n=1}^{\infty} (-1)^n \frac{t^{2n+1}}{n(2n+1)} \Big|_0^x = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n+1}}{n(2n+1)}.$$

56. $\int_0^x \frac{dt}{\sqrt{1-t^4}}$, for $|x| < 1$

SOLUTION From Example 10, we know that for $|t| < 1$

$$\frac{1}{\sqrt{1-t^2}} = \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} t^{2n} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n} (n!)^2} t^{2n},$$

therefore,

$$\frac{1}{\sqrt{1-t^4}} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n} (n!)^2} (t^2)^{2n} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n} (n!)^2} t^{4n},$$

and

$$\int_0^x \frac{dt}{\sqrt{1-t^4}} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} \frac{t^{4n+1}}{4n+1} \Big|_0^x = \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} \frac{x^{4n+1}}{4n+1}.$$

57. Which function has Maclaurin series $\sum_{n=0}^{\infty} (-1)^n 2^n x^n$?

SOLUTION We recognize that

$$\sum_{n=0}^{\infty} (-1)^n 2^n x^n = \sum_{n=0}^{\infty} (-2x)^n$$

is the Maclaurin series for $\frac{1}{1-x}$ with x replaced by $-2x$. Therefore,

$$\sum_{n=0}^{\infty} (-1)^n 2^n x^n = \frac{1}{1-(-2x)} = \frac{1}{1+2x}.$$

58. Which function has Maclaurin series

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{3^{k+1}} (x-3)^k?$$

For which values of x is the expansion valid?

SOLUTION Write the series as

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{3^{k+1}} (x-3)^k = \frac{1}{3} \sum_{k=0}^{\infty} \left(-\frac{x-3}{3} \right)^k,$$

which we recognize as $\frac{1}{3}$ times the Maclaurin series for $\frac{1}{1-x}$ with x replaced by $-\frac{x-3}{3}$. Therefore,

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{3^{k+1}} (x-3)^k = \frac{1}{3} \cdot \frac{1}{1+\frac{x-3}{3}} = \frac{1}{3+x-3} = \frac{1}{x}.$$

The series is valid for $\left| \frac{x-3}{3} \right| < 1$, or $|x-3| < 3$.

In Exercises 59–62, use Theorem 2 to prove that the $f(x)$ is represented by its Maclaurin series on the interval I .

59. $f(x) = \sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{3}\right)$,

SOLUTION All derivatives of $f(x)$ consist of sin or cos applied to each of $x/2$ and $x/3$ and added together, so each summand is bounded by 1. Thus $|f^{(n)}(x)| \leq 2$ for all n and x . By Theorem 2, $f(x)$ is represented by its Taylor series for every x .

60. $f(x) = e^{-x}$,

SOLUTION For any c , choose any $R > 0$ and consider the interval $(c-R, c+R)$. For $f(x) = e^{-x}$, we have

$$|f^{(n)}(x)| = |(-1)^n e^{-x}| = e^{-x}$$

and on $(c-R, c+R)$, e^{-x} is bounded above by $e^{-(c-R)} = e^{R-c}$. Thus all derivatives of $f(x)$ are bounded by e^{R-c} for any $x \in (c-R, c+R)$, so by Theorem 2, $f(x)$ is represented by its Taylor series centered at c .

61. $f(x) = \sinh x$,

SOLUTION By definition, $\sinh x = \frac{1}{2}(e^x - e^{-x})$, so if both e^x and e^{-x} are represented by their Taylor series centered at c , then so is $\sinh x$. But the previous exercise shows that e^{-x} is so represented, and the text shows that e^x is.

62. $f(x) = (1+x)^{100}$

SOLUTION $f(x)$ is a polynomial, so it is equal to its Taylor series and thus is obviously represented by its Taylor series.

In Exercises 63–66, find the functions with the following Maclaurin series (refer to Table 1 on page 599).

$$63. 1 + x^3 + \frac{x^6}{2!} + \frac{x^9}{3!} + \frac{x^{12}}{4!} + \cdots$$

SOLUTION We recognize

$$1 + x^3 + \frac{x^6}{2!} + \frac{x^9}{3!} + \frac{x^{12}}{4!} + \cdots = \sum_{n=0}^{\infty} \frac{x^{3n}}{n!} = \sum_{n=0}^{\infty} \frac{(x^3)^n}{n!}$$

as the Maclaurin series for e^x with x replaced by x^3 . Therefore,

$$1 + x^3 + \frac{x^6}{2!} + \frac{x^9}{3!} + \frac{x^{12}}{4!} + \cdots = e^{x^3}.$$

$$64. 1 - 4x + 4^2x^2 - 4^3x^3 + 4^4x^4 - 4^5x^5 + \cdots$$

SOLUTION We recognize

$$1 - 4x + 4^2x^2 - 4^3x^3 + 4^4x^4 - 4^5x^5 + \cdots = \sum_{n=0}^{\infty} (-4x)^n$$

as the Maclaurin series for $\frac{1}{1-x}$ with x replaced by $-4x$. Therefore,

$$1 - 4x + 4^2x^2 - 4^3x^3 + 4^4x^4 - 4^5x^5 + \cdots = \frac{1}{1 - (-4x)} = \frac{1}{1 + 4x}.$$

$$65. 1 - \frac{5^3x^3}{3!} + \frac{5^5x^5}{5!} - \frac{5^7x^7}{7!} + \cdots$$

SOLUTION Note

$$\begin{aligned} 1 - \frac{5^3x^3}{3!} + \frac{5^5x^5}{5!} - \frac{5^7x^7}{7!} + \cdots &= 1 - 5x + \left(5x - \frac{5^3x^3}{3!} + \frac{5^5x^5}{5!} - \frac{5^7x^7}{7!} + \cdots \right) \\ &= 1 - 5x + \sum_{n=0}^{\infty} (-1)^n \frac{(5x)^{2n+1}}{(2n+1)!}. \end{aligned}$$

The series is the Maclaurin series for $\sin x$ with x replaced by $5x$, so

$$1 - \frac{5^3x^3}{3!} + \frac{5^5x^5}{5!} - \frac{5^7x^7}{7!} + \cdots = 1 - 5x + \sin(5x).$$

$$66. x^4 - \frac{x^{12}}{3} + \frac{x^{20}}{5} - \frac{x^{28}}{7} + \cdots$$

SOLUTION We recognize

$$x^4 - \frac{x^{12}}{3} + \frac{x^{20}}{5} - \frac{x^{28}}{7} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{(x^4)^{2n+1}}{2n+1}$$

as the Maclaurin series for $\tan^{-1} x$ with x replaced by x^4 . Therefore,

$$x^4 - \frac{x^{12}}{3} + \frac{x^{20}}{5} - \frac{x^{28}}{7} + \cdots = \tan^{-1}(x^4).$$

In Exercises 67 and 68, let

$$f(x) = \frac{1}{(1-x)(1-2x)}$$

67. Find the Maclaurin series of $f(x)$ using the identity

$$f(x) = \frac{2}{1-2x} - \frac{1}{1-x}$$

SOLUTION Substituting $2x$ for x in the Maclaurin series for $\frac{1}{1-x}$ gives

$$\frac{1}{1-2x} = \sum_{n=0}^{\infty} (2x)^n = \sum_{n=0}^{\infty} 2^n x^n$$

which is valid for $|2x| < 1$, or $|x| < \frac{1}{2}$. Because the Maclaurin series for $\frac{1}{1-x}$ is valid for $|x| < 1$, the two series together are valid for $|x| < \frac{1}{2}$. Thus, for $|x| < \frac{1}{2}$,

$$\begin{aligned} \frac{1}{(1-2x)(1-x)} &= \frac{2}{1-2x} - \frac{1}{1-x} = 2 \sum_{n=0}^{\infty} 2^n x^n - \sum_{n=0}^{\infty} x^n \\ &= \sum_{n=0}^{\infty} 2^{n+1} x^n - \sum_{n=0}^{\infty} x^n = \sum_{n=0}^{\infty} (2^{n+1} - 1) x^n. \end{aligned}$$

68. Find the Taylor series for $f(x)$ at $c = 2$. *Hint:* Rewrite the identity of Exercise 67 as

$$f(x) = \frac{2}{-3-2(x-2)} - \frac{1}{-1-(x-2)}$$

SOLUTION Using the given identity,

$$f(x) = \frac{2}{-3-2(x-2)} - \frac{1}{-1-(x-2)} = -\frac{2}{3} \frac{1}{1+\frac{2}{3}(x-2)} + \frac{1}{1+(x-2)}.$$

Substituting $-\frac{2}{3}(x-2)$ for x in the Maclaurin series for $\frac{1}{1-x}$ yields

$$\frac{1}{1+\frac{2}{3}(x-2)} = \sum_{n=0}^{\infty} (-1)^n \left(\frac{2}{3}\right)^n (x-2)^n,$$

and substituting $-(x-2)$ for x in the same Maclaurin series yields

$$\frac{1}{1+(x-2)} = \sum_{n=0}^{\infty} (-1)^n (x-2)^n.$$

The first series is valid for $\left|-\frac{2}{3}(x-2)\right| < 1$, or $|x-2| < \frac{3}{2}$, and the second series is valid for $|x-2| < 1$; therefore, the two series together are valid for $|x-2| < 1$. Finally, for $|x-2| < 1$,

$$f(x) = -\frac{2}{3} \sum_{n=0}^{\infty} (-1)^n \left(\frac{2}{3}\right)^n (x-2)^n + \sum_{n=0}^{\infty} (-1)^n (x-2)^n = \sum_{n=0}^{\infty} (-1)^n \left[1 - \left(\frac{2}{3}\right)^{n+1}\right] (x-2)^n.$$

69. When a voltage V is applied to a series circuit consisting of a resistor R and an inductor L , the current at time t is

$$I(t) = \left(\frac{V}{R}\right) (1 - e^{-Rt/L})$$

Expand $I(t)$ in a Maclaurin series. Show that $I(t) \approx \frac{Vt}{L}$ for small t .

SOLUTION Substituting $-\frac{Rt}{L}$ for t in the Maclaurin series for e^t gives

$$e^{-Rt/L} = \sum_{n=0}^{\infty} \frac{\left(-\frac{Rt}{L}\right)^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left(\frac{R}{L}\right)^n t^n = 1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \left(\frac{R}{L}\right)^n t^n$$

Thus,

$$1 - e^{-Rt/L} = 1 - \left(1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \left(\frac{R}{L}\right)^n t^n\right) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n!} \left(\frac{Rt}{L}\right)^n,$$

and

$$I(t) = \frac{V}{R} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n!} \left(\frac{Rt}{L}\right)^n = \frac{Vt}{L} + \frac{V}{R} \sum_{n=2}^{\infty} \frac{(-1)^{n+1}}{n!} \left(\frac{Rt}{L}\right)^n.$$

If t is small, then we can approximate $I(t)$ by the first (linear) term, and ignore terms with higher powers of t ; then we find

$$V(t) \approx \frac{Vt}{L}.$$

70. Use the result of Exercise 69 and your knowledge of alternating series to show that

$$\frac{Vt}{L} \left(1 - \frac{R}{2L}t\right) \leq I(t) \leq \frac{Vt}{L} \quad (\text{for all } t)$$

SOLUTION Since the series for $I(t)$ is an alternating series, we know that the true value lies between any two successive partial sums. Since the term for $n = 2$ is negative, we have

$$S_2 \leq I(t) \leq S_1 \quad \text{for all } t$$

Clearly $S_1 = \frac{Vt}{L}$, and

$$S_2 = \frac{Vt}{L} + \frac{V}{R} \left(\frac{-1}{n!} \cdot \frac{R^2 t^2}{L^2} \right) = \frac{Vt}{L} - \frac{VR^2 t^2}{2RL^2} = \frac{Vt}{L} \left(1 - \frac{R}{2L}t\right)$$

71. Find the Maclaurin series for $f(x) = \cos(x^3)$ and use it to determine $f^{(6)}(0)$.

SOLUTION The Maclaurin series for $\cos x$ is

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

Substituting x^3 for x gives

$$\cos(x^3) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{6n}}{(2n)!}$$

Now, the coefficient of x^6 in this series is

$$-\frac{1}{2!} = -\frac{1}{2} = \frac{f^{(6)}(0)}{6!}$$

so

$$f^{(6)}(0) = -\frac{6!}{2} = -360$$

72. Find $f^{(7)}(0)$ and $f^{(8)}(0)$ for $f(x) = \tan^{-1} x$ using the Maclaurin series.

SOLUTION The Maclaurin series for $f(x) = \tan^{-1} x$ is:

$$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}.$$


The coefficient of x^7 in this series is

$$\frac{(-1)^3}{7} = -\frac{1}{7} = \frac{f^{(7)}(0)}{7!},$$

so

$$f^{(7)}(0) = -\frac{7!}{7} = -6! = -720.$$

The coefficient of x^8 is 0, so $f^{(8)}(0) = 0$.

73.  Use substitution to find the first three terms of the Maclaurin series for $f(x) = e^{x^{20}}$. How does the result show that $f^{(k)}(0) = 0$ for $1 \leq k \leq 19$?

SOLUTION Substituting x^{20} for x in the Maclaurin series for e^x yields

$$e^{x^{20}} = \sum_{n=0}^{\infty} \frac{(x^{20})^n}{n!} = \sum_{n=0}^{\infty} \frac{x^{20n}}{n!};$$

the first three terms in the series are then

$$1 + x^{20} + \frac{1}{2}x^{40}.$$

Recall that the coefficient of x^k in the Maclaurin series for f is $\frac{f^{(k)}(0)}{k!}$. For $1 \leq k \leq 19$, the coefficient of x^k in the Maclaurin series for $f(x) = e^{x^{20}}$ is zero; it therefore follows that

$$\frac{f^{(k)}(0)}{k!} = 0 \quad \text{or} \quad f^{(k)}(0) = 0$$

for $1 \leq k \leq 19$.

74. Use the binomial series to find $f^{(8)}(0)$ for $f(x) = \sqrt{1-x^2}$.

SOLUTION We obtain the Maclaurin series for $f(x) = \sqrt{1-x^2}$ by substituting $-x^2$ for x in the binomial series with $a = \frac{1}{2}$. This gives

$$\sqrt{1-x^2} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} (-x^2)^n = \sum_{n=0}^{\infty} (-1)^n \binom{\frac{1}{2}}{n} x^{2n}.$$

The coefficient of x^8 is

$$(-1)^4 \binom{\frac{1}{2}}{4} = \frac{\frac{1}{2} \left(\frac{1}{2} - 1\right) \left(\frac{1}{2} - 2\right) \left(\frac{1}{2} - 3\right)}{4!} = -\frac{15}{16 \cdot 4!} = \frac{f^{(8)}(0)}{8!},$$

so


$$f^{(8)}(0) = \frac{-15 \cdot 8!}{16 \cdot 4!} = -1575.$$

75. Does the Maclaurin series for $f(x) = (1+x)^{3/4}$ converge to $f(x)$ at $x = 2$? Give numerical evidence to support your answer.

SOLUTION The Taylor series for $f(x) = (1+x)^{3/4}$ converges to $f(x)$ for $|x| < 1$; because $x = 2$ is not contained on this interval, the series does not converge to $f(x)$ at $x = 2$. The graph below displays


$$S_N = \sum_{n=0}^N \binom{\frac{3}{4}}{n} 2^n$$

for $0 \leq N \leq 14$. The divergent nature of the sequence of partial sums is clear.



76.  Explain the steps required to verify that the Maclaurin series for $f(x) = e^x$ converges to $f(x)$ for all x .

SOLUTION To show that the Maclaurin series for e^x converges to e^x for all x , we show that for any real number c , the Maclaurin series converges to e^x on an interval containing c . To do this, it suffices to show that for any interval $I = (-R, R)$, the Maclaurin series for e^x converges to e^x on I , since each real number is contained in some such interval. By Theorem 2, it suffices to show that there is a number K that bounds all derivatives of e^x for all numbers in the interval $(-R, R)$. But each derivative of e^x is also e^x , so it suffices to show that there is a number K that bounds e^x for all $x \in (-R, R)$. But e^x is an increasing function, so that $e^x < e^R$ for all $x \in (-R, R)$. Thus $K = e^R$ is the bound we want. Theorem 2 then assures us that the Maclaurin series for e^x converges to e^x on I .

77.  Let $f(x) = \sqrt{1+x}$.

(a) Use a graphing calculator to compare the graph of f with the graphs of the first five Taylor polynomials for f . What do they suggest about the interval of convergence of the Taylor series?

(b) Investigate numerically whether or not the Taylor expansion for f is valid for $x = 1$ and $x = -1$.

SOLUTION

(a) The five first terms of the Binomial series with $a = \frac{1}{2}$ are

$$\begin{aligned}\sqrt{1+x} &= 1 + \frac{1}{2}x + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)}{2!}x^2 + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)\left(\frac{1}{2}-2\right)}{3!}x^3 + \frac{\frac{1}{2}\left(\frac{1}{2}-1\right)\left(\frac{1}{2}-2\right)\left(\frac{1}{2}-3\right)}{4!}x^4 + \dots \\ &= 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{9}{4}x^3 - \frac{45}{2}x^4 + \dots\end{aligned}$$

Therefore, the first five Taylor polynomials are

$$T_0(x) = 1;$$


$$T_1(x) = 1 + \frac{1}{2}x;$$

$$T_2(x) = 1 + \frac{1}{2}x - \frac{1}{8}x^2;$$

$$T_3(x) = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{8}x^3;$$

$$T_4(x) = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{8}x^3 - \frac{5}{128}x^4.$$

The figure displays the graphs of these Taylor polynomials, along with the graph of the function $f(x) = \sqrt{1+x}$, which is shown in red.



The graphs suggest that the interval of convergence for the Taylor series is $-1 < x < 1$.

(b) Using a computer algebra system to calculate $S_N = \sum_{n=0}^N \left(\frac{1}{2}\right)^n x^n$ for $x = 1$ we find

$$S_{10} = 1.409931183, \quad S_{100} = 1.414073048, \quad S_{1000} = 1.414209104,$$

which appears to be converging to $\sqrt{2}$ as expected. At $x = -1$ we calculate $S_N = \sum_{n=0}^N \left(\frac{1}{2}\right)^n \cdot (-1)^n$, and find

$$S_{10} = 0.176197052, \quad S_{100} = 0.056348479, \quad S_{1000} = 0.017839011,$$

which appears to be converging to zero, though slowly.

78. Use the first five terms of the Maclaurin series for the elliptic function $E(k)$ to estimate the period T of a 1-meter pendulum released at an angle $\theta = \frac{\pi}{4}$ (see Example 11).

SOLUTION The period T of a pendulum of length L released from an angle θ is

$$T = 4\sqrt{\frac{L}{g}}E(k),$$

where $g \approx 9.8 \text{ m/s}^2$ is the acceleration due to gravity, $E(k)$ is the elliptic function of the first kind and $k = \sin \frac{\theta}{2}$. From Example 11, we know that

$$E(k) = \frac{\pi}{2} \sum_{n=0}^{\infty} \left(\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \right)^2 k^{2n}.$$

With $\theta = \frac{\pi}{4}$,

$$k = \sin \frac{\pi}{8} = \frac{\sqrt{2-\sqrt{2}}}{2},$$

and using the first five terms of the series for $E(k)$, we find

$$\begin{aligned} E\left(\sin \frac{\pi}{8}\right) &\approx \frac{\pi}{2} \left(1 + \left(\frac{1}{2}\right)^2 \sin^2 \frac{\pi}{8} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \sin^4 \frac{\pi}{8} + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \sin^6 \frac{\pi}{8} + \left(\frac{1 \cdot 3 \cdot 5 \cdot 7}{2 \cdot 4 \cdot 6 \cdot 8}\right)^2 \sin^8 \frac{\pi}{8}\right) \\ &= 1.633578996 \end{aligned}$$

Therefore,

$$T \approx 4\sqrt{\frac{1}{9.8}} \cdot 1.633578996 = 2.09 \text{ seconds.}$$

79. Use Example 11 and the approximation $\sin x \approx x$ to show that the period T of a pendulum released at an angle θ has the following second-order approximation:

$$T \approx 2\pi \sqrt{\frac{L}{g}} \left(1 + \frac{\theta^2}{16}\right)$$

SOLUTION The period T of a pendulum of length L released from an angle θ is

$$T = 4\sqrt{\frac{L}{g}} E(k),$$

where $g \approx 9.8 \text{ m/s}^2$ is the acceleration due to gravity, $E(k)$ is the elliptic function of the first kind and $k = \sin \frac{\theta}{2}$. From Example 11, we know that

$$E(k) = \frac{\pi}{2} \sum_{n=0}^{\infty} \left(\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)}\right)^2 k^{2n}.$$

Using the approximation $\sin x \approx x$, we have

$$k = \sin \frac{\theta}{2} \approx \frac{\theta}{2};$$

moreover, using the first two terms of the series for $E(k)$, we find

$$E(k) \approx \frac{\pi}{2} \left[1 + \left(\frac{1}{2}\right)^2 \left(\frac{\theta}{2}\right)^2\right] = \frac{\pi}{2} \left(1 + \frac{\theta^2}{16}\right).$$

Therefore,

$$T = 4\sqrt{\frac{L}{g}} E(k) \approx 2\pi \sqrt{\frac{L}{g}} \left(1 + \frac{\theta^2}{16}\right).$$

In Exercises 80–83, find the Maclaurin series of the function and use it to calculate the limit.

$$\mathbf{80.} \quad \lim_{x \rightarrow 0} \frac{\cos x - 1 + \frac{x^2}{2}}{x^4}$$

SOLUTION Using the Maclaurin series for $\cos x$, we find

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2} + \frac{x^4}{24} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}.$$

Thus,

$$\cos x - 1 + \frac{x^2}{2} = \frac{x^4}{24} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

and

$$\frac{\cos x - 1 + \frac{x^2}{2}}{x^4} = \frac{1}{24} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n-4}}{(2n)!}$$

Note that the radius of convergence for this series is infinite, and recall from the previous section that a convergent power series is continuous within its radius of convergence. Thus to calculate the limit of this power series as $x \rightarrow 0$ it suffices to evaluate it at $x = 0$:

$$\lim_{x \rightarrow 0} \frac{\cos x - 1 + \frac{x^2}{2}}{x^4} = \lim_{x \rightarrow 0} \left(\frac{1}{24} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n-4}}{(2n)!} \right) = \frac{1}{24} + 0 = \frac{1}{24}.$$

$$81. \lim_{x \rightarrow 0} \frac{\sin x - x + \frac{x^3}{6}}{x^5}$$

SOLUTION Using the Maclaurin series for $\sin x$, we find

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{6} + \frac{x^5}{120} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}.$$

Thus,

$$\sin x - x + \frac{x^3}{6} = \frac{x^5}{120} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

and

$$\frac{\sin x - x + \frac{x^3}{6}}{x^5} = \frac{1}{120} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n-4}}{(2n+1)!}$$

Note that the radius of convergence for this series is infinite, and recall from the previous section that a convergent power series is continuous within its radius of convergence. Thus to calculate the limit of this power series as $x \rightarrow 0$ it suffices to evaluate it at $x = 0$:

$$\lim_{x \rightarrow 0} \frac{\sin x - x + \frac{x^3}{6}}{x^5} = \lim_{x \rightarrow 0} \left(\frac{1}{120} + \sum_{n=3}^{\infty} (-1)^n \frac{x^{2n-4}}{(2n+1)!} \right) = \frac{1}{120} + 0 = \frac{1}{120}$$

$$82. \lim_{x \rightarrow 0} \frac{\tan^{-1} x - x \cos x - \frac{1}{6}x^3}{x^5}$$

SOLUTION Start with the Maclaurin series for $\tan^{-1} x$ and $\cos x$:

$$\tan^{-1} x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad \cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

Then

$$x \cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n)!}$$

so that

$$\begin{aligned} \tan^{-1} x - x \cos x &= \sum_{n=0}^{\infty} (-1)^n \left(\frac{1}{2n+1} - \frac{1}{(2n)!} \right) x^{2n+1} \\ &= \frac{1}{6}x^3 + \frac{19}{120}x^5 + \sum_{n=3}^{\infty} (-1)^n \left(\frac{1}{2n+1} - \frac{1}{(2n)!} \right) x^{2n+1} \end{aligned}$$

and

$$\frac{\tan^{-1} x - x \cos x - \frac{1}{6}x^3}{x^5} = \frac{19}{120} + \sum_{n=3}^{\infty} (-1)^n \left(\frac{1}{2n+1} - \frac{1}{(2n)!} \right) x^{2n-4}$$

Since the radius of convergence of the series for $\tan^{-1} x$ is 1 and that of $\cos x$ is infinite, the radius of convergence of this series is 1. Recall from the previous section that a convergent power series is continuous within its radius of convergence. Thus to calculate the limit of this power series as $x \rightarrow 0$ it suffices to evaluate it at $x = 0$:

$$\lim_{x \rightarrow 0} \frac{\tan^{-1} x - x \cos x - \frac{1}{6}x^3}{x^5} = \lim_{x \rightarrow 0} \left(\frac{19}{120} + \sum_{n=3}^{\infty} (-1)^n \left(\frac{1}{2n+1} - \frac{1}{(2n)!} \right) x^{2n-4} \right) = \frac{19}{120} + 0 = \frac{19}{120}$$

83. $\lim_{x \rightarrow 0} \left(\frac{\sin(x^2)}{x^4} - \frac{\cos x}{x^2} \right)$

SOLUTION We start with

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad \cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

so that

$$\begin{aligned} \frac{\sin(x^2)}{x^4} &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{(2n+1)!x^4} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n-2}}{(2n+1)!} \\ \frac{\cos x}{x^2} &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n-2}}{(2n)!} \end{aligned}$$

Expanding the first few terms gives

$$\begin{aligned} \frac{\sin(x^2)}{x^4} &= \frac{1}{x^2} - \sum_{n=1}^{\infty} (-1)^n \frac{x^{4n-2}}{(2n+1)!} \\ \frac{\cos x}{x^2} &= \frac{1}{x^2} - \frac{1}{2} + \sum_{n=2}^{\infty} (-1)^n \frac{x^{2n-2}}{(2n)!} \end{aligned}$$

so that

$$\frac{\sin(x^2)}{x^4} - \frac{\cos x}{x^2} = \frac{1}{2} - \sum_{n=1}^{\infty} (-1)^n \frac{x^{4n-2}}{(2n+1)!} - \sum_{n=2}^{\infty} (-1)^n \frac{x^{2n-2}}{(2n)!}$$

Note that all terms under the summation signs have positive powers of x . Now, the radius of convergence of the series for both \sin and \cos is infinite, so the radius of convergence of this series is infinite. Recall from the previous section that a convergent power series is continuous within its radius of convergence. Thus to calculate the limit of this power series as $x \rightarrow 0$ it suffices to evaluate it at $x = 0$:

$$\lim_{x \rightarrow 0} \left(\frac{\sin(x^2)}{x^4} - \frac{\cos x}{x^2} \right) = \lim_{x \rightarrow 0} \left(\frac{1}{2} - \sum_{n=1}^{\infty} (-1)^n \frac{x^{4n-2}}{(2n+1)!} - \sum_{n=2}^{\infty} (-1)^n \frac{x^{2n-2}}{(2n)!} \right) = \frac{1}{2} + 0 = \frac{1}{2}$$

Further Insights and Challenges

84. In this exercise we show that the Maclaurin expansion of $f(x) = \ln(1+x)$ is valid for $x = 1$.

(a) Show that for all $x \neq -1$,

$$\frac{1}{1+x} = \sum_{n=0}^N (-1)^n x^n + \frac{(-1)^{N+1} x^{N+1}}{1+x}$$

(b) Integrate from 0 to 1 to obtain

$$\ln 2 = \sum_{n=1}^N \frac{(-1)^{n-1}}{n} + (-1)^{N+1} \int_0^1 \frac{x^{N+1} dx}{1+x}$$

(c) Verify that the integral on the right tends to zero as $N \rightarrow \infty$ by showing that it is smaller than $\int_0^1 x^{N+1} dx$.

(d) Prove the formula

$$\ln 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots$$

SOLUTION

(a) Substituting $-x$ for x in the Maclaurin series for $\frac{1}{1-x}$ yields

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n.$$

Now, rewrite the series as

$$\sum_{n=0}^N (-1)^n x^n + \sum_{n=N+1}^{\infty} (-1)^n x^n,$$

and use the formula for the sum of a geometric series on the second term to obtain

$$\frac{1}{1+x} = \sum_{n=0}^N (-1)^n x^n + \frac{(-1)^{N+1} x^{N+1}}{1+x}.$$

(b) Integrate the equation derived in part (a) from 0 to 1 to obtain

$$\ln(1+x) \Big|_0^1 = \sum_{n=0}^N (-1)^n \frac{x^{n+1}}{n+1} \Big|_0^1 + (-1)^{N+1} \int_0^1 \frac{x^{N+1}}{1+x} dx,$$

or

$$\ln 2 = \sum_{n=0}^N \frac{(-1)^n}{n+1} + (-1)^{N+1} \int_0^1 \frac{x^{N+1}}{1+x} dx = \sum_{n=1}^{N+1} \frac{(-1)^{n-1}}{n} + (-1)^{N+1} \int_0^1 \frac{x^{N+1}}{1+x} dx.$$

(c) For $0 < x < 1$,

$$0 \leq \frac{x^{N+1}}{1+x} \leq x^{N+1} \quad \text{so} \quad 0 \leq \int_0^1 \frac{x^{N+1}}{1+x} dx \leq \int_0^1 x^{N+1} dx.$$

Now,

$$\int_0^1 x^{N+1} dx = \frac{x^{N+2}}{N+2} \Big|_0^1 = \frac{1}{N+2} \rightarrow 0 \text{ as } N \rightarrow \infty.$$

Thus, by the Squeeze Theorem,

$$\lim_{N \rightarrow \infty} \int_0^1 \frac{x^{N+1}}{1+x} dx = 0.$$

(d) Taking the limit as $N \rightarrow \infty$ of the equation derived in part (b) and using the result from part (c), we find

$$\ln 2 = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

85. Let $g(t) = \frac{1}{1+t^2} - \frac{t}{1+t^2}$.

(a) Show that $\int_0^1 g(t) dt = \frac{\pi}{4} - \frac{1}{2} \ln 2$.

(b) Show that $g(t) = 1 - t - t^2 + t^3 - t^4 + t^5 + \dots$.

(c) Evaluate $S = 1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} - \frac{1}{6} + \dots$.

SOLUTION

(a)

$$\int_0^1 g(t) dt = \left(\tan^{-1} t - \frac{1}{2} \ln(t^2 + 1) \right) \Big|_0^1 = \tan^{-1} 1 - \frac{1}{2} \ln 2 = \frac{\pi}{4} - \frac{1}{2} \ln 2$$

(b) Start with the Taylor series for $\frac{1}{1+t}$:

$$\frac{1}{1+t} = \sum_{n=0}^{\infty} (-1)^n t^n$$

and substitute t^2 for t to get

$$\frac{1}{1+t^2} = \sum_{n=0}^{\infty} (-1)^n t^{2n} = 1 - t^2 + t^4 - t^6 + \dots$$

so that

$$\frac{t}{1+t^2} = \sum_{n=0}^{\infty} (-1)^n t^{2n+1} = t - t^3 + t^5 - t^7 + \dots$$

Finally,

$$g(t) = \frac{1}{1+t^2} - \frac{t}{1+t^2} = 1 - t - t^2 + t^3 + t^4 - t^5 - t^6 + t^7 + \dots$$

(c) We have

$$\int g(t) dt = \int (1 - t - t^2 + t^3 + t^4 - t^5 - \dots) dt = t - \frac{1}{2}t^2 - \frac{1}{3}t^3 + \frac{1}{4}t^4 + \frac{1}{5}t^5 - \frac{1}{6}t^6 - \dots + C$$

The radius of convergence of the series for $g(t)$ is 1, so the radius of convergence of this series is also 1. However, this series converges at the right endpoint, $t = 1$, since

$$\left(1 - \frac{1}{2}\right) - \left(\frac{1}{3} - \frac{1}{4}\right) + \left(\frac{1}{5} - \frac{1}{6}\right) - \dots$$

is an alternating series with general term decreasing to zero. Thus by part (a),

$$1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} - \dots = \frac{\pi}{4} - \frac{1}{2} \ln 2$$

In Exercises 86 and 87, we investigate the convergence of the binomial series

$$T_a(x) = \sum_{n=0}^{\infty} \binom{a}{n} x^n$$

86. Prove that $T_a(x)$ has radius of convergence $R = 1$ if a is not a whole number. What is the radius of convergence if a is a whole number?

SOLUTION Suppose that a is not a whole number. Then

$$\binom{a}{n} = \frac{a(a-1)\cdots(a-n+1)}{n!}$$

is never zero. Moreover,

$$\left| \frac{\binom{a}{n+1}}{\binom{a}{n}} \right| = \left| \frac{a(a-1)\cdots(a-n+1)(a-n)}{(n+1)!} \cdot \frac{n!}{a(a-1)\cdots(a-n+1)} \right| = \left| \frac{a-n}{n+1} \right|,$$

so, by the formula for the radius of convergence

$$r = \lim_{n \rightarrow \infty} \left| \frac{a-n}{n+1} \right| = 1.$$

The radius of convergence of $T_a(x)$ is therefore $R = r^{-1} = 1$.

If a is a whole number, then $\binom{a}{n} = 0$ for all $n > a$. The infinite series then reduces to a polynomial of degree a , so it converges for all x (i.e. $R = \infty$).

87. By Exercise 86, $T_a(x)$ converges for $|x| < 1$, but we do not yet know whether $T_a(x) = (1+x)^a$.

(a) Verify the identity

$$a \binom{a}{n} = n \binom{a}{n} + (n+1) \binom{a}{n+1}$$

(b) Use (a) to show that $y = T_a(x)$ satisfies the differential equation $(1+x)y' = ay$ with initial condition $y(0) = 1$.

(c) Prove that $T_a(x) = (1+x)^a$ for $|x| < 1$ by showing that the derivative of the ratio $\frac{T_a(x)}{(1+x)^a}$ is zero.

SOLUTION**(a)**

$$\begin{aligned} n \binom{a}{n} + (n+1) \binom{a}{n+1} &= n \cdot \frac{a(a-1)\cdots(a-n+1)}{n!} + (n+1) \cdot \frac{a(a-1)\cdots(a-n+1)(a-n)}{(n+1)!} \\ &= \frac{a(a-1)\cdots(a-n+1)}{(n-1)!} + \frac{a(a-1)\cdots(a-n+1)(a-n)}{n!} \\ &= \frac{a(a-1)\cdots(a-n+1)(n+(a-n))}{n!} = a \cdot \binom{a}{n} \end{aligned}$$

(b) Differentiating $T_a(x)$ term-by-term yields

$$T'_a(x) = \sum_{n=1}^{\infty} n \binom{a}{n} x^{n-1}.$$

Thus,

$$\begin{aligned} (1+x)T'_a(x) &= \sum_{n=1}^{\infty} n \binom{a}{n} x^{n-1} + \sum_{n=1}^{\infty} n \binom{a}{n} x^n = \sum_{n=0}^{\infty} (n+1) \binom{a}{n+1} x^n + \sum_{n=0}^{\infty} n \binom{a}{n} x^n \\ &= \sum_{n=0}^{\infty} \left[(n+1) \binom{a}{n+1} + n \binom{a}{n} \right] x^n = a \sum_{n=0}^{\infty} \binom{a}{n} x^n = aT_a(x). \end{aligned}$$

Moreover,

$$T_a(0) = \binom{a}{0} = 1.$$

(c)

$$\frac{d}{dx} \left(\frac{T_a(x)}{(1+x)^a} \right) = \frac{(1+x)^a T'_a(x) - a(1+x)^{a-1} T_a(x)}{(1+x)^{2a}} = \frac{(1+x)T'_a(x) - aT_a(x)}{(1+x)^{a+1}} = 0.$$

Thus,

$$\frac{T_a(x)}{(1+x)^a} = C,$$

for some constant C . For $x = 0$,

$$\frac{T_a(0)}{(1+0)^a} = \frac{1}{1} = 1, \text{ so } C = 1.$$

Finally, $T_a(x) = (1+x)^a$.**88.** The function $G(k) = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 t} dt$ is called an **elliptic function of the second kind**. Prove that for $|k| < 1$,

$$G(k) = \frac{\pi}{2} - \frac{\pi}{2} \sum_{n=1}^{\infty} \left(\frac{1 \cdot 3 \cdots (2n-1)}{2 \cdots 4 \cdots (2n)} \right)^2 \frac{k^{2n}}{2n-1}$$

SOLUTION For $|k| < 1$, $|k^2 \sin^2 t| < 1$ for all t . Substituting $-k^2 \sin^2 t$ for t in the binomial series for $a = \frac{1}{2}$, we find

$$\begin{aligned} \sqrt{1-k^2 \sin^2 t} &= 1 + \sum_{n=1}^{\infty} \binom{\frac{1}{2}}{n} (-k^2 \sin^2 t)^n \\ &= 1 + \sum_{n=1}^{\infty} (-1)^n \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n+1\right)}{n!} k^{2n} \sin^{2n} t \\ &= 1 + \sum_{n=1}^{\infty} (-1)^n \frac{1(1-2)(1-4)\cdots(1-2(n-1))}{2^n n!} k^{2n} \sin^{2n} t \\ &= 1 + \sum_{n=1}^{\infty} (-1)^n (-1)^{n-1} \frac{(2-1)(4-1)\cdots(2n-3)}{2^n n!} k^{2n} \sin^{2n} t \\ &= 1 - \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2 \cdot 4 \cdot 6 \cdots (2n)} k^{2n} \sin^{2n} t. \end{aligned}$$

Integrating from 0 to $\frac{\pi}{2}$ term-by-term, we obtain

$$G(k) = \frac{\pi}{2} - \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2 \cdot 4 \cdot 6 \cdots (2n)} k^{2n} \int_0^{\pi/2} \sin^{2n} t \, dt.$$

Finally, using the formula

$$\int_0^{\pi/2} \sin^{2n} t \, dt = \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \frac{\pi}{2},$$

we arrive at

$$G(k) = \frac{\pi}{2} - \frac{\pi}{2} \sum_{n=1}^{\infty} \left(\frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2 \cdot 4 \cdot 6 \cdots (2n)} \right)^2 (2n-1) k^{2n} = \frac{\pi}{2} - \frac{\pi}{2} \sum_{n=1}^{\infty} \left(\frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)} \right)^2 \frac{k^{2n}}{2n-1}.$$

89. Assume that $a < b$ and let L be the arc length (circumference) of the ellipse $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$ shown in Figure 5. There is no explicit formula for L , but it is known that $L = 4bG(k)$, with $G(k)$ as in Exercise 88 and $k = \sqrt{1 - a^2/b^2}$. Use the first three terms of the expansion of Exercise 88 to estimate L when $a = 4$ and $b = 5$.




FIGURE 5 The ellipse $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$.

SOLUTION With $a = 4$ and $b = 5$,

$$k = \sqrt{1 - \frac{4^2}{5^2}} = \frac{3}{5},$$

and the arc length of the ellipse $\left(\frac{x}{4}\right)^2 + \left(\frac{y}{5}\right)^2 = 1$ is

$$L = 20G\left(\frac{3}{5}\right) = 20 \left(\frac{\pi}{2} - \frac{\pi}{2} \sum_{n=1}^{\infty} \left(\frac{1 \cdot 3 \cdots (2n-1)}{2 \cdot 4 \cdots (2n)} \right)^2 \frac{\left(\frac{3}{5}\right)^{2n}}{2n-1} \right).$$

Using the first three terms in the series for $G(k)$ gives

$$L \approx 10\pi - 10\pi \left(\left(\frac{1}{2}\right)^2 \cdot \frac{(3/5)^2}{1} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \cdot \frac{(3/5)^4}{3} \right) = 10\pi \left(1 - \frac{9}{100} - \frac{243}{40,000} \right) = \frac{36,157\pi}{4000} \approx 28.398.$$

90. Use Exercise 88 to prove that if $a < b$ and a/b is near 1 (a nearly circular ellipse), then

$$L \approx \frac{\pi}{2} \left(3b + \frac{a^2}{b} \right)$$

Hint: Use the first two terms of the series for $G(k)$.

SOLUTION From the previous exercise, we know that

$$L = 4bG(k), \quad \text{where } k = \sqrt{1 - \frac{a^2}{b^2}}.$$

Following the hint and using only the first two terms of the series expansion for $G(k)$ from Exercise 88, we find

$$L \approx 4b \left(\frac{\pi}{2} - \frac{\pi}{2} \left(\frac{1}{2}\right)^2 k^2 \right) = \frac{\pi}{2} \left(4b - b \left(1 - \frac{a^2}{b^2} \right) \right) = \frac{\pi}{2} \left(3b + \frac{a^2}{b} \right).$$

91. Irrationality of e Prove that e is an irrational number using the following argument by contradiction. Suppose that $e = M/N$, where M, N are nonzero integers.

(a) Show that $M!e^{-1}$ is a whole number.

(b) Use the power series for e^x at $x = -1$ to show that there is an integer B such that $M!e^{-1}$ equals

$$B + (-1)^{M+1} \left(\frac{1}{M+1} - \frac{1}{(M+1)(M+2)} + \cdots \right)$$

(c) Use your knowledge of alternating series with decreasing terms to conclude that $0 < |M!e^{-1} - B| < 1$ and observe that this contradicts (a). Hence, e is not equal to M/N .

SOLUTION Suppose that $e = M/N$, where M, N are nonzero integers.

(a) With $e = M/N$,

$$M!e^{-1} = M! \frac{N}{M} = (M-1)!N,$$

which is a whole number.

(b) Substituting $x = -1$ into the Maclaurin series for e^x and multiplying the resulting series by $M!$ yields

$$M!e^{-1} = M! \left(1 - 1 + \frac{1}{2!} - \frac{1}{3!} + \cdots + \frac{(-1)^k}{k!} + \cdots \right).$$

For all $k \leq M$, $\frac{M!}{k!}$ is a whole number, so

$$M! \left(1 - 1 + \frac{1}{2!} - \frac{1}{3!} + \cdots + \frac{(-1)^k}{M!} \right)$$

is an integer. Denote this integer by B . Thus,

$$M!e^{-1} = B + M! \left(\frac{(-1)^{M+1}}{(M+1)!} + \frac{(-1)^{M+2}}{(M+2)!} + \cdots \right) = B + (-1)^{M+1} \left(\frac{1}{M+1} - \frac{1}{(M+1)(M+2)} + \cdots \right).$$

(c) The series for $M!e^{-1}$ obtained in part (b) is an alternating series with $a_n = \frac{M!}{n!}$. Using the error bound for an alternating series and noting that $B = S_M$, we have

$$|M!e^{-1} - B| \leq a_{M+1} = \frac{1}{M+1} < 1.$$

This inequality implies that $M!e^{-1} - B$ is not a whole number; however, B is a whole number so $M!e^{-1}$ cannot be a whole number. We get a contradiction to the result in part (a), which proves that the original assumption that e is a rational number is false.

92. Use the result of Exercise 73 in Section 4.5 to show that the Maclaurin series of the function

$$f(x) = \begin{cases} e^{-1/x^2} & \text{for } x \neq 0 \\ 0 & \text{for } x = 0 \end{cases}$$

is $T(x) = 0$. This provides an example of a function $f(x)$ whose Maclaurin series converges but does not converge to $f(x)$ (except at $x = 0$).

SOLUTION By the referenced exercise, $f(x)$ has continuous derivatives of all orders at 0, and $f^{(n)}(0) = 0$ for all $n > 0$. But then the Maclaurin series is

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = 0$$

CHAPTER REVIEW EXERCISES

- Let $a_n = \frac{n-3}{n!}$ and $b_n = a_{n+3}$. Calculate the first three terms in each sequence.
 - a_n^2
 - b_n
 - $a_n b_n$
 - $2a_{n+1} - 3a_n$

SOLUTION

(a)

$$a_1^2 = \left(\frac{1-3}{1!}\right)^2 = (-2)^2 = 4;$$

$$a_2^2 = \left(\frac{2-3}{2!}\right)^2 = \left(-\frac{1}{2}\right)^2 = \frac{1}{4};$$

$$a_3^2 = \left(\frac{3-3}{3!}\right)^2 = 0.$$

(b)

$$b_1 = a_4 = \frac{4-3}{4!} = \frac{1}{24};$$

$$b_2 = a_5 = \frac{5-3}{5!} = \frac{1}{60};$$

$$b_3 = a_6 = \frac{6-3}{6!} = \frac{1}{240}.$$

(c) Using the formula for a_n and the values in (b) we obtain:

$$a_1 b_1 = \frac{1-3}{1!} \cdot \frac{1}{24} = -\frac{1}{12};$$

$$a_2 b_2 = \frac{2-3}{2!} \cdot \frac{1}{60} = -\frac{1}{120};$$

$$a_3 b_3 = \frac{3-3}{3!} \cdot \frac{1}{240} = 0.$$

(d)

$$2a_2 - 3a_1 = 2\left(-\frac{1}{2}\right) - 3(-2) = 5;$$

$$2a_3 - 3a_2 = 2 \cdot 0 - 3\left(-\frac{1}{2}\right) = \frac{3}{2};$$

$$2a_4 - 3a_3 = 2 \cdot \frac{1}{24} - 3 \cdot 0 = \frac{1}{12}.$$

2. Prove that $\lim_{n \rightarrow \infty} \frac{2n-1}{3n+2} = \frac{2}{3}$ using the limit definition.

SOLUTION Note

$$\left| \frac{2n-1}{3n+2} - \frac{2}{3} \right| = \left| \frac{6n-3-2(3n+2)}{3(3n+2)} \right| = \left| -\frac{7}{3(3n+2)} \right| = \frac{7}{3(3n+2)} < \frac{7}{9n}.$$

Therefore, to have $\left| a_n - \frac{2}{3} \right| < \epsilon$, we need

$$\frac{7}{9n} < \epsilon \quad \text{or} \quad n > \frac{7}{9\epsilon}.$$

Thus, let $\epsilon > 0$ and take $M = \frac{7}{9\epsilon}$. Then, whenever $n > M$,

$$\left| \frac{2n-1}{3n+2} - \frac{2}{3} \right| = \frac{7}{3(3n+2)} < \frac{7}{9n} < \frac{7}{9} \cdot \frac{9\epsilon}{7} = \epsilon.$$

In Exercises 3–8, compute the limit (or state that it does not exist) assuming that $\lim_{n \rightarrow \infty} a_n = 2$.

3. $\lim_{n \rightarrow \infty} (5a_n - 2a_n^2)$

SOLUTION

$$\lim_{n \rightarrow \infty} (5a_n - 2a_n^2) = 5 \lim_{n \rightarrow \infty} a_n - 2 \lim_{n \rightarrow \infty} a_n^2 = 5 \lim_{n \rightarrow \infty} a_n - 2 \left(\lim_{n \rightarrow \infty} a_n \right)^2 = 5 \cdot 2 - 2 \cdot 2^2 = 2.$$

$$4. \lim_{n \rightarrow \infty} \frac{1}{a_n}$$

$$\text{SOLUTION} \quad \lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{1}{\lim_{n \rightarrow \infty} a_n} = \frac{1}{2}.$$

$$5. \lim_{n \rightarrow \infty} e^{a_n}$$

SOLUTION The function $f(x) = e^x$ is continuous, hence:

$$\lim_{n \rightarrow \infty} e^{a_n} = e^{\lim_{n \rightarrow \infty} a_n} = e^2.$$

$$6. \lim_{n \rightarrow \infty} \cos(\pi a_n)$$

SOLUTION The function $f(x) = \cos(\pi x)$ is continuous, hence:

$$\lim_{n \rightarrow \infty} \cos(\pi a_n) = \cos\left(\pi \lim_{n \rightarrow \infty} a_n\right) = \cos(2\pi) = 1.$$

$$7. \lim_{n \rightarrow \infty} (-1)^n a_n$$

SOLUTION Because $\lim_{n \rightarrow \infty} a_n \neq 0$, it follows that $\lim_{n \rightarrow \infty} (-1)^n a_n$ does not exist.

$$8. \lim_{n \rightarrow \infty} \frac{a_n + n}{a_n + n^2}$$

SOLUTION Because the sequence $\{a_n\}$ converges, $\{a_n\}$ is bounded and

$$\lim_{n \rightarrow \infty} \frac{a_n}{n^2} = 0.$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{a_n + n}{a_n + n^2} = \lim_{n \rightarrow \infty} \frac{\frac{a_n}{n^2} + \frac{1}{n}}{\frac{a_n}{n^2} + 1} = \frac{0 + 0}{0 + 1} = 0.$$

In Exercises 9–22, determine the limit of the sequence or show that the sequence diverges.

$$9. a_n = \sqrt{n+5} - \sqrt{n+2}$$

SOLUTION First rewrite a_n as follows:

$$a_n = \frac{(\sqrt{n+5} - \sqrt{n+2})(\sqrt{n+5} + \sqrt{n+2})}{\sqrt{n+5} + \sqrt{n+2}} = \frac{(n+5) - (n+2)}{\sqrt{n+5} + \sqrt{n+2}} = \frac{3}{\sqrt{n+5} + \sqrt{n+2}}.$$

Thus,

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{3}{\sqrt{n+5} + \sqrt{n+2}} = 0.$$

$$10. a_n = \frac{3n^3 - n}{1 - 2n^3}$$

$$\text{SOLUTION} \quad \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{3n^3 - n}{1 - 2n^3} = -\frac{3}{2}.$$

$$11. a_n = 2^{1/n^2}$$

SOLUTION The function $f(x) = 2^x$ is continuous, so

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} 2^{1/n^2} = 2^{\lim_{n \rightarrow \infty} (1/n^2)} = 2^0 = 1.$$

$$12. a_n = \frac{10^n}{n!}$$

SOLUTION For $n > 10$, write a_n as

$$0 \leq a_n = \underbrace{\left(\frac{10}{1} \cdot \frac{10}{2} \cdots \frac{10}{10}\right)}_{\text{equals } \frac{10^{10}}{10!}} \underbrace{\left(\frac{10}{11}\right) \cdot \left(\frac{10}{12}\right) \cdots \left(\frac{10}{n}\right)}_{\text{each factor is less than 1}} < \frac{10^{10}}{10!} \cdot \frac{10}{n} = \frac{10^{10}}{9!n};$$

Thus, by the Squeeze Theorem, $\lim_{n \rightarrow \infty} a_n = 0$.

$$13. b_m = 1 + (-1)^m$$

SOLUTION Because $1 + (-1)^m$ is equal to 0 for m odd and is equal to 2 for m even, the sequence $\{b_m\}$ does not approach one limit; hence this sequence diverges.

$$14. b_m = \frac{1 + (-1)^m}{m}$$

SOLUTION The numerator is equal to zero for m odd and is equal to 2 for m even. Therefore,

$$0 \leq \frac{1 + (-1)^m}{m} \leq \frac{2}{m},$$

and by the Squeeze Theorem, $\lim_{m \rightarrow \infty} b_m = 0$.

$$15. b_n = \tan^{-1} \left(\frac{n+2}{n+5} \right)$$

SOLUTION The function $\tan^{-1}x$ is continuous, so

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \tan^{-1} \left(\frac{n+2}{n+5} \right) = \tan^{-1} \left(\lim_{n \rightarrow \infty} \frac{n+2}{n+5} \right) = \tan^{-1} 1 = \frac{\pi}{4}.$$

$$16. a_n = \frac{100^n}{n!} - \frac{3 + \pi^n}{5^n}$$

SOLUTION For $n > 100$,

$$0 \leq \frac{100^n}{n!} = \left(\frac{100}{1} \cdot \frac{100}{2} \cdots \frac{100}{100} \right) \frac{100}{101} \cdot \frac{100}{102} \cdots \frac{100}{n} < \frac{100^{100}}{99!n};$$

therefore,

$$\lim_{n \rightarrow \infty} \frac{100^n}{n!} = 0$$

by the Squeeze Theorem. Moreover,

$$\lim_{n \rightarrow \infty} \left(\frac{3 + \pi^n}{5^n} \right) = \lim_{n \rightarrow \infty} \frac{3}{5^n} + \lim_{n \rightarrow \infty} \left(\frac{\pi}{5} \right)^n = 0 + 0 = 0.$$

Thus,

$$\lim_{n \rightarrow \infty} a_n = 0 + 0 = 0.$$

$$17. b_n = \sqrt{n^2 + n} - \sqrt{n^2 + 1}$$

SOLUTION Rewrite b_n as

$$b_n = \frac{(\sqrt{n^2 + n} - \sqrt{n^2 + 1})(\sqrt{n^2 + n} + \sqrt{n^2 + 1})}{\sqrt{n^2 + n} + \sqrt{n^2 + 1}} = \frac{(n^2 + n) - (n^2 + 1)}{\sqrt{n^2 + n} + \sqrt{n^2 + 1}} = \frac{n - 1}{\sqrt{n^2 + n} + \sqrt{n^2 + 1}}.$$

Then

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{\frac{n}{n} - \frac{1}{n}}{\sqrt{\frac{n^2}{n^2} + \frac{n}{n^2}} + \sqrt{\frac{n^2}{n^2} + \frac{1}{n^2}}} = \lim_{n \rightarrow \infty} \frac{1 - \frac{1}{n}}{\sqrt{1 + \frac{1}{n}} + \sqrt{1 + \frac{1}{n^2}}} = \frac{1 - 0}{\sqrt{1 + 0} + \sqrt{1 + 0}} = \frac{1}{2}.$$

$$18. c_n = \sqrt{n^2 + n} - \sqrt{n^2 - n}$$

SOLUTION Rewrite c_n as

$$c_n = \frac{(\sqrt{n^2 + n} - \sqrt{n^2 - n})(\sqrt{n^2 + n} + \sqrt{n^2 - n})}{\sqrt{n^2 + n} + \sqrt{n^2 - n}} = \frac{(n^2 + n) - (n^2 - n)}{\sqrt{n^2 + n} + \sqrt{n^2 - n}} = \frac{2n}{\sqrt{n^2 + n} + \sqrt{n^2 - n}}.$$

Then

$$\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \frac{\frac{2n}{n}}{\sqrt{\frac{n^2}{n^2} + \frac{n}{n^2}} + \sqrt{\frac{n^2}{n^2} - \frac{n}{n^2}}} = \lim_{n \rightarrow \infty} \frac{2}{\sqrt{1 + \frac{1}{n}} + \sqrt{1 - \frac{1}{n}}} = \frac{2}{\sqrt{1 + 0} + \sqrt{1 - 0}} = 1.$$

$$19. b_m = \left(1 + \frac{1}{m}\right)^{3m}$$

$$\text{SOLUTION } \lim_{m \rightarrow \infty} b_m = \lim_{m \rightarrow \infty} \left(1 + \frac{1}{m}\right)^m = e.$$

$$20. c_n = \left(1 + \frac{3}{n}\right)^n$$

SOLUTION Write

$$c_n = \left(1 + \frac{1}{n/3}\right)^n = \left[\left(1 + \frac{1}{n/3}\right)^{n/3}\right]^3.$$

Then, because x^3 is a continuous function,

$$\lim_{n \rightarrow \infty} c_n = \left[\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n/3}\right)^{n/3}\right]^3 = e^3.$$

$$21. b_n = n(\ln(n+1) - \ln n)$$

SOLUTION Write

$$b_n = n \ln\left(\frac{n+1}{n}\right) = \frac{\ln\left(1 + \frac{1}{n}\right)}{\frac{1}{n}}.$$

Using L'Hôpital's Rule, we find

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{\ln\left(1 + \frac{1}{n}\right)}{\frac{1}{n}} = \lim_{x \rightarrow \infty} \frac{\ln\left(1 + \frac{1}{x}\right)}{\frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{\left(1 + \frac{1}{x}\right)^{-1} \cdot \left(-\frac{1}{x^2}\right)}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{-1} = 1.$$

$$22. c_n = \frac{\ln(n^2 + 1)}{\ln(n^3 + 1)}$$

SOLUTION Using L'Hôpital's Rule, we find

$$\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} \frac{\ln(n^2 + 1)}{\ln(n^3 + 1)} = \lim_{n \rightarrow \infty} \frac{2n/(n^2 + 1)}{3n^2/(n^3 + 1)} = \lim_{n \rightarrow \infty} \frac{2n^4 + 2n}{3n^4 + 3n^2} = \lim_{n \rightarrow \infty} \frac{2 + 2n^{-3}}{3 + 3n^{-2}} = \frac{2}{3}$$

$$23. \text{ Use the Squeeze Theorem to show that } \lim_{n \rightarrow \infty} \frac{\arctan(n^2)}{\sqrt{n}} = 0.$$

SOLUTION For all x ,

$$-\frac{\pi}{2} < \arctan x < \frac{\pi}{2},$$

so

$$-\frac{\pi/2}{\sqrt{n}} < \frac{\arctan(n^2)}{\sqrt{n}} < \frac{\pi/2}{\sqrt{n}},$$

for all n . Because

$$\lim_{n \rightarrow \infty} \left(-\frac{\pi/2}{\sqrt{n}}\right) = \lim_{n \rightarrow \infty} \frac{\pi/2}{\sqrt{n}} = 0,$$

it follows by the Squeeze Theorem that

$$\lim_{n \rightarrow \infty} \frac{\arctan(n^2)}{\sqrt{n}} = 0.$$

24. Give an example of a divergent sequence $\{a_n\}$ such that $\{\sin a_n\}$ is convergent.

SOLUTION Let $a_n = (-1)^n \pi$. This is an alternating series, which does not approach 0, hence it diverges. However, a_n is a multiple of π for every n , and thus, $\sin a_n = 0$. Since $\{\sin a_n\}$ is a constant sequence, it converges.

25. Calculate $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$, where $a_n = \frac{1}{2}3^n - \frac{1}{3}2^n$.

SOLUTION Because

$$\frac{1}{2}3^n - \frac{1}{3}2^n \geq \frac{1}{2}3^n - \frac{1}{3}3^n = \frac{3^n}{6}$$

and

$$\lim_{n \rightarrow \infty} \frac{3^n}{6} = \infty,$$

we conclude that $\lim_{n \rightarrow \infty} a_n = \infty$, so L'Hôpital's rule may be used:

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{2}3^{n+1} - \frac{1}{3}2^{n+1}}{\frac{1}{2}3^n - \frac{1}{3}2^n} = \lim_{n \rightarrow \infty} \frac{3^{n+2} - 2^{n+2}}{3^{n+1} - 2^{n+1}} = \lim_{n \rightarrow \infty} \frac{3 - 2\left(\frac{2}{3}\right)^{n+1}}{1 - \left(\frac{2}{3}\right)^{n+1}} = \frac{3 - 0}{1 - 0} = 3.$$

26. Define $a_{n+1} = \sqrt{a_n + 6}$ with $a_1 = 2$.

- (a) Compute a_n for $n = 2, 3, 4, 5$.
 (b) Show that $\{a_n\}$ is increasing and is bounded by 3.
 (c) Prove that $\lim_{n \rightarrow \infty} a_n$ exists and find its value.

SOLUTION

(a) We compute the first four values of a_n recursively:

$$a_2 = \sqrt{a_1 + 6} = \sqrt{2 + 6} = \sqrt{8} = 2\sqrt{2} \approx 2.828427;$$

$$a_3 = \sqrt{a_2 + 6} = \sqrt{2\sqrt{2} + 6} \approx 2.971267;$$

$$a_4 = \sqrt{a_3 + 6} = \sqrt{\sqrt{2\sqrt{2} + 6} + 6} \approx 2.995207;$$

$$a_5 = \sqrt{a_4 + 6} = \sqrt{\sqrt{\sqrt{2\sqrt{2} + 6} + 6} + 6} \approx 2.999201.$$

(b) By part (a) and the given data, $a_2 \approx 2.8$ and $a_1 = 2$, so $a_2 > a_1$. Now, suppose that $a_k > a_{k-1}$; then

$$a_{k+1} = \sqrt{a_k + 6} > \sqrt{a_{k-1} + 6} = a_k.$$

Thus, by mathematical induction, $a_{n+1} > a_n$ for all n and $\{a_n\}$ is increasing.

Next, note that $a_1 = 2 < 3$. Suppose $a_k < 3$, then

$$a_{k+1} = \sqrt{a_k + 6} < \sqrt{3 + 6} = 3.$$

Thus, by mathematical induction, $a_n < 3$ for all n .

(c) Since $\{a_n\}$ is increasing and has an upper bound, $\{a_n\}$ converges. Let

$$L = \lim_{n \rightarrow \infty} a_n.$$

Then,

$$L = \sqrt{L + 6}$$

$$L^2 = L + 6$$

$$L^2 - L - 6 = 0$$

$$(L - 3)(L + 2) = 0$$

so $L = 3$ or $L = -2$; however, the sequence is increasing and its first term is positive, so -2 cannot be the limit. Therefore,

$$\lim_{n \rightarrow \infty} a_n = 3.$$

27. Calculate the partial sums S_4 and S_7 of the series $\sum_{n=1}^{\infty} \frac{n-2}{n^2+2n}$.

SOLUTION

$$S_4 = -\frac{1}{3} + 0 + \frac{1}{15} + \frac{2}{24} = -\frac{11}{60} = -0.183333;$$

$$S_7 = -\frac{1}{3} + 0 + \frac{1}{15} + \frac{2}{24} + \frac{3}{35} + \frac{4}{48} + \frac{5}{63} = \frac{287}{4410} = 0.065079.$$

28. Find the sum $1 - \frac{1}{4} + \frac{1}{4^2} - \frac{1}{4^3} + \cdots$.

SOLUTION This is a geometric series with $r = -\frac{1}{4}$. Therefore,

$$1 - \frac{1}{4} + \frac{1}{4^2} - \frac{1}{4^3} + \cdots = \frac{1}{1 - (-\frac{1}{4})} = \frac{4}{5}.$$

29. Find the sum $\frac{4}{9} + \frac{8}{27} + \frac{16}{81} + \frac{32}{243} + \cdots$.

SOLUTION This is a geometric series with common ratio $r = \frac{2}{3}$. Therefore,

$$\frac{4}{9} + \frac{8}{27} + \frac{16}{81} + \frac{32}{243} + \cdots = \frac{\frac{4}{9}}{1 - \frac{2}{3}} = \frac{4}{3}.$$

30. Find the sum $\sum_{n=2}^{\infty} \left(\frac{2}{e}\right)^n$.

SOLUTION This is a geometric series with common ratio $r = \frac{2}{e}$. Therefore,

$$\sum_{n=2}^{\infty} \left(\frac{2}{e}\right)^n = \frac{\left(\frac{2}{e}\right)^2}{1 - \frac{2}{e}} = \frac{\frac{4}{e^2}}{1 - \frac{2}{e}} = \frac{4}{e^2 - 2e}.$$

31. Find the sum $\sum_{n=-1}^{\infty} \frac{2^{n+3}}{3^n}$.

SOLUTION Note

$$\sum_{n=-1}^{\infty} \frac{2^{n+3}}{3^n} = 2^3 \sum_{n=-1}^{\infty} \frac{2^n}{3^n} = 8 \sum_{n=-1}^{\infty} \left(\frac{2}{3}\right)^n;$$

therefore,

$$\sum_{n=-1}^{\infty} \frac{2^{n+3}}{3^n} = 8 \cdot \frac{3}{2} \cdot \frac{1}{1 - \frac{2}{3}} = 36.$$

32. Show that $\sum_{n=1}^{\infty} (b - \tan^{-1} n^2)$ diverges if $b \neq \frac{\pi}{2}$.

SOLUTION Note

$$\lim_{n \rightarrow \infty} (b - \tan^{-1} n^2) = b - \lim_{n \rightarrow \infty} \tan^{-1} n^2 = b - \frac{\pi}{2}.$$

If $b \neq \frac{\pi}{2}$, then the limit of the terms in the series is not 0; hence, the series diverges by the Divergence Test.

33. Give an example of divergent series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ such that $\sum_{n=1}^{\infty} (a_n + b_n) = 1$.

SOLUTION Let $a_n = \left(\frac{1}{2}\right)^n + 1$, $b_n = -1$. The corresponding series diverge by the Divergence Test; however,

$$\sum_{n=1}^{\infty} (a_n + b_n) = \sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n = \frac{\frac{1}{2}}{1 - \frac{1}{2}} = 1.$$

34. Let $S = \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+2} \right)$. Compute S_N for $N = 1, 2, 3, 4$. Find S by showing that

$$S_N = \frac{3}{2} - \frac{1}{N+1} - \frac{1}{N+2}$$

SOLUTION

$$S_1 = 1 - \frac{1}{3} = \frac{2}{3};$$

$$S_2 = \left(1 - \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{1}{4}\right) = \frac{3}{2} - \frac{7}{12} = \frac{11}{12};$$

$$S_3 = \left(1 - \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) = \frac{3}{2} - \frac{9}{20} = \frac{21}{20};$$

$$S_4 = \left(1 - \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) + \left(\frac{1}{4} - \frac{1}{6}\right) = \frac{3}{2} - \frac{11}{30} = \frac{17}{15}.$$

The general term in the sequence of partial sums is

$$\begin{aligned} S_N &= \left(1 - \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) + \left(\frac{1}{4} - \frac{1}{6}\right) + \cdots + \left(\frac{1}{N-1} - \frac{1}{N+1}\right) + \left(\frac{1}{N} - \frac{1}{N+2}\right) \\ &= 1 + \frac{1}{2} - \frac{1}{N+1} - \frac{1}{N+2} = \frac{3}{2} - \left(\frac{1}{N+1} + \frac{1}{N+2}\right). \end{aligned}$$

Finally,

$$S = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} \left[\frac{3}{2} - \left(\frac{1}{N+1} + \frac{1}{N+2}\right) \right] = \frac{3}{2}.$$

35. Evaluate $S = \sum_{n=3}^{\infty} \frac{1}{n(n+3)}$.

SOLUTION Note that

$$\frac{1}{n(n+3)} = \frac{1}{3} \left(\frac{1}{n} - \frac{1}{n+3} \right)$$

so that

$$\begin{aligned} \sum_{n=3}^N \frac{1}{n(n+3)} &= \frac{1}{3} \sum_{n=3}^N \left(\frac{1}{n} - \frac{1}{n+3} \right) \\ &= \frac{1}{3} \left(\left(\frac{1}{3} - \frac{1}{6} \right) + \left(\frac{1}{4} - \frac{1}{7} \right) + \left(\frac{1}{5} - \frac{1}{8} \right) \right. \\ &\quad \left. + \left(\frac{1}{6} - \frac{1}{9} \right) + \cdots + \left(\frac{1}{N-1} - \frac{1}{N+2} \right) + \left(\frac{1}{N} - \frac{1}{N+3} \right) \right) \\ &= \frac{1}{3} \left(\frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{N+1} - \frac{1}{N+2} - \frac{1}{N+3} \right) \end{aligned}$$

Thus

$$\begin{aligned} \sum_{n=3}^{\infty} \frac{1}{n(n+3)} &= \frac{1}{3} \lim_{N \rightarrow \infty} \sum_{n=3}^N \left(\frac{1}{n} - \frac{1}{n+3} \right) \\ &= \frac{1}{3} \left(\frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{N+1} - \frac{1}{N+2} - \frac{1}{N+3} \right) = \frac{1}{3} \left(\frac{1}{3} + \frac{1}{4} + \frac{1}{5} \right) = \frac{47}{180} \end{aligned}$$

36. Find the total area of the infinitely many circles on the interval $[0, 1]$ in Figure 1.




FIGURE 1

SOLUTION The diameter of the largest circle is $\frac{1}{2}$, and the diameter of each smaller circle is $\frac{1}{2}$ the diameter of the previous circle; thus, the diameter of the n th circle (for $n \geq 1$) is $\frac{1}{2^n}$ and the area is

$$\pi \left(\frac{1}{2^{n+1}} \right)^2 = \frac{\pi}{4^{n+1}}.$$

The total area of the circles is

$$\sum_{n=1}^{\infty} \frac{\pi}{4^{n+1}} = \frac{\pi}{4} \sum_{n=1}^{\infty} \left(\frac{1}{4} \right)^n = \frac{\pi}{4} \cdot \frac{\frac{1}{4}}{1 - \frac{1}{4}} = \frac{\pi}{12}.$$

In Exercises 37–40, use the Integral Test to determine whether the infinite series converges.

$$37. \sum_{n=1}^{\infty} \frac{n^2}{n^3 + 1}$$

SOLUTION Let $f(x) = \frac{x^2}{x^3+1}$. This function is continuous and positive for $x \geq 1$. Because

$$f'(x) = \frac{(x^3 + 1)(2x) - x^2(3x^2)}{(x^3 + 1)^2} = \frac{x(2 - x^3)}{(x^3 + 1)^2},$$

we see that $f'(x) < 0$ and f is decreasing on the interval $x \geq 2$. Therefore, the Integral Test applies on the interval $x \geq 2$. Now,

$$\int_2^{\infty} \frac{x^2}{x^3 + 1} dx = \lim_{R \rightarrow \infty} \int_2^R \frac{x^2}{x^3 + 1} dx = \frac{1}{3} \lim_{R \rightarrow \infty} (\ln(R^3 + 1) - \ln 9) = \infty.$$

The integral diverges; hence, the series $\sum_{n=2}^{\infty} \frac{n^2}{n^3 + 1}$ diverges, as does the series $\sum_{n=1}^{\infty} \frac{n^2}{n^3 + 1}$.

$$38. \sum_{n=1}^{\infty} \frac{n^2}{(n^3 + 1)^{1.01}}$$

SOLUTION Let $f(x) = \frac{x^2}{(x^3+1)^{1.01}}$. This function is continuous and positive for $x \geq 1$. Because

$$f'(x) = \frac{(x^3 + 1)^{1.01}(2x) - x^2 \cdot 1.01(x^3 + 1)^{0.01}(3x^2)}{(x^3 + 1)^{2.02}} = \frac{x(x^3 + 1)^{0.01}(2 - 1.03x^3)}{(x^3 + 1)^{2.02}},$$

we see that $f'(x) < 0$ and f is decreasing on the interval $x \geq 2$. Therefore, the Integral Test applies on the interval $x \geq 2$. Now,

$$\int_2^{\infty} \frac{x^2}{(x^3 + 1)^{1.01}} dx = \lim_{R \rightarrow \infty} \int_2^R \frac{x^2}{(x^3 + 1)^{1.01}} dx = -\frac{1}{0.03} \lim_{R \rightarrow \infty} \left(\frac{1}{(R^3 + 1)^{0.01}} - \frac{1}{9^{0.01}} \right) = \frac{1}{0.03 \cdot 9^{0.01}}.$$

The integral converges; hence, the series $\sum_{n=2}^{\infty} \frac{n^2}{(n^3 + 1)^{1.01}}$ converges, as does the series $\sum_{n=1}^{\infty} \frac{n^2}{(n^3 + 1)^{1.01}}$.

$$39. \sum_{n=1}^{\infty} \frac{1}{(n+2)(\ln(n+2))^3}$$

SOLUTION Let $f(x) = \frac{1}{(x+2)\ln^3(x+2)}$. Using the substitution $u = \ln(x+2)$, so that $du = \frac{1}{x+2} dx$, we have

$$\begin{aligned} \int_0^{\infty} f(x) dx &= \int_{\ln 2}^{\infty} \frac{1}{u^3} du = \lim_{R \rightarrow \infty} \int_{\ln 2}^R \frac{1}{u^3} du = \lim_{R \rightarrow \infty} \left(-\frac{1}{2u^2} \Big|_{\ln 2}^R \right) \\ &= \lim_{R \rightarrow \infty} \left(\frac{1}{2(\ln 2)^2} - \frac{1}{2(\ln R)^2} \right) = \frac{1}{2(\ln 2)^2} \end{aligned}$$

Since the integral of $f(x)$ converges, so does the series.

$$40. \sum_{n=1}^{\infty} \frac{n^3}{e^{n^4}}$$

SOLUTION Let $f(x) = x^3 e^{-x^4}$. This function is continuous and positive for $x \geq 1$. Because

$$f'(x) = x^3(-4x^3 e^{-x^4}) + 3x^2 e^{-x^4} = x^2 e^{-x^4}(3 - 4x^4),$$

we see that $f'(x) < 0$ and f is decreasing on the interval $x \geq 1$. Therefore, the Integral Test applies on the interval $x \geq 1$. Now,

$$\int_1^{\infty} x^3 e^{-x^4} dx = \lim_{R \rightarrow \infty} \int_1^R x^3 e^{-x^4} dx = -\frac{1}{4} \lim_{R \rightarrow \infty} (e^{-R^4} - e^{-1}) = \frac{1}{4e}.$$

The integral converges; hence, the series $\sum_{n=1}^{\infty} \frac{n^3}{e^{n^4}}$ also converges.

In Exercises 41–48, use the Comparison or Limit Comparison Test to determine whether the infinite series converges.

$$41. \sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$$

SOLUTION For all $n \geq 1$,

$$0 < \frac{1}{n+1} < \frac{1}{n} \quad \text{so} \quad \frac{1}{(n+1)^2} < \frac{1}{n^2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series, so the series $\sum_{n=1}^{\infty} \frac{1}{(n+1)^2}$ converges by the Comparison Test.

$$42. \sum_{n=1}^{\infty} \frac{1}{\sqrt{n} + n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{1}{\sqrt{n} + n}$ and $b_n = \frac{1}{n}$. Now,

$$L = \lim_{n \rightarrow \infty} \frac{\frac{1}{\sqrt{n} + n}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n}{\sqrt{n} + n} = \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{\sqrt{n}} + 1} = 1.$$

Because $L > 0$ and $\sum_{n=1}^{\infty} \frac{1}{n}$ is the divergent harmonic series, we conclude by the Limit Comparison Test that the series

$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n} + n}$ also diverges.

$$43. \sum_{n=2}^{\infty} \frac{n^2 + 1}{n^{3.5} - 2}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{n^2 + 1}{n^{3.5} - 2}$ and $b_n = \frac{1}{n^{1.5}}$. Now,

$$L = \lim_{n \rightarrow \infty} \frac{\frac{n^2 + 1}{n^{3.5} - 2}}{\frac{1}{n^{1.5}}} = \lim_{n \rightarrow \infty} \frac{n^{3.5} + n^{1.5}}{n^{3.5} - 2} = 1.$$

Because L exists and $\sum_{n=1}^{\infty} \frac{1}{n^{1.5}}$ is a convergent p -series, we conclude by the Limit Comparison Test that the series

$\sum_{n=2}^{\infty} \frac{n^2 + 1}{n^{3.5} - 2}$ also converges.

$$44. \sum_{n=1}^{\infty} \frac{1}{n - \ln n}$$

SOLUTION Since $0 \leq \ln n \leq n$ for all $n \geq 1$, we have $0 \leq n - \ln n \leq n$ and

$$\frac{1}{n} \leq \frac{1}{n - \ln n}$$

The harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges, so we conclude by the Comparison Test that $\sum_{n=1}^{\infty} \frac{1}{n - \ln n}$ also diverges.

$$45. \sum_{n=2}^{\infty} \frac{n}{\sqrt{n^5+5}}$$

SOLUTION For all $n \geq 2$,

$$\frac{n}{\sqrt{n^5+5}} < \frac{n}{n^{5/2}} = \frac{1}{n^{3/2}}.$$

The series $\sum_{n=2}^{\infty} \frac{1}{n^{3/2}}$ is a convergent p -series, so the series $\sum_{n=2}^{\infty} \frac{n}{\sqrt{n^5+5}}$ converges by the Comparison Test.

$$46. \sum_{n=1}^{\infty} \frac{1}{3^n - 2^n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{1}{3^n - 2^n}$ and $b_n = \frac{1}{3^n}$. Then,

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{3^n}{3^n - 2^n} = \lim_{n \rightarrow \infty} \frac{1}{1 - \left(\frac{2}{3}\right)^n} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{3^n}$ is a convergent geometric series; because L exists, we may therefore conclude by the Limit Comparison

Test that the series $\sum_{n=1}^{\infty} \frac{1}{3^n - 2^n}$ also converges.

$$47. \sum_{n=1}^{\infty} \frac{n^{10} + 10^n}{n^{11} + 11^n}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{n^{10} + 10^n}{n^{11} + 11^n}$ and $b_n = \left(\frac{10}{11}\right)^n$. Then,

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{n^{10} + 10^n}{n^{11} + 11^n}}{\left(\frac{10}{11}\right)^n} = \lim_{n \rightarrow \infty} \frac{\frac{n^{10} + 10^n}{10^n}}{\frac{n^{11} + 11^n}{11^n}} = \lim_{n \rightarrow \infty} \frac{\frac{n^{10}}{10^n} + 1}{\frac{n^{11}}{11^n} + 1} = 1.$$

The series $\sum_{n=1}^{\infty} \left(\frac{10}{11}\right)^n$ is a convergent geometric series; because L exists, we may therefore conclude by the Limit

Comparison Test that the series $\sum_{n=1}^{\infty} \frac{n^{10} + 10^n}{n^{11} + 11^n}$ also converges.

$$48. \sum_{n=1}^{\infty} \frac{n^{20} + 21^n}{n^{21} + 20^n}$$

SOLUTION Apply the Limit Comparison Theorem with $a_n = \frac{n^{20} + 21^n}{n^{21} + 20^n}$ and $b_n = \left(\frac{21}{20}\right)^n$. Then

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{n^{20} + 21^n}{n^{21} + 20^n}}{\left(\frac{21}{20}\right)^n} = \lim_{n \rightarrow \infty} \frac{\frac{n^{20} + 21^n}{21^n}}{\frac{n^{21} + 20^n}{20^n}} = \lim_{n \rightarrow \infty} \frac{\frac{n^{20}}{21^n} + 1}{\frac{n^{21}}{20^n} + 1} = 1$$

The series $\sum_{n=1}^{\infty} \left(\frac{21}{20}\right)^n$ is a divergent geometric series. Since $L = 1$, the two series either both converge or both diverge;

thus, we may conclude from the Limit Comparison Test that the series $\sum_{n=1}^{\infty} \frac{n^{20} + 21^n}{n^{21} + 20^n}$ diverges.

$$49. \text{ Determine the convergence of } \sum_{n=1}^{\infty} \frac{2^n + n}{3^n - 2} \text{ using the Limit Comparison Test with } b_n = \left(\frac{2}{3}\right)^n.$$

SOLUTION With $a_n = \frac{2^n + n}{3^n - 2}$, we have

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2^n + n}{3^n - 2} \cdot \frac{3^n}{2^n} = \lim_{n \rightarrow \infty} \frac{6^n + n3^n}{6^n - 2^{n+1}} = \lim_{n \rightarrow \infty} \frac{1 + n\left(\frac{1}{2}\right)^n}{1 - 2\left(\frac{1}{3}\right)^n} = 1$$

Since $L = 1$, the two series either both converge or both diverge. Since $\sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n$ is a convergent geometric series, the

Limit Comparison Test tells us that $\sum_{n=1}^{\infty} \frac{2^n + n}{3^n - 2}$ also converges.

50. Determine the convergence of $\sum_{n=1}^{\infty} \frac{\ln n}{1.5^n}$ using the Limit Comparison Test with $b_n = \frac{1}{1.4^n}$.

SOLUTION With $a_n = \frac{\ln n}{1.5^n}$, and using L'Hôpital's Rule,

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{\ln n}{1.5^n}}{\frac{1}{1.4^n}} = \lim_{n \rightarrow \infty} \frac{\ln n}{\left(\frac{1.5}{1.4}\right)^n} \\ &= \lim_{n \rightarrow \infty} \frac{1/n}{\ln(1.5/1.4) \left(\frac{1.5}{1.4}\right)^n} = \frac{1}{\ln(1.5/1.4)} \lim_{n \rightarrow \infty} \frac{\left(\frac{1.4}{1.5}\right)^n}{n} = 0 \end{aligned}$$

Since $L < \infty$ and $\sum_{n=1}^{\infty} b_n$ is a convergent geometric series, it follows from the Limit Comparison Test that $\sum_{n=1}^{\infty} \frac{\ln n}{1.5^n}$ also converges.

51. Let $a_n = 1 - \sqrt{1 - \frac{1}{n}}$. Show that $\lim_{n \rightarrow \infty} a_n = 0$ and that $\sum_{n=1}^{\infty} a_n$ diverges. *Hint:* Show that $a_n \geq \frac{1}{2n}$.

SOLUTION

$$\begin{aligned} 1 - \sqrt{1 - \frac{1}{n}} &= 1 - \sqrt{\frac{n-1}{n}} = \frac{\sqrt{n} - \sqrt{n-1}}{\sqrt{n}} = \frac{n - (n-1)}{\sqrt{n}(\sqrt{n} + \sqrt{n-1})} = \frac{1}{n + \sqrt{n^2 - n}} \\ &\geq \frac{1}{n + \sqrt{n^2}} = \frac{1}{2n}. \end{aligned}$$

The series $\sum_{n=2}^{\infty} \frac{1}{2n}$ diverges, so the series $\sum_{n=2}^{\infty} \left(1 - \sqrt{1 - \frac{1}{n}}\right)$ also diverges by the Comparison Test.

52. Determine whether $\sum_{n=2}^{\infty} \left(1 - \sqrt{1 - \frac{1}{n^2}}\right)$ converges.

SOLUTION

$$\begin{aligned} 1 - \sqrt{1 - \frac{1}{n^2}} &= 1 - \sqrt{\frac{n^2 - 1}{n^2}} = \frac{n - \sqrt{n^2 - 1}}{n} = \frac{n^2 - (n^2 - 1)}{n(n + \sqrt{n^2 - 1})} \\ &= \frac{1}{n(n + \sqrt{n^2 - 1})} = \frac{1}{n^2 + n\sqrt{n^2 - 1}} \leq \frac{1}{n^2} \end{aligned}$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series, so the series $\sum_{n=2}^{\infty} \left(1 - \sqrt{1 - \frac{1}{n^2}}\right)$ also converges by the Comparison Test.

53. Let $S = \sum_{n=1}^{\infty} \frac{n}{(n^2 + 1)^2}$.

(a) Show that S converges.

(b) $\square \square \square$ Use Eq. (4) in Exercise 83 of Section 10.3 with $M = 99$ to approximate S . What is the maximum size of the error?

SOLUTION

(a) For $n \geq 1$,

$$\frac{n}{(n^2 + 1)^2} < \frac{n}{(n^2)^2} = \frac{1}{n^3}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^3}$ is a convergent p -series, so the series $\sum_{n=1}^{\infty} \frac{n}{(n^2 + 1)^2}$ also converges by the Comparison Test.

(b) With $a_n = \frac{n}{(n^2+1)^2}$, $f(x) = \frac{x}{(x^2+1)^2}$ and $M = 99$, Eq. (4) in Exercise 83 of Section 10.3 becomes

$$\sum_{n=1}^{99} \frac{n}{(n^2+1)^2} + \int_{100}^{\infty} \frac{x}{(x^2+1)^2} dx \leq S \leq \sum_{n=1}^{100} \frac{n}{(n^2+1)^2} + \int_{100}^{\infty} \frac{x}{(x^2+1)^2} dx,$$

or

$$0 \leq S - \left(\sum_{n=1}^{99} \frac{n}{(n^2+1)^2} + \int_{100}^{\infty} \frac{x}{(x^2+1)^2} dx \right) \leq \frac{100}{(100^2+1)^2}.$$

Now,

$$\begin{aligned} \sum_{n=1}^{99} \frac{n}{(n^2+1)^2} &= 0.397066274; \text{ and} \\ \int_{100}^{\infty} \frac{x}{(x^2+1)^2} dx &= \lim_{R \rightarrow \infty} \int_{100}^R \frac{x}{(x^2+1)^2} dx = \frac{1}{2} \lim_{R \rightarrow \infty} \left(-\frac{1}{R^2+1} + \frac{1}{100^2+1} \right) \\ &= \frac{1}{20002} = 0.000049995; \end{aligned}$$

thus,

$$S \approx 0.397066274 + 0.000049995 = 0.397116269.$$

The bound on the error in this approximation is

$$\frac{100}{(100^2+1)^2} = 9.998 \times 10^{-7}.$$

In Exercises 54–57, determine whether the series converges absolutely. If it does not, determine whether it converges conditionally.

$$54. \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt[3]{n} + 2n}$$

SOLUTION Both $\sqrt[3]{n}$ and $2n$ are increasing functions, so $\sqrt[3]{n} + 2n$ is also increasing. Therefore, $\frac{1}{\sqrt[3]{n} + 2n}$ is decreasing. Moreover,

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt[3]{n} + 2n} = 0,$$

so the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt[3]{n} + 2n}$ converges by the Leibniz Test.

The corresponding positive series is $\sum_{n=1}^{\infty} \frac{1}{\sqrt[3]{n} + 2n}$. Because

$$\frac{1}{\sqrt[3]{n} + 2n} > \frac{1}{n + 2n} = \frac{1}{3} \cdot \frac{1}{n}$$

and the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges, $\sum_{n=1}^{\infty} \frac{1}{\sqrt[3]{n} + 2n}$ also diverges by the Comparison Test. Thus, $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt[3]{n} + 2n}$ converges conditionally.

$$55. \sum_{n=1}^{\infty} \frac{(-1)^n}{n^{1.1} \ln(n+1)}$$

SOLUTION Consider the corresponding positive series $\sum_{n=1}^{\infty} \frac{1}{n^{1.1} \ln(n+1)}$. Because

$$\frac{1}{n^{1.1} \ln(n+1)} < \frac{1}{n^{1.1}}$$

and $\sum_{n=1}^{\infty} \frac{1}{n^{1.1}}$ is a convergent p -series, we can conclude by the Comparison Test that $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^{1.1} \ln(n+1)}$ also converges.

Thus, $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^{1.1} \ln(n+1)}$ converges absolutely.

$$56. \sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + \pi n\right)}{\sqrt{n}}$$

SOLUTION Note

$$\cos\left(\frac{\pi}{4} + \pi n\right) = \cos \frac{\pi}{4} \cos n\pi - \sin \frac{\pi}{4} \sin n\pi = (-1)^n \frac{\sqrt{2}}{2}.$$

Therefore,

$$\sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + \pi n\right)}{\sqrt{n}} = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}} \frac{2}{\sqrt{2}} = \frac{2}{\sqrt{2}} \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}.$$

Now, the sequence $\left\{\frac{1}{\sqrt{n}}\right\}$ is decreasing and converges to 0 as $n \rightarrow \infty$. Therefore, $\sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + \pi n\right)}{\sqrt{n}}$ converges by the

Leibniz Test. However, the corresponding positive series is a divergent p -series ($p = \frac{1}{2}$), so the original series converges conditionally.

$$57. \sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + 2\pi n\right)}{\sqrt{n}}$$

SOLUTION $\cos\left(\frac{\pi}{4} + 2\pi n\right) = \cos \frac{\pi}{4} = \frac{\sqrt{2}}{2}$, so

$$\sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + 2\pi n\right)}{\sqrt{n}} = \frac{\sqrt{2}}{2} \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}.$$

This is a divergent p -series, so the series $\sum_{n=1}^{\infty} \frac{\cos\left(\frac{\pi}{4} + 2\pi n\right)}{\sqrt{n}}$ diverges.

58. **CAS** Use a computer algebra system to approximate $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 + \sqrt{n}}$ to within an error of at most 10^{-5} .

SOLUTION The sequence $\left\{\frac{1}{n^3 + \sqrt{n}}\right\}$ is decreasing and converges to 0, so the series $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 + \sqrt{n}}$ converges by the Leibniz Test. Using the error bound for an alternating series,

$$\left| S_N - \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 + \sqrt{n}} \right| \leq a_{N+1} = \frac{1}{(N+1)^3 + \sqrt{N+1}}.$$

If we want an approximation with an error of at most 10^{-5} , we must choose N such that

$$\frac{1}{(N+1)^3 + \sqrt{N+1}} < 10^{-5} \quad \text{or} \quad (N+1)^3 + \sqrt{N+1} > 10^5.$$

For $N = 45$, $(N+1)^3 + \sqrt{N+1} = 97,342.8 < 10^5$, and for $N = 46$, $(N+1)^3 + \sqrt{N+1} = 103,829.9 > 10^5$. The smallest acceptable value for N is therefore $N = 46$. Using a computer algebra system, we find

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n^3 + \sqrt{n}} \approx S_{46} = -0.418452236.$$

59. Catalan's constant is defined by $K = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}$.

(a) How many terms of the series are needed to calculate K with an error of less than 10^{-6} ?

(b) **CAS** Carry out the calculation.

SOLUTION Using the error bound for an alternating series, we have

$$|S_N - K| \leq \frac{1}{(2(N+1)+1)^2} = \frac{1}{(2N+3)^2}.$$

For accuracy to three decimal places, we must choose N so that

$$\frac{1}{(2N+3)^2} < 5 \times 10^{-3} \quad \text{or} \quad (2N+3)^2 > 2000.$$

Solving for N yields

$$N > \frac{1}{2}(\sqrt{2000} - 3) \approx 20.9.$$

Thus,

$$K \approx \sum_{k=0}^{21} \frac{(-1)^k}{(2k+1)^2} = 0.915707728.$$

60. Give an example of conditionally convergent series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ such that $\sum_{n=1}^{\infty} (a_n + b_n)$ converges absolutely.

SOLUTION Let $a_n = \frac{(-1)^n}{n}$ and $b_n = \frac{(-1)^{n+1}}{n}$. The corresponding alternating series converge by the Leibniz Test; however, the corresponding positive series are the divergent harmonic series. Thus, $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ converge conditionally.

On the other hand, the series

$$\sum_{n=1}^{\infty} (a_n + b_n) = \sum_{n=1}^{\infty} \left(\frac{(-1)^n}{n} + \frac{(-1)^{n+1}}{n} \right) = \sum_{n=1}^{\infty} (-1)^n \left(\frac{1}{n} + \frac{-1}{n} \right) = \sum_{n=1}^{\infty} 0$$

converges absolutely.

61. Let $\sum_{n=1}^{\infty} a_n$ be an absolutely convergent series. Determine whether the following series are convergent or divergent:

(a) $\sum_{n=1}^{\infty} \left(a_n + \frac{1}{n^2} \right)$

(b) $\sum_{n=1}^{\infty} (-1)^n a_n$

(c) $\sum_{n=1}^{\infty} \frac{1}{1+a_n^2}$

(d) $\sum_{n=1}^{\infty} \frac{|a_n|}{n}$

SOLUTION Because $\sum_{n=1}^{\infty} a_n$ converges absolutely, we know that $\sum_{n=1}^{\infty} a_n$ converges and that $\sum_{n=1}^{\infty} |a_n|$ converges.

(a) Because we know that $\sum_{n=1}^{\infty} a_n$ converges and the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series, the sum of these two series,

$$\sum_{n=1}^{\infty} \left(a_n + \frac{1}{n^2} \right) \text{ also converges.}$$

(b) We have,

$$\sum_{n=1}^{\infty} |(-1)^n a_n| = \sum_{n=1}^{\infty} |a_n|$$

Because $\sum_{n=1}^{\infty} |a_n|$ converges, it follows that $\sum_{n=1}^{\infty} (-1)^n a_n$ converges absolutely, which implies that $\sum_{n=1}^{\infty} (-1)^n a_n$ converges.

(c) Because $\sum_{n=1}^{\infty} a_n$ converges, $\lim_{n \rightarrow \infty} a_n = 0$. Therefore,

$$\lim_{n \rightarrow \infty} \frac{1}{1+a_n^2} = \frac{1}{1+0^2} = 1 \neq 0,$$

and the series $\sum_{n=1}^{\infty} \frac{1}{1+a_n^2}$ diverges by the Divergence Test.

(d) $\frac{|a_n|}{n} \leq |a_n|$ and the series $\sum_{n=1}^{\infty} |a_n|$ converges, so the series $\sum_{n=1}^{\infty} \frac{|a_n|}{n}$ also converges by the Comparison Test.

62. Let $\{a_n\}$ be a positive sequence such that $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \frac{1}{2}$. Determine whether the following series converge or diverge:

(a) $\sum_{n=1}^{\infty} 2a_n$

(b) $\sum_{n=1}^{\infty} 3^n a_n$

(c) $\sum_{n=1}^{\infty} \sqrt{a_n}$

SOLUTION

(a)

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{2a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{2} \sqrt[n]{a_n} = 1 \cdot \frac{1}{2} = \frac{1}{2}.$$

Because $L < 1$, the series converges by the Root Test.

(b)

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{3^n a_n} = \lim_{n \rightarrow \infty} 3 \sqrt[n]{a_n} = 3 \cdot \frac{1}{2} = \frac{3}{2}.$$

Because $L > 1$, the series diverges by the Root Test.

(c)

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{\sqrt{a_n}} = \lim_{n \rightarrow \infty} \sqrt{\sqrt[n]{a_n}} = \sqrt{\frac{1}{2}}.$$

Because $L < 1$, the series converges by the Root Test.

In Exercises 63–70, apply the Ratio Test to determine convergence or divergence, or state that the Ratio Test is inconclusive.

63. $\sum_{n=1}^{\infty} \frac{n^5}{5^n}$

SOLUTION With $a_n = \frac{n^5}{5^n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^5}{5^{n+1}} \cdot \frac{5^n}{n^5} = \frac{1}{5} \left(1 + \frac{1}{n} \right)^5,$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{5} \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^5 = \frac{1}{5} \cdot 1 = \frac{1}{5}.$$

Because $\rho < 1$, the series converges by the Ratio Test.

64. $\sum_{n=1}^{\infty} \frac{\sqrt{n+1}}{n^8}$

SOLUTION With $a_n = \frac{\sqrt{n+1}}{n^8}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{\sqrt{n+2}}{(n+1)^8} \cdot \frac{n^8}{\sqrt{n+1}} = \sqrt{\frac{n+2}{n+1}} \left(\frac{n}{n+1} \right)^8,$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot 1^8 = 1.$$

Because $\rho = 1$, the Ratio Test is inconclusive.

65. $\sum_{n=1}^{\infty} \frac{1}{n2^n + n^3}$

SOLUTION With $a_n = \frac{1}{n2^n + n^3}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{n2^n + n^3}{(n+1)2^{n+1} + (n+1)^3} = \frac{n2^n \left(1 + \frac{n^2}{2^n} \right)}{(n+1)2^{n+1} \left(1 + \frac{(n+1)^2}{2^{n+1}} \right)} = \frac{1}{2} \cdot \frac{n}{n+1} \cdot \frac{1 + \frac{n^2}{2^n}}{1 + \frac{(n+1)^2}{2^{n+1}}},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2} \cdot 1 \cdot 1 = \frac{1}{2}.$$

Because $\rho < 1$, the series converges by the Ratio Test.

$$66. \sum_{n=1}^{\infty} \frac{n^4}{n!}$$

SOLUTION With $a_n = \frac{n^4}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^4}{(n+1)!} \cdot \frac{n!}{n^4} = \frac{(n+1)^3}{n^4} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 0.$$

Because $\rho < 1$, the series converges by the Ratio Test.

$$67. \sum_{n=1}^{\infty} \frac{2^{n^2}}{n!}$$

SOLUTION With $a_n = \frac{2^{n^2}}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{(n+1)^2}}{(n+1)!} \cdot \frac{n!}{2^{n^2}} = \frac{2^{2n+1}}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty.$$

Because $\rho > 1$, the series diverges by the Ratio Test.

$$68. \sum_{n=4}^{\infty} \frac{\ln n}{n^{3/2}}$$

SOLUTION With $a_n = \frac{\ln n}{n^{3/2}}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{\ln(n+1)}{(n+1)^{3/2}} \cdot \frac{n^{3/2}}{\ln n} = \left(\frac{n}{n+1} \right)^{3/2} \frac{\ln(n+1)}{\ln n},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1^{3/2} \cdot 1 = 1.$$

Because $\rho = 1$, the Ratio Test is inconclusive.

$$69. \sum_{n=1}^{\infty} \left(\frac{n}{2} \right)^n \frac{1}{n!}$$

SOLUTION With $a_n = \left(\frac{n}{2} \right)^n \frac{1}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \left(\frac{n+1}{2} \right)^{n+1} \frac{1}{(n+1)!} \cdot \left(\frac{2}{n} \right)^n n! = \frac{1}{2} \left(\frac{n+1}{n} \right)^n = \frac{1}{2} \left(1 + \frac{1}{n} \right)^n,$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{2}e.$$

Because $\rho = \frac{e}{2} > 1$, the series diverges by the Ratio Test.

$$70. \sum_{n=1}^{\infty} \left(\frac{n}{4} \right)^n \frac{1}{n!}$$

SOLUTION With $a_n = \left(\frac{n}{4} \right)^n \frac{1}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \left(\frac{n+1}{4} \right)^{n+1} \frac{1}{(n+1)!} \cdot \left(\frac{4}{n} \right)^n n! = \frac{1}{4} \left(\frac{n+1}{n} \right)^n = \frac{1}{4} \left(1 + \frac{1}{n} \right)^n,$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}e.$$

Because $\rho = \frac{e}{4} < 1$, the series converges by the Ratio Test.

In Exercises 71–74, apply the Root Test to determine convergence or divergence, or state that the Root Test is inconclusive.

$$71. \sum_{n=1}^{\infty} \frac{1}{4^n}$$

SOLUTION With $a_n = \frac{1}{4^n}$,

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\frac{1}{4^n}} = \frac{1}{4}.$$

Because $L < 1$, the series converges by the Root Test.

$$72. \sum_{n=1}^{\infty} \left(\frac{2}{n}\right)^n$$

SOLUTION With $a_n = \left(\frac{2}{n}\right)^n$,

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{\left(\frac{2}{n}\right)^n} = \lim_{n \rightarrow \infty} \frac{2}{n} = 0.$$

Because $L < 1$, the series converges by the Root Test.

$$73. \sum_{n=1}^{\infty} \left(\frac{3}{4n}\right)^n$$

SOLUTION With $a_n = \left(\frac{3}{4n}\right)^n$,

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\left(\frac{3}{4n}\right)^n} = \lim_{n \rightarrow \infty} \frac{3}{4n} = 0.$$

Because $L < 1$, the series converges by the Root Test.

$$74. \sum_{n=1}^{\infty} \left(\cos \frac{1}{n}\right)^{n^3}$$

SOLUTION With $a_n = \left(\cos \frac{1}{n}\right)^{n^3}$,

$$L = \lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\cos \left(\frac{1}{n}\right)^{n^3}} = \lim_{n \rightarrow \infty} \cos \left(\frac{1}{n}\right)^{n^2} = \lim_{x \rightarrow \infty} \cos \left(\frac{1}{x}\right)^{x^2}.$$

Now,

$$\begin{aligned} \ln L &= \lim_{x \rightarrow \infty} x^2 \ln \cos \left(\frac{1}{x}\right) = \lim_{x \rightarrow \infty} \frac{\ln \cos \left(\frac{1}{x}\right)}{\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{\frac{1}{\cos \left(\frac{1}{x}\right)} \left(-\sin \left(\frac{1}{x}\right)\right) \left(-\frac{1}{x^2}\right)}{-\frac{2}{x^3}} \\ &= -\frac{1}{2} \lim_{x \rightarrow \infty} \frac{1}{\cos \left(\frac{1}{x}\right)} \cdot \lim_{x \rightarrow \infty} \frac{\sin \left(\frac{1}{x}\right)}{\frac{1}{x}} = -\frac{1}{2} \cdot 1 \cdot 1 = -\frac{1}{2}. \end{aligned}$$

Therefore, $L = e^{-1/2}$. Because $L < 1$, the series converges by the Root Test.

In Exercises 75–92, determine convergence or divergence using any method covered in the text.

$$75. \sum_{n=1}^{\infty} \left(\frac{2}{3}\right)^n$$

SOLUTION This is a geometric series with ratio $r = \frac{2}{3} < 1$; hence, the series converges.

$$76. \sum_{n=1}^{\infty} \frac{\pi^{7n}}{e^{8n}}$$

SOLUTION This is a geometric series with ratio $r = \frac{\pi^7}{e^8} \approx 1.013$, so it diverges.

$$77. \sum_{n=1}^{\infty} e^{-0.02n}$$

SOLUTION This is a geometric series with common ratio $r = \frac{1}{e^{0.02}} \approx 0.98 < 1$; hence, the series converges.

$$78. \sum_{n=1}^{\infty} ne^{-0.02n}$$

SOLUTION With $a_n = ne^{-0.02n}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)e^{-0.02(n+1)}}{ne^{-0.02n}} = \frac{n+1}{n} e^{-0.02},$$

and

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1 \cdot e^{-0.02} = e^{-0.02}.$$

Because $\rho < 1$, the series converges by the Ratio Test.

$$79. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n} + \sqrt{n+1}}$$

SOLUTION In this alternating series, $a_n = \frac{1}{\sqrt{n} + \sqrt{n+1}}$. The sequence $\{a_n\}$ is decreasing, and

$$\lim_{n \rightarrow \infty} a_n = 0;$$

therefore the series converges by the Leibniz Test.

$$80. \sum_{n=10}^{\infty} \frac{1}{n(\ln n)^{3/2}}$$

SOLUTION Let $f(x) = \frac{1}{x(\ln x)^{3/2}}$. This function is continuous, positive and decreasing for $x > e^{-3/2}$ and thus for $x \geq 10$; therefore, the Integral Test applies. Now,

$$\begin{aligned} \int_{10}^{\infty} \frac{dx}{x(\ln x)^{3/2}} &= \lim_{R \rightarrow \infty} \int_{10}^R \frac{dx}{x(\ln x)^{3/2}} = \lim_{R \rightarrow \infty} \int_{\ln 10}^{\ln R} \frac{1}{u^{3/2}} du \\ &= \lim_{R \rightarrow \infty} \left(\frac{-2}{\sqrt{u}} \Big|_{\ln 10}^{\ln R} \right) = 2 \lim_{R \rightarrow \infty} \left(\frac{1}{\sqrt{\ln 10}} - \frac{1}{\sqrt{\ln R}} \right) = 2. \end{aligned}$$

The integral converges; hence, the series converges as well.

$$81. \sum_{n=2}^{\infty} \frac{(-1)^n}{\ln n}$$

SOLUTION The sequence $a_n = \frac{1}{\ln n}$ is decreasing for $n \geq 10$ and

$$\lim_{n \rightarrow \infty} a_n = 0;$$

therefore, the series converges by the Leibniz Test.

$$82. \sum_{n=1}^{\infty} \frac{e^n}{n!}$$

SOLUTION With $a_n = \frac{e^n}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{e^{n+1}}{(n+1)!} \cdot \frac{n!}{e^n} = \frac{e}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0.$$

Because $\rho < 1$, the series converges by the Ratio Test.

$$83. \sum_{n=1}^{\infty} \frac{1}{n\sqrt{n} + \ln n}$$

SOLUTION For $n \geq 1$,

$$\frac{1}{n\sqrt{n} + \ln n} \leq \frac{1}{n\sqrt{n}} = \frac{1}{n^{3/2}}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$ is a convergent p -series, so the series $\sum_{n=1}^{\infty} \frac{1}{n\sqrt{n} + \ln n}$ converges by the Comparison Test.

$$84. \sum_{n=1}^{\infty} \frac{1}{\sqrt[3]{n}(1 + \sqrt{n})}$$

SOLUTION Apply the Limit Comparison Test with $a_n = \frac{1}{\sqrt[3]{n}(1 + \sqrt{n})}$ and $b_n = \frac{1}{n^{5/6}}$. Then,

$$L = \lim_{n \rightarrow \infty} \frac{\frac{1}{\sqrt[3]{n}(1 + \sqrt{n})}}{\frac{1}{n^{5/6}}} = \lim_{n \rightarrow \infty} \frac{n^{5/6}}{\sqrt[3]{n} + n^{5/6}} = \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{\sqrt{n}} + 1} = 1.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^{5/6}}$ is a divergent p -series. Because $L > 0$, the series $\sum_{n=1}^{\infty} \frac{1}{\sqrt[3]{n}(1 + \sqrt{n})}$ also diverges by the Limit Comparison Test.

$$85. \sum_{n=1}^{\infty} \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}} \right)$$

SOLUTION This series telescopes:

$$\sum_{n=1}^{\infty} \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}} \right) = \left(1 - \frac{1}{\sqrt{2}} \right) + \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{3}} \right) + \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{4}} \right) + \dots$$

so that the n^{th} partial sum S_n is

$$S_n = \left(1 - \frac{1}{\sqrt{2}} \right) + \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{3}} \right) + \left(\frac{1}{\sqrt{3}} - \frac{1}{\sqrt{4}} \right) + \dots + \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}} \right) = 1 - \frac{1}{\sqrt{n+1}}$$

and then

$$\sum_{n=1}^{\infty} \left(\frac{1}{\sqrt{n}} - \frac{1}{\sqrt{n+1}} \right) = \lim_{n \rightarrow \infty} S_n = 1 - \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n+1}} = 1$$

$$86. \sum_{n=1}^{\infty} (\ln n - \ln(n+1))$$

SOLUTION This series telescopes:

$$\sum_{n=1}^{\infty} (\ln n - \ln(n+1)) = (\ln 1 - \ln 2) + (\ln 2 - \ln 3) + (\ln 3 - \ln 4) + \dots$$

so that the n^{th} partial sum S_n is

$$\begin{aligned} S_n &= (\ln 1 - \ln 2) + (\ln 2 - \ln 3) + (\ln 3 - \ln 4) + \dots + (\ln n - \ln(n+1)) \\ &= \ln 1 - \ln(n+1) = -\ln(n+1) \end{aligned}$$

and then

$$\sum_{n=1}^{\infty} (\ln n - \ln(n+1)) = \lim_{n \rightarrow \infty} S_n = -\lim_{n \rightarrow \infty} \ln(n+1) = \infty$$

so the series diverges.

$$87. \sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}}$$

SOLUTION For $n \geq 1$, $\sqrt{n} \leq n$, so that

$$\sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}} \geq \sum_{n=1}^{\infty} \frac{1}{2n}$$

which diverges since it is a constant multiple of the harmonic series. Thus $\sum_{n=1}^{\infty} \frac{1}{n + \sqrt{n}}$ diverges as well, by the Comparison Test.

$$88. \sum_{n=2}^{\infty} \frac{\cos(\pi n)}{n^{2/3}}$$

SOLUTION $\cos(\pi n) = (-1)^n$, so

$$\sum_{n=2}^{\infty} \frac{\cos(\pi n)}{n^{2/3}} = \sum_{n=2}^{\infty} \frac{(-1)^n}{n^{2/3}}.$$

The sequence $a_n = \frac{1}{n^{2/3}}$ is decreasing and

$$\lim_{n \rightarrow \infty} a_n = 0;$$

therefore, the series converges by the Leibniz Test.

$$89. \sum_{n=2}^{\infty} \frac{1}{n^{\ln n}}$$

SOLUTION For $n \geq N$ large enough, $\ln n \geq 2$ so that

$$\sum_{n=N}^{\infty} \frac{1}{n^{\ln n}} \leq \sum_{n=N}^{\infty} \frac{1}{n^2}$$

which is a convergent p -series. Thus by the Comparison Test, $\sum_{n=N}^{\infty} \frac{1}{n^{\ln n}}$ also converges; adding back in the terms for $n < N$ does not affect convergence.

$$90. \sum_{n=2}^{\infty} \frac{1}{\ln^3 n}$$

SOLUTION For N large enough, $\ln n \leq n^{1/4}$ when $n \geq N$ so that

$$\sum_{n=N}^{\infty} \frac{1}{\ln^3 n} > \sum_{n=N}^{\infty} \frac{1}{n^{3/4}}$$

which is a divergent p -series. Thus by the Comparison Test, $\sum_{n=N}^{\infty} \frac{1}{\ln^3 n}$ diverges; adding back in the terms for $n < N$ does not affect this result.

$$91. \sum_{n=1}^{\infty} \sin^2 \frac{\pi}{n}$$

SOLUTION For all $x > 0$, $\sin x < x$. Therefore, $\sin^2 x < x^2$, and for $x = \frac{\pi}{n}$,

$$\sin^2 \frac{\pi}{n} < \frac{\pi^2}{n^2} = \pi^2 \cdot \frac{1}{n^2}.$$

The series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series, so the series $\sum_{n=1}^{\infty} \sin^2 \frac{\pi}{n}$ also converges by the Comparison Test.

$$92. \sum_{n=0}^{\infty} \frac{2^{2n}}{n!}$$

SOLUTION With $a_n = \frac{2^{2n}}{n!}$,

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{2^{2(n+1)}}{(n+1)!} \cdot \frac{n!}{2^{2n}} = \frac{4}{n+1} \quad \text{and} \quad \rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 0.$$

Because $\rho < 1$, the series converges by the Ratio Test.

In Exercises 93–98, find the interval of convergence of the power series.

$$93. \sum_{n=0}^{\infty} \frac{2^n x^n}{n!}$$

SOLUTION With $a_n = \frac{2^n x^n}{n!}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{2^{n+1} x^{n+1}}{(n+1)!} \cdot \frac{n!}{2^n x^n} \right| = \lim_{n \rightarrow \infty} \left| x \cdot \frac{2}{n} \right| = 0$$

Then $\rho < 1$ for all x , so that the radius of convergence is $R = \infty$, and the series converges for all x .

$$94. \sum_{n=0}^{\infty} \frac{x^n}{n+1}$$

SOLUTION With $a_n = \frac{x^n}{n+1}$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{n+2} \cdot \frac{n+1}{x^n} \right| = \lim_{n \rightarrow \infty} \left| x \cdot \frac{n+1}{n+2} \right| = \lim_{n \rightarrow \infty} \left| x \cdot \frac{1+1/n}{1+2/n} \right| = |x|$$

Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is 1, and the series converges absolutely for $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=0}^{\infty} \frac{1}{n+1} = \sum_{n=1}^{\infty} \frac{1}{n}$, which is the divergent harmonic series.

For the endpoint $x = -1$, the series becomes $\sum_{n=0}^{\infty} \frac{(-1)^n}{n+1}$, which converges by the Leibniz Test. The series $\sum_{n=0}^{\infty} \frac{x^n}{n+1}$ therefore converges for $-1 \leq x < 1$.

$$95. \sum_{n=0}^{\infty} \frac{n^6}{n^8 + 1} (x-3)^n$$

SOLUTION With $a_n = \frac{n^6(x-3)^n}{n^8+1}$,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)^6(x-3)^{n+1}}{(n+1)^8-1} \cdot \frac{n^8+1}{n^6(x-3)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| (x-3) \cdot \frac{(n+1)^6(n^8+1)}{n^6((n+1)^8+1)} \right| \\ &= \lim_{n \rightarrow \infty} \left| (x-3) \cdot \frac{n^{14} + \text{terms of lower degree}}{n^{14} + \text{terms of lower degree}} \right| = |x-3| \end{aligned}$$

Then $\rho < 1$ when $|x-3| < 1$, so the radius of convergence is 1, and the series converges absolutely for $|x-3| < 1$, or $2 < x < 4$. For the endpoint $x = 4$, the series becomes $\sum_{n=0}^{\infty} \frac{n^6}{n^8+1}$, which converges by the Comparison Test comparing

with the convergent p -series $\sum_{n=1}^{\infty} \frac{1}{n^2}$. For the endpoint $x = 2$, the series becomes $\sum_{n=0}^{\infty} \frac{n^6(-1)^n}{n^8+1}$, which converges by the

Leibniz Test. The series $\sum_{n=0}^{\infty} \frac{n^6(x-3)^n}{n^8+1}$ therefore converges for $2 \leq x \leq 4$.

$$96. \sum_{n=0}^{\infty} nx^n$$

SOLUTION With $a_n = nx^n$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)x^{n+1}}{nx^n} \right| = \lim_{n \rightarrow \infty} \left| x \cdot \frac{n+1}{n} \right| = |x|$$

Then $\rho < 1$ when $|x| < 1$, so the radius of convergence is 1, and the series converges for $|x| < 1$, or $-1 < x < 1$. For the endpoint $x = 1$, the series becomes $\sum_{n=0}^{\infty} n$, which diverges by the Divergence Test. For the endpoint $x = -1$, the series becomes $\sum_{n=0}^{\infty} (-1)^n n$, which also diverges by the Divergence Test. The series $\sum_{n=0}^{\infty} nx^n$ therefore converges for

$-1 < x < 1$.

$$97. \sum_{n=0}^{\infty} (nx)^n$$

SOLUTION With $a_n = n^n x^n$, and assuming $x \neq 0$,

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n+1)^{n+1} x^{n+1}}{n^n x^n} \right| = \lim_{n \rightarrow \infty} \left| x(n+1) \cdot \left(\frac{n+1}{n} \right)^n \right| = \infty$$

since $\left(\frac{n+1}{n} \right)^n = \left(1 + \frac{1}{n} \right)^n$ converges to e and the $(n+1)$ term diverges to ∞ . Thus $\rho < 1$ only when $x = 0$, so the series converges only for $x = 0$.

$$98. \sum_{n=0}^{\infty} \frac{(2x-3)^n}{n \ln n}$$

SOLUTION With $a_n = \frac{(2x-3)^n}{n \ln n}$, and using L'Hôpital's Rule,

$$\begin{aligned} \rho &= \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(2x-3)^{n+1}}{(n+1) \ln(n+1)} \cdot \frac{n \ln n}{(2x-3)^n} \right| \\ &= \lim_{n \rightarrow \infty} \left| (2x-3) \frac{n \ln n}{(n+1) \ln(n+1)} \right| = \lim_{n \rightarrow \infty} \left| (2x-3) \frac{1 + \ln n}{1 + \ln(n+1)} \right| \\ &= \lim_{n \rightarrow \infty} \left| (2x-3) \frac{1/n}{1/(n+1)} \right| = \lim_{n \rightarrow \infty} \left| (2x-3) \frac{n+1}{n} \right| = |2x-3| \end{aligned}$$

Then $\rho < 1$ when $|2x-3| < 1$, so the radius of convergence is 1, and the series converges absolutely for $|2x-3| < 1$, or $1 < x < 2$. For the endpoint $x = 2$, the series becomes $\sum_{n=0}^{\infty} \frac{1}{n \ln n}$, which diverges by the Integral Test. For the endpoint

$x = -1$, the series becomes $\sum_{n=0}^{\infty} \frac{(-1)^n}{n \ln n}$, which converges by the Leibniz Test. The series $\sum_{n=0}^{\infty} \frac{(2x-3)^n}{n \ln n}$ therefore converges for $1 \leq x < 2$.

99. Expand $f(x) = \frac{2}{4-3x}$ as a power series centered at $c = 0$. Determine the values of x for which the series converges.

SOLUTION Write

$$\frac{2}{4-3x} = \frac{1}{2} \frac{1}{1 - \frac{3}{4}x}$$

Substituting $\frac{3}{4}x$ for x in the Maclaurin series for $\frac{1}{1-x}$, we obtain

$$\frac{1}{1 - \frac{3}{4}x} = \sum_{n=0}^{\infty} \left(\frac{3}{4} \right)^n x^n$$

This series converges for $\left| \frac{3}{4}x \right| < 1$, or $|x| < \frac{4}{3}$. Hence, for $|x| < \frac{4}{3}$,

$$\frac{2}{4-3x} = \frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{3}{4} \right)^n x^n$$

100. Prove that

$$\sum_{n=0}^{\infty} n e^{-nx} = \frac{e^{-x}}{(1 - e^{-x})^2}$$

Hint: Express the left-hand side as the derivative of a geometric series.

SOLUTION For $x > 0$, $\sum_{n=0}^{\infty} e^{-nx} = \sum_{n=0}^{\infty} (e^{-x})^n$ is a convergent geometric series with ratio $r = e^{-x}$; hence,

$$\sum_{n=0}^{\infty} e^{-nx} = \frac{1}{1 - e^{-x}}.$$

Differentiating term-by-term then yields

$$\sum_{n=0}^{\infty} (-n e^{-nx}) = -\frac{e^{-x}}{(1 - e^{-x})^2}.$$

Therefore, for $x > 0$,

$$\sum_{n=0}^{\infty} n e^{-nx} = \frac{e^{-x}}{(1 - e^{-x})^2}.$$

101. Let $F(x) = \sum_{k=0}^{\infty} \frac{x^{2k}}{2^k \cdot k!}$.

(a) Show that $F(x)$ has infinite radius of convergence.

(b) Show that $y = F(x)$ is a solution of

$$y'' = xy' + y, \quad y(0) = 1, \quad y'(0) = 0$$

(c)  Plot the partial sums S_N for $N = 1, 3, 5, 7$ on the same set of axes.

SOLUTION

(a) With $a_k = \frac{x^{2k}}{2^k \cdot k!}$,

$$\left| \frac{a_{k+1}}{a_k} \right| = \frac{|x|^{2k+2}}{2^{k+1} \cdot (k+1)!} \cdot \frac{2^k \cdot k!}{|x|^{2k}} = \frac{x^2}{2(k+1)},$$

and

$$\rho = \lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| = x^2 \cdot 0 = 0.$$

Because $\rho < 1$ for all x , we conclude that the series converges for all x ; that is, $R = \infty$.

(b) Let

$$y = F(x) = \sum_{k=0}^{\infty} \frac{x^{2k}}{2^k \cdot k!}.$$

Then

$$y' = \sum_{k=1}^{\infty} \frac{2kx^{2k-1}}{2^k k!} = \sum_{k=1}^{\infty} \frac{x^{2k-1}}{2^{k-1}(k-1)!},$$

$$y'' = \sum_{k=1}^{\infty} \frac{(2k-1)x^{2k-2}}{2^{k-1}(k-1)!},$$

and


$$\begin{aligned} xy' + y &= x \sum_{k=1}^{\infty} \frac{x^{2k-1}}{2^{k-1}(k-1)!} + \sum_{k=0}^{\infty} \frac{x^{2k}}{2^k k!} = \sum_{k=1}^{\infty} \frac{x^{2k}}{2^{k-1}(k-1)!} + 1 + \sum_{k=1}^{\infty} \frac{x^{2k}}{2^k k!} \\ &= 1 + \sum_{k=1}^{\infty} \frac{(2k+1)x^{2k}}{2^k k!} = \sum_{k=0}^{\infty} \frac{(2k+1)x^{2k}}{2^k k!} = \sum_{k=1}^{\infty} \frac{(2k-1)x^{2k-2}}{2^{k-1}(k-1)!} = y''. \end{aligned}$$

Moreover,

$$y(0) = 1 + \sum_{k=1}^{\infty} \frac{0^{2k}}{2^k k!} = 1 \quad \text{and} \quad y'(0) = \sum_{k=1}^{\infty} \frac{0^{2k-1}}{2^{k-1}(k-1)!} = 0.$$

Thus, $\sum_{k=0}^{\infty} \frac{x^{2k}}{2^k k!}$ is the solution to the equation $y'' = xy' + y$ satisfying $y(0) = 1$, $y'(0) = 0$.

(c) The partial sums S_1 , S_3 , S_5 and S_7 are plotted in the figure below.



102. Find a power series $P(x) = \sum_{n=0}^{\infty} a_n x^n$ that satisfies the Laguerre differential equation

$$xy'' + (1-x)y' - y = 0$$

with initial condition satisfying $P(0) = 1$.

SOLUTION Let

$$y = P(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Then,

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}, \quad y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2},$$

and

$$\begin{aligned} xy'' + (1-x)y' - y &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-1} + \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n \\ &= \sum_{n=1}^{\infty} (n+1) n a_{n+1} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n \\ &= (a_1 - a_0) + \sum_{n=1}^{\infty} [(n+1)^2 a_{n+1} - (n+1) a_n] x^n. \end{aligned}$$

In order for this series to be equal to zero, the coefficient of x^n must be equal to zero for each n ; thus

$$a_1 = a_0 \quad \text{and} \quad a_{n+1} = \frac{a_n}{n+1}.$$

Now, $y(0) = P(0) = a_0$, so to satisfy the initial condition $P(0) = 1$, we must set $a_0 = 1$. Then,

$$\begin{aligned} a_1 &= a_0 = 1; \\ a_2 &= \frac{a_1}{2} = \frac{1}{2}; \\ a_3 &= \frac{a_2}{3} = \frac{1}{6} = \frac{1}{3!}; \\ a_4 &= \frac{a_3}{4} = \frac{1}{4!}; \end{aligned}$$

and, in general, $a_n = \frac{1}{n!}$. Thus,

$$P(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x.$$

In Exercises 103–112, find the Taylor series centered at c .

103. $f(x) = e^{4x}$, $c = 0$

SOLUTION Substituting $4x$ for x in the Maclaurin series for e^x yields

$$e^{4x} = \sum_{n=0}^{\infty} \frac{(4x)^n}{n!} = \sum_{n=0}^{\infty} \frac{4^n}{n!} x^n.$$

104. $f(x) = e^{2x}$, $c = -1$

SOLUTION Write:

$$e^{2x} = e^{2(x+1)-2} = e^{-2} e^{2(x+1)}.$$

Substituting $2(x+1)$ for x in the Maclaurin series for e^x yields

$$e^{2(x+1)} = \sum_{n=0}^{\infty} \frac{(2(x+1))^n}{n!} = \sum_{n=0}^{\infty} \frac{2^n}{n!} (x+1)^n;$$

hence,

$$e^{2x} = e^{-2} \sum_{n=0}^{\infty} \frac{2^n (x+1)^n}{n!}.$$

105. $f(x) = x^4$, $c = 2$

SOLUTION We have

$$f'(x) = 4x^3 \quad f''(x) = 12x^2 \quad f'''(x) = 24x \quad f^{(4)}(x) = 24$$

and all higher derivatives are zero, so that

$$f(2) = 2^4 = 16 \quad f'(2) = 4 \cdot 2^3 = 32 \quad f''(2) = 12 \cdot 2^2 = 48 \quad f'''(2) = 24 \cdot 2 = 48 \quad f^{(4)}(2) = 24$$

Thus the Taylor series centered at $c = 2$ is

$$\begin{aligned} \sum_{n=0}^4 \frac{f^{(n)}(2)}{n!} (x-2)^n &= 16 + \frac{32}{1!} (x-2) + \frac{48}{2!} (x-2)^2 + \frac{48}{3!} (x-2)^3 + \frac{24}{4!} (x-2)^4 \\ &= 16 + 32(x-2) + 24(x-2)^2 + 8(x-2)^3 + (x-2)^4 \end{aligned}$$

106. $f(x) = x^3 - x$, $c = -2$

SOLUTION We have

$$f'(x) = 3x^2 - 1 \quad f''(x) = 6x \quad f'''(x) = 6$$

and all higher derivatives are zero, so that

$$f(-2) = -8 + 2 = -6 \quad f'(-2) = 3(-2)^2 - 1 = 11 \quad f''(-2) = 6(-2) = -12 \quad f'''(-2) = 6$$

Thus the Taylor series centered at $c = -2$ is

$$\begin{aligned} \sum_{n=0}^3 \frac{f^{(n)}(-2)}{n!} (x+2)^n &= -6 + \frac{11}{1!} (x+2) + \frac{-12}{2!} (x+2)^2 + \frac{6}{3!} (x+2)^3 \\ &= -6 + 11(x+2) - 6(x+2)^2 + (x+2)^3 \end{aligned}$$

107. $f(x) = \sin x$, $c = \pi$

SOLUTION We have

$$f^{(4n)}(x) = \sin x \quad f^{(4n+1)}(x) = \cos x \quad f^{(4n+2)}(x) = -\sin x \quad f^{(4n+3)}(x) = -\cos x$$

so that

$$f^{(4n)}(\pi) = \sin \pi = 0 \quad f^{(4n+1)}(\pi) = \cos \pi = -1 \quad f^{(4n+2)}(\pi) = -\sin \pi = 0 \quad f^{(4n+3)}(\pi) = -\cos \pi = 1$$

Then the Taylor series centered at $c = \pi$ is

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{f^{(n)}(\pi)}{n!} (x - \pi)^n &= \frac{-1}{1!} (x - \pi) + \frac{1}{3!} (x - \pi)^3 + \frac{-1}{5!} (x - \pi)^5 + \frac{1}{7!} (x - \pi)^7 - \dots \\ &= -(x - \pi) + \frac{1}{6} (x - \pi)^3 - \frac{1}{120} (x - \pi)^5 + \frac{1}{5040} (x - \pi)^7 - \dots\end{aligned}$$

108. $f(x) = e^{x-1}$, $c = -1$

SOLUTION Write

$$e^{x-1} = e^{x+1-1-1} = e^{-2} e^{x+1}.$$

Substituting $x + 1$ for x in the Maclaurin series for e^x yields

$$e^{x+1} = \sum_{n=0}^{\infty} \frac{(x+1)^n}{n!};$$

hence,

$$e^{x-1} = e^{-2} \sum_{n=0}^{\infty} \frac{(x+1)^n}{n!} = \sum_{n=0}^{\infty} \frac{(x+1)^n}{n! e^2}.$$

109. $f(x) = \frac{1}{1-2x}$, $c = -2$

SOLUTION Write

$$\frac{1}{1-2x} = \frac{1}{5-2(x+2)} = \frac{1}{5} \frac{1}{1-\frac{2}{5}(x+2)}.$$

Substituting $\frac{2}{5}(x+2)$ for x in the Maclaurin series for $\frac{1}{1-x}$ yields

$$\frac{1}{1-\frac{2}{5}(x+2)} = \sum_{n=0}^{\infty} \frac{2^n}{5^n} (x+2)^n;$$

hence,

$$\frac{1}{1-2x} = \frac{1}{5} \sum_{n=0}^{\infty} \frac{2^n}{5^n} (x+2)^n = \sum_{n=0}^{\infty} \frac{2^n}{5^{n+1}} (x+2)^n.$$

110. $f(x) = \frac{1}{(1-2x)^2}$, $c = -2$

SOLUTION Note that

$$\frac{d}{dx} \frac{1}{1-2x} = \frac{2}{1-2x}$$

so that we can derive the Taylor series for $f(x)$ by differentiating the Taylor series for $\frac{1}{1-2x}$, computed in the previous exercise, and dividing by 2. Thus

$$\begin{aligned}\frac{1}{(1-2x)^2} &= \frac{1}{2} \cdot \frac{d}{dx} \left(\sum_{n=0}^{\infty} \frac{2^n}{5^{n+1}} (x+2)^n \right) \\ &= \frac{1}{2} \sum_{n=1}^{\infty} \frac{n 2^n}{5^{n+1}} (x+2)^{n-1} = \frac{2}{50} \sum_{n=1}^{\infty} \frac{n 2^{n-1}}{5^{n-1}} (x+2)^{n-1} \\ &= \frac{1}{25} \sum_{k=0}^{\infty} \frac{(k+1) 2^k}{5^k} (x+2)^k\end{aligned}$$

111. $f(x) = \ln \frac{x}{2}$, $c = 2$

SOLUTION Write

$$\ln \frac{x}{2} = \ln \left(\frac{(x-2)+2}{2} \right) = \ln \left(1 + \frac{x-2}{2} \right).$$

Substituting $\frac{x-2}{2}$ for x in the Maclaurin series for $\ln(1+x)$ yields

$$\ln \frac{x}{2} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \left(\frac{x-2}{2} \right)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} (x-2)^n}{n \cdot 2^n}.$$

This series is valid for $|x-2| < 2$.

112. $f(x) = x \ln \left(1 + \frac{x}{2} \right)$, $c = 0$

SOLUTION Substituting $\frac{x}{2}$ for x in the Maclaurin series for $\ln(1+x)$ yields

$$\ln \left(1 + \frac{x}{2} \right) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} \left(\frac{x}{2} \right)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n 2^n}.$$

Thus,

$$x \ln \left(1 + \frac{x}{2} \right) = x \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n 2^n} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^{n+1}}{n 2^n}.$$

In Exercises 113–116, find the first three terms of the Maclaurin series of $f(x)$ and use it to calculate $f^{(3)}(0)$.

113. $f(x) = (x^2 - x)e^{x^2}$

SOLUTION Substitute x^2 for x in the Maclaurin series for e^x to get

$$e^{x^2} = 1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6 + \dots$$

so that the Maclaurin series for $f(x)$ is

$$(x^2 - x)e^{x^2} = x^2 + x^4 + \frac{1}{2}x^6 + \dots - x - x^3 - \frac{1}{2}x^5 - \dots = -x + x^2 - x^3 + x^4 + \dots$$

The coefficient of x^3 is

$$\frac{f'''(0)}{3!} = -1$$

so that $f'''(0) = -6$.

114. $f(x) = \tan^{-1}(x^2 - x)$

SOLUTION Substitute $x^2 - x$ for x in the Maclaurin series for $\tan^{-1} x$ to get

$$\tan^{-1}(x^2 - x) = (x^2 - x) - \frac{1}{3}(x^2 - x)^3 + \dots = -x + x^2 + \frac{1}{3}x^3 + \dots$$

The coefficient of x^3 is

$$\frac{f'''(0)}{3!} = \frac{1}{3}$$

so that $f'''(0) = 3! \frac{1}{3} = 2$.

115. $f(x) = \frac{1}{1 + \tan x}$

SOLUTION Substitute $-\tan x$ in the Maclaurin series for $\frac{1}{1-x}$ to get

$$\frac{1}{1 + \tan x} = 1 - \tan x + (\tan x)^2 - (\tan x)^3 + \dots$$

We have not yet encountered the Maclaurin series for $\tan x$. We need only the terms up through x^3 , so compute

$$\tan'(x) = \sec^2 x \quad \tan''(x) = 2(\tan x) \sec^2 x \quad \tan'''(x) = 2(1 + \tan^2 x) \sec^2 x + 4(\tan^2 x) \sec^2 x$$

so that

$$\tan'(0) = 1 \quad \tan''(0) = 0 \quad \tan'''(0) = 2$$

Then the Maclaurin series for $\tan x$ is

$$\tan x = \tan 0 + \frac{\tan'(0)}{1!}x + \frac{\tan''(0)}{2!}x^2 + \frac{\tan'''(0)}{3!}x^3 + \cdots = x + \frac{1}{3}x^3 + \cdots$$

Substitute these into the series above to get

$$\begin{aligned} \frac{1}{1 + \tan x} &= 1 - \left(x + \frac{1}{3}x^3\right) + \left(x + \frac{1}{3}x^3\right)^2 - \left(x + \frac{1}{3}x^3\right)^3 + \cdots \\ &= 1 - x - \frac{1}{3}x^3 + x^2 - x^3 + \text{higher degree terms} \\ &= 1 - x + x^2 - \frac{4}{3}x^3 + \text{higher degree terms} \end{aligned}$$

The coefficient of x^3 is

$$\frac{f'''(0)}{3!} = -\frac{4}{3}$$

so that

$$f'''(0) = -6 \cdot \frac{4}{3} = -8$$

116. $f(x) = (\sin x)\sqrt{1+x}$

SOLUTION The binomial series for $\sqrt{1+x}$ is

$$\begin{aligned} \sqrt{1+x} &= (1+x)^{1/2} = \binom{1/2}{0} + \binom{1/2}{1}x + \binom{1/2}{2}x^2 + \binom{1/2}{3}x^3 + \cdots \\ &= 1 + \frac{1}{2}x + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)}{2}x^2 + \frac{\frac{1}{2}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}x^3 + \cdots \\ &= 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 + \cdots \end{aligned}$$

So, multiply the first few terms of the two Maclaurin series together:

$$\begin{aligned} (\sin x)\sqrt{1+x} &= \left(x - \frac{x^3}{6}\right) \left(1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3\right) \\ &= x + \frac{1}{2}x^2 - \frac{1}{8}x^3 - \frac{1}{6}x^3 + \text{higher degree terms} \\ &= x + \frac{1}{2}x^2 - \frac{7}{24}x^3 + \text{higher degree terms} \end{aligned}$$

The coefficient of x^3 is

$$\frac{f'''(0)}{3!} = -\frac{7}{24}$$

so that

$$f'''(0) = -6 \cdot \frac{7}{24} = -\frac{7}{4}$$

117. Calculate $\frac{\pi}{2} - \frac{\pi^3}{2^3 3!} + \frac{\pi^5}{2^5 5!} - \frac{\pi^7}{2^7 7!} + \cdots$.

SOLUTION We recognize that

$$\frac{\pi}{2} - \frac{\pi^3}{2^3 3!} + \frac{\pi^5}{2^5 5!} - \frac{\pi^7}{2^7 7!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{(\pi/2)^{2n+1}}{(2n+1)!}$$

is the Maclaurin series for $\sin x$ with x replaced by $\pi/2$. Therefore,

$$\frac{\pi}{2} - \frac{\pi^3}{2^3 3!} + \frac{\pi^5}{2^5 5!} - \frac{\pi^7}{2^7 7!} + \cdots = \sin \frac{\pi}{2} = 1.$$

118. Find the Maclaurin series of the function $F(x) = \int_0^x \frac{e^t - 1}{t} dt$.

SOLUTION Subtracting 1 from the Maclaurin series for e^t yields

$$e^t - 1 = \sum_{n=0}^{\infty} \frac{t^n}{n!} - 1 = 1 + \sum_{n=1}^{\infty} \frac{t^n}{n!} - 1 = \sum_{n=1}^{\infty} \frac{t^n}{n!}.$$

Thus,

$$\frac{e^t - 1}{t} = \frac{1}{t} \sum_{n=1}^{\infty} \frac{t^n}{n!} = \sum_{n=1}^{\infty} \frac{t^{n-1}}{n!}.$$

Finally, integrating term-by-term yields

$$\int_0^x \frac{e^t - 1}{t} dt = \int_0^x \sum_{n=1}^{\infty} \frac{t^{n-1}}{n!} dt = \sum_{n=1}^{\infty} \int_0^x \frac{t^{n-1}}{n!} dt = \sum_{n=1}^{\infty} \frac{x^n}{n! n}.$$